AUTHOR INDEX

Animal Feed Science and Technology, volumes 1–25

A

Abate, A., Kayongo-Male, H. and Karue, C.N.
Dry matter, protein, energy and fibre intake by dairy heifers grazing a Rhodes grass (*Chloris gayana*) pasture 6(1981)15

Abate, A. and Pfeffer, E.
Changes in nutrient intake and performance by goats fed coffee pulp-based diets followed by a commercial concentrate 14(1986)1

Abate, A., see Kategile, J.A. 20(1988)97

Abate, A.N. and Gomez, M.
Substitution of finger millet (*Eleusine coracana*) and bulrush millet (*Pennisetum typhoides*) for maize in broiler feeds 10(1983/1984)291

Abd El Ati, N., see Abel, Hj. 19(1988)111

Abd El Hafiz, G., see Abel, Hj. 19(1988)111

Abd El-Razek, M., see Abel, Hj. 19(1988)111

Abdel-Raheim, F.M., see El-Tayeb, A.E. 22 (1988)333

Abdullah, N. and Hutagalung, R.I.
Rumen fermentation, urease activity and performance of cattle given palm kernel cake-based diet 20(1988)79

Abdullah, N., see Ho, Y.W. 22(1988)161

Abel, Hj., Abd El Ati, N., Abd El Hafiz, G., Abd El-Razek, M., El-Din, T. and El Shazly, K.
Investigations on the nutritional value for ruminants of various fats 19(1988)111

Abidin, Z. and Kempton, T.J.
Effects of treatment of barley straw with anhydrous ammonia and supplementation with heat-treated protein meals on feed intake and liveweight performance of growing lambs 6(1981)145

Abouheif, M.A., Basmaeil, S., Metwally, H. and Masoud, S.
Chemical preparation of NaOH–keratin hydrolysate for improving the nutritive value of wheat straw 13(1985)215

Abrams, S.M.
Sources of error in predicting digestible dry matter from the acid-detergent fiber content of forages 21(1988)205

Acamovic, T., see D'Mello, J.P.F. 7(1982)247

Adams, K.L. and Jensen, A.H.
Effect of processing on the utilization by young pigs of the fat in soya beans and sunflower seeds 12(1984/1985)267

Adams, K.L. and Jensen, A.H.
Effect of dietary protein and fat levels on the utilization of the fat in sunflower seeds by the young pig 13(1985)159

Adams, K.L. and Jensen, A.H.
High-fat maize in diets for pigs and sows 17 (1987)201

Adamson, A.H., see Ibbotson, C.F. 10(1983/1984)223

Adamson, A.H., see Givens, D.I. 19(1988)173

Adamson, A.H., see Givens, D.I. 24(1989)27

Adamu, A.M., Russell, J.R. and Trenkle, A.
Effects of ruminal degradability of the protein supplement on the utilization of maize-stover silage by growing beef cattle 20(1988) 241

Adamu, A.M., Russell, J.R., McGilliard, A.D. and Trenkle, A.
Effects of added dietary urea on the utilization of maize stover silage by growing beef cattle 22(1988)227

Adebowale, E.A.
Organic waste ash as possible source of alkali for animal feed treatment 13(1985)237

Adegbola, A.A., see Smith, O.B. 13(1985)249

Adegbola, A.A., see Smith, O.B. 20(1988)189

Adeneye, J.A.
A note on the nutrient and mineral composition of *Leucaena leucocephala* in Western Nigeria 4(1979)221

Aderibigbe, A.O., see Razzaque, M.A. 14 (1986)265

Aderibigbe, A.O. and Church, D.C.
Evaluation of cardboard and dried poultry waste as feed ingredients for ruminants 18 (1987)209

Aduku, A.O., see Okoh, P.N. 7(1982)359

Aerts, J.V., see Carlier, L.A. 1(1976)607

Aerts, J.V., see Boucque, Ch.V. 1(1976)643

Aerts, J.V., de Brabander, D.L., Cottyn, B.G. and Buysse, F.X.
Comparison of laboratory methods for predicting the organic matter digestibility of forages 2(1977)337

Aerts, J.V., De Brabander, D.L., Cottyn, B.G., Buysse, F.X., Carlier, L.A. and Moermans, R.J.
Some remarks on the analytical procedure of Van Soest for the prediction of forage digestibility 3(1978)309

Aerts, J.V., see Cottyn, B.G. 11(1984)137

Aerts, J.V., De Boever, J.L., Cottyn, B.G., De Brabander, D.L. and Buysse, F.X.
Comparative digestibility of feedstuffs by sheep and cows 12(1984/1985)47

Aguilera, A., see Alcántara, E. 23(1989)323

Aguilera, J.F., see Molina, E. 9(1983)1

Aguilera, J.F., Molina, E. and Prieto, C.
Digestibility and energy value of sweet lupin seed (*Lupinus albus* var. multolupa) in pigs 12(1984/1985)171

Aguilera, J.F., Prieto, C., Fonolla, J. and Gil, F.
Protein and energy utilisation in rats of diets based on lupin seed (*Lupinus albus* var. multolupa) 15(1986)33

Aherne, F.X. and Lewis, A.J.
The nutritive value of tower rapeseed meal for swine 3(1978)235

Aherne, F.X., see Baidoo, S.K. 18(1987)37

Aherne, F.X., see Baidoo, S.K. 18(1987)45

Aherne, F.X., see Gibson, D.M. 19(1988)203

Aii, T. and Stobbs, the late T.H.
Solubility of the protein of tropical pasture species and the rate of its digestion in the rumen 5(1980)183

Aitken, W.M., see Wright, D.E. 2(1977)93

Akin, D.E.
Association of rumen fungi with various forage grasses 16(1986/1987)273
Akin, D.E., Rigsby, L.L., Theodorou, M.K. and Hartley, R.D.
Population changes of fibrolytic rumen bacteria in the presence of phenolic acids and plant extracts 19(1988)261
Akin, D.E.
Biological structure of lignocellulose and its degradation in the rumen 21(1988)295
Akin, D.E., Gordon, G.L.R. and Rigsby, L.L.
Comparative fiber degradation by mixed rumen fungi from Australian and U.S.A. cattle 23(1989)305
Akin, D.E., see Horn, G.W. 24(1989)201
Al-Athari, A.K. and Guenter, W.
Nutritional value of triticale (Carman) for broiler diets 22(1988)119
Al-Athari, A.K. and Guenter, W.
The effect of fat level and type on the utilization of triticale (cultivar Carman) by broiler chicks 22(1988)273
Al-Awadi, A., see Ilian, M.A. 19(1988)327
Al-Nasser, A., see Razzaque, M.A. 14(1986) 265
Alawa, J.P. and Owen, E.
The effect of milling and sodium hydroxide treatment on intake and digestibility of wheat straw by sheep 11(1984)149
Alawa, J.P. and Owen, E.
The effect of supplementing grass silage with sodium hydroxide-treated chopped or milled straw on intake and digestibility by wether sheep 12(1984/1985)215
Alawa, J.P., Fishwick, G. and Hemingway, R.G.
Fresh and dried brewer's grains as protein supplements to barley straw diets given to pregnant beef cows 19(1988)33
Alcántara, E., Aguilera, A., Elliott, R. and Shimada, A.
Fermentation and utilization by lambs of sugarcane harvested fresh and ensiled with and without NaOH. 4. Ruminal kinetics 23 (1989)323
Alexander, J.C., see Gregory, K.F. 2(1977)7
Alexander, J.C., see Santos, J. 8(1983)313
Alhassan, W.S., Bello, S.A. and Obilana, A.B.
Yield and potential feeding value of straws of grain sorghum cultivars developed in Nigeria 17(1987)285
Ali, O.M., see Mohammed, T.A. 22(1988)131
Alibes, X., Muñoz, F. and Rodriguez, J.
Feeding value of apple pomace silage for sheep 11(1984)189
Alibés, X., Maestre, M.R., Muñoz, F., Combellas, J. and Rodriguez, J.
Nutritive value of almond hulls for sheep 8 (1983)63
Alibés, X., Muñoz, F. and Faci, R.
Anhydrous ammonia-treated cereal straw for animal feeding. Some results from the Mediterranean area 10(1983/1984)239
Aliu, Y.O., see Kumaresan, A. 11(1984)45
Allan, S.J., see Partridge, G.G. 13(1985)299
Allee, G.L., see Pollmann, D.S. 8(1983)99
Alli, I., Baker, B.E. and Garcia, G.
Studies on the fermentation of chopped sugarcane 7(1982)411
Alli, I., Fairbairn, R., Baker, B.E. and Garcia, G.
The effects of ammonia on the fermentation of chopped sugarcane 9(1983)291
Allison, M.J.
Characterization of the flora of the large bowel of pigs: a status report 23(1989)79
Alumot, E. and Mandel, E.
Long term feeding of concentrate fumigated with carbon tetrachloride to sheep 2(1977) 77
Aluyi, H.S.A., see Aregheore, E.M. 20(1988) 233
Alvira, P., see Rebolé, A. 16(1986/1987)89
Alvira, P., see Rebolé, A. 22(1988)173
Alzueta, C. and Gonzalez, G.
Yield, composition and nutritive value of a leaf protein concentrate obtained by fractionating the shelled aerial part of *Pisum savitum* L. 8(1983)155
Alzueta, C., see González, G. 20(1988)177
Åman, P., see Graham, H. 10(1983/1984)199
Åman, P., see Graham, H. 12(1984/1985)195
Åman, P., see Hesselman, K. 15(1986)83
Åman, P. and Graham, H.
Whole-crop peas. I. Changes in botanical and chemical composition and rumen in vitro degradability during maturation 17 (1987)15
Åman, P., see Graham, H. 17(1987)33
Åman, P., see Pettersson, D. 17(1987)145
Åman, P., see Nordkvist, E. 17(1987)295
Åman, P., see Pettersson, D. 20(1988)313
Åman, P., see Theander, O. 23(1989)205
Åman, P., see Narasimhalu, P. 24(1989)159
Amaning-Kwarteng, K., Kellaway, R.C., Spragg, J.C. and Kirby, A.C.
Relative intakes, digestibility and bacterial protein synthesis by sheep and cattle fed high-roughage diets 16(1986/1987)75
Ammerman, C.B., see Potter, S.G. 12(1984/ 1985)275
Anderson, D.H., see Henderson, A.R. 7(1982) 303
Anderson, D.M., see Bowland, J.P. 12(1984/ 1985)75
Anderson, P., see Zervas, G. 21(1988)23
Anderson, R.
Effect of prolonged wilting in poor conditions on the fermentation quality, metabolisability and net energy value of silage given to sheep 12(1984/1985)109
Andries, J.I., Buysse, F.X., de Brabander, D.L. and Cottyn, B.G.
Isoacids in ruminant nutrition: their role in ruminal and intermediary metabolism and

possible influences on performances – a review 18(1987)169
Andries, J.I., see De Boever, J.L. 19(1988)247
Andrieu, J.
Agronomic factors affecting the growth and composition of the maize plant 1(1976)263
Andrieu, J.
Factors influencing the composition and nutritive value of ensiled whole-crop maize 1 (1976)381
Angemar, Y., see Holzer, Z. 12(1984/1985)253
Anglesea, J.D. and Jackson, A.J.
Thiaminase activity in fish silage and moist fish feed 13(1985)39
Anonymous
Requirements for future research 1(1976) 555
Antongiovanni, M., Giorgetti, A. and Franci, O.
Amino acid composition of two varieties of sorghum grain 5(1980)169
Anwar, A., see Mohamed, K. 11(1984)199
Aregheore, E.M., Job, T.A. and Aluyi, H.S.A.
The maize replacement value of cassava flour in rations for growing ewe lambs 20 (1988)233
Argamenteria, A., see Caja, G. 2(1977)267
Arieli, A. and Werner, D.
A comparison between fermentation heat of forages and organic matter digestibility determined by in vitro incubation with rumen fluid 23(1989)333
Armstrong, D.G., see McMeniman, N.P. 2 (1977)255
Armstrong, D.G., see Rooke, J.A. 13(1985)255
Armstrong, D.G., see Rooke, J.A. 13(1985)269
Armstrong, D.G., see Lee, N.-H. 15(1986)301
Arora, S.K., see Das, B. 4(1979)17
Arp, S.C., see Schaefer, D.M. 25(1989)23
Asahida, Y., see Sekine, J. 25(1989)207
Ash, A.J., see McMeniman, N.P. 19(1988)43
Ashbell, G., see Levy, D. 16(1986/1987)63
Ashbell, G., see Weinberg, Z.G. 20(1988)335
Ashes, J.R. and Peck, N.J.
A simple device for dehulling seeds and grain 3(1978)109
Ashes, J.R., see Farrell, D.J. 9(1983)99
Asplund, J.M., see Bowman, J.G.P. 20(1988) 19
Asplund, J.M., see Bowman, J.G.P. 20(1988) 33
Atinkpahoun, H., see Szylit, O. 3(1978)73
Atkinson, T., see Livingstone, R.M. 4(1979) 295
Atkinson, T., see Livingstone, R.M. 5(1980)69
Atuahene, C.C., Donkoh, A. and Swatson, H.
Oil palm slurry (OPS) as a partial replacement for maize in the diets of broiler chickens 17(1987)157
Atuahene, C.C., see Donkoh, A. 22(1988)139
Aufrere, J. and Michalet-Doreau, B.
Comparison of methods for predicting digestibility of feeds 20(1988)203
Auld, D.L., see Thomas, V.M. 11(1984)269
Aumaitre, A., see Le Dividich, J. 3(1978)15
Aumaitre, A., see Perez, J.M. 4(1979)109
Aumaitre, A., Bourdon, D., Peiniau, J. and Bengala Freire, J.
Effect of graded levels of raw and processed rapeseed on feed digestibility and nutrient utilization in young pigs 24(1989)275
Awadel-Kareim, A.M., see El-Tayeb, A.E. 22 (1988)333
Ayyaluswami, P., see Thanu, K. 9(1983)263

B

Babatunde, G.M., Fetuga, B.L., Odebunmi, J.O. and Oduye, O.O.
The effects of air dried poultry manure on performance, organ weights and haematological measurements of weaner pigs reared in the tropics 4(1979)23
Babatunde, G.M. and Pond, W.G.
Nutritive value of Nigerian rubber seed (*Hevea brasiliensis*) meal and oil. III. Performance characteristics, relative organ weights, hematocrit and plasma metabolites of growing female rats fed corn diets containing rubber seed meal, soya bean meal or casein 20(1988)125
Bačvanski, S.
Maize grain or ears in concentrate diets for young fattening bulls 1(1976)393
Badir, N.A., see El Khidir, O.A. 9(1983)301
Badve, V.C., see Rangnekar, D.V. 7(1982)61
Baidoo, S.K., Aherne, F.X., Mitaru, B.N. and Blair, R.
Canola meal as a protein supplement for growing–finishing pigs 18(1987)37
Baidoo, S.K., Mitaru, B.N., Aherne, F.X. and Blair, R.
The nutritive value of canola meal for early-weaned pigs 18(1987)45
Baird, B.A., see Livingstone, R.M. 4(1979)295
Baird, B.A., see Livingstone, R.M. 5(1980)69
Baker, B.E., see Alli, I. 7(1982)411
Baker, B.E., see Alli, I. 9(1983)291
Baker, E.C., see Lynch, G.L. 14(1986)127
Bakshi, M.P.S., see Langar, P.N. 13(1985)103
Balagopal, R., see Natanam, R. 25(1989)201
Baldwin, A., see O'Shea, J. 15(1986)13
Ballance, G.M., see Campbell, G.L. 16(1986/ 1987)243
Balogun, O.O. and Fetuga, B.L.
Lysine requirements of imported European pigs in the humid lowland tropics from 23 to 62 kg liveweight 7(1982)233
Balogun, O.O. and Fetuga, B.L.A.
Lysine requirement of weanling European pigs fed on yellow maize and groundnut cake diets in the humid lowland tropics 8(1983) 197
Bandyk, C.A., see Pollmann, D.S. 11(1984) 261
Bar-Tsur, A., see Levy, D. 16(1986/1987)63

Baran, M., Boďa, K. and Siroka, P.
The effect of monensin on rumen fermentation in sheep fed on all-roughage and barley/roughage diets 15(1986)7
Barber, R.S., Braude, R. and Mitchell, K.G.
The value of 'Pekilo protein' for growing pigs 2(1977)161
Barber, R.S., Braude, R., Mitchell, K.G. and Pittman, R.J.
The nutritive value of liquid whey, either sour or sweet, when given in restricted amounts to the growing pig 3(1978)163
Barber, R.S., Braude, R., Hosking, Z.D. and Mitchell, K.G.
Olaquindox as performance-promoting feed additive for growing pigs 4(1979)117
Barber, R.S., Braude, R., Mitchell, K.G., Partridge, I.G. and Pittman, R.J.
Value of lucerne juice and grass juice as sources of protein for the growing pig 4 (1979)233
Barber, R.S., Braude, R., Mitchell, K.G., Partridge, I.G. and Pittman, R.J.
Value of freshly produced lucerne juice as a source of supplemental protein for the growing pig 5(1980)215
Barber, R.S., Braude, R., Mitchell, K.G. and Partridge, I.G.
Lucerne juice as a protein supplement for growing pigs: effects of mineral content of the diet and of the water supply 6(1981)35
Barber, W.P., see Givens, D.I. 16(1986/1987) 311
Barge, M.T., see Pace, V. 11(1984)125
Barker, D.J., see May, P.J. 12(1984/1985)57
Barnes, B.J. and Ørskov, E.R.
Utilization of alkali-treated grain. 2. Utilization by steers of diets based on hay or straw and mixed with either NaOH-treated or rolled barley 6(1981)347
Barnes, B.J., see Ørskov, E.R. 6(1981)355
Barnes, D.A., see Brown, D.L. 18(1987)247
Barnes, R.J.
Near infra-red spectra of ammonia-treated straw and of isolated cell walls 21(1988)209
Barnett, C.W. and Batterham, E.S.
Lupinus angustifolius cv. Unicrop as a protein- and energy-source for weaner pigs 6(1981)27
Barnett, R.D., see Myer, R.O. 22(1988)217
Barro, C., see González, G. 20(1988)177
Barry, J.L., see Cherbut, C. 20(1988)327
Barry, J.L., see Champ, M. 23(1989)195
Barry, P., see Grenet, E. 19(1988)25
Bartling, R.R., see Umunna, N.N. 12(1984/1985)151
Bartling, R.R., see Umunna, N.N. 14(1986)161
Barton, F.E., see Coelho, M. 20(1988)219
Barton, G.M., see Hunt, J.R. 3(1978)63
Barton II, F.E., Wolsink, J.H. and Vedder, H.M.
Near infrared reflectance spectroscopy of untreated and ammoniated barley straw 15 (1986)189
Barton II, F.E.
Chemistry of lignocellulose: methods of analysis and consequences of structure 21 (1988)279
Bartsch, B.D., see Valentine, S.C. 16(1986/1987)143
Bartsch, B.D., see Valentine, S.C. 16(1986/1987)261
Basmaeil, S., see Abouheif, M.A. 13(1985)215
Bass, J.M., Parkins, J.J. and Fishwick, G.
The effect of calcium hydroxide treatment on the digestibility of chopped oat straw supplemented with a solution containing urea, calcium, phosphorus, sodium, trace elements and vitamins 7(1982)93
Batchelder, D.G., see Horn, G.W. 8(1983)35
Batterham, E.S., see Barnett, C.W. 6(1981)27
Batterham, E.S., Gorman, T.B.S. and Chvojka, R.
Nutritional value and mercury content of fish silage for growing pigs 9(1983)169
Batterham, E.S. and Watson, C.
Tryptophan content of feeds, limitations in diets and requirement for growing pigs 13 (1985)171
Baucells, M.D., see Gasa, J. 25(1989)67
Bazylak, G., see Maslowska, J. 13(1985)227
Beames, R.M., see Pringle, D. 9(1983)89
Beames, R.M., Tait, R.M., Pringle, D.B. and Owen, B.D.
The nutritive value of barley for monogastric animals and for ruminants as affected by artificial drying and alkali and anaerobic storage 15(1986)285
Beard, J., see Clark, J. 2(1977)153
Beaumatin, P., see Durand, M. 21(1988)197
Becker, H.N., see Potter, S.G. 12(1984/1985) 275
Becker, K., see Ohlde, G. 7(1982)191
Beever, D.E.
The digestion and metabolism of the energy and protein of maize 1(1976)455
Behnke, K.C., see Skoch, E.R. 6(1981)83
Behnke, K.C., see Pollmann, D.S. 8(1983)99
Bell, J.M., see Thacker, P.A. 19(1988)191
Bell, J.M. and Keith, M.O.
Factors effecting the digestibility by pigs of energy and protein in wheat, barley and sorghum diets supplemented with canola meal 24(1989)253
Bell, S.L., see Neill, A.R. 9(1983)71
Bell, S.L., see Rooke, J.A. 13(1985)269
Bello, S.A., see Alhassan, W.S. 17(1987)285
Belyavin, C.G.
Effects of the addition of silica to the diet of the laying hen, with particular reference to egg-shell quality 6(1981)157
Belyea, R.L., Martz, F.A., Madhisetty, M. and Davis, T.P.
Intake, digestibility and energy utilization of corrugated paper diets 17(1987)57
Ben-Ghedalia, D. and Miron, J.
The response of wheat straw varieties to mild sulphur dioxide treatment 10(1983/

1984)269
Ben-Ghedalia, D. and Rubinstein, A.
The response of screened manure fibre to ozone and sodium hydroxide treatments 15 (1986)47
Ben-Ghedalia, D. and Solomon, R.
The effect of dietary barley on carbohydrate digestibility of sulphur dioxide-treated wheat straw by sheep 18(1987)55
Ben-Ghedalia, D., Miron, J., Est, Y. and Yosef, E.
SO_2 treatment for converting straw into a concentrate-like feed: a growth study with lambs 19(1988)219
Ben-Ghedalia, D. and Solomon, R.
Amino acids flow to and absorption from the small intestine of sheep fed barley and sulfur dioxide-treated straw at different ratios 22(1988)147
Ben-Ghedalia, D. and Yosef, E.
SO_2-treated straw as a silage additive: fermentation data on lucerne with particular reference to protein degradation 22(1988) 247
Ben-Ghedalia, D. and Yosef, E.
SO_2-treated straw as a silage additive: the participation of soluble and cell wall monosaccharide residues in lucerne silage fermentation 22(1988)255
Ben-Ghedalia, D., Yosef, E., Miron, J. and Est, Y.
The effects of starch and pectin-rich diets on quantitative aspects of digestion in sheep 24 (1989)289
Benevenga, N.J., see Cieslak, D.G. 14(1986) 171
Bengala Freire, J., see Aumaitre, A. 24(1989) 275
Berger, L.L., see Mora, M.I. 9(1983)205
Berger, L.L., see Lynch, G.L. 14(1986)127
Berger, L.L., see Lewis, S.M. 17(1987)179
Berkovitz, S., see Levy, D. 16(1986/1987)63
Bertrand, D., see Sauvant, D. 13(1985)7
Bertrand, D., see Robert, P. 16(1986/1987)215
Berzins, R., see Ha, J.K. 14(1986)117
Bhargava, P.K., Ørskov, E.R. and Walli, T.K.
Effect of soaking, ensilage and hydrogen peroxide treatment of barley straw on rumen degradability 22(1988)295
Bickel, H., see Wyss, U. 20(1988)325
Bienfait, J.M.. see Istasse, L. 25(1989)1
Binder, S.F., see Skoch, E.R. 6(1981)83
Biston, L., see Istasse, L. 25(1989)1
Blair, R., see Baidoo, S.K. 18(1987)37
Blair, R., see Baidoo, S.K. 18(1987)45
Blanchart, G., see Laurent, F. 13(1985)131
Blanco, J.L.
Selection objectives for maximum yields of feed with hybrid maize 1(1976)245
Bland, B., see Panigrahi, S. 22(1988)285
Boďa, K., see Baran, M. 15(1986)7
Boekholt, H.A., see Van Es, A.J.S. 1(1976)481
Boer, H., see Tamminga, S. 25(1989)89
Bokori, J., see Hegedüs, M. 9(1983)79
Boling, J.A., see Pendlum, L.C. 2(1977)225
Bolsen, K.K., Tetlow, R.M. and Wilson, R.F.
The effect of calcium and sodium hydroxides and of sodium acrylate on the fermentation and digestibility in vitro of ensiled whole-crop wheat and barley harvested at different stages of maturity 9(1983)37
Bolsen, K.K., see Tetlow, R.M. 12(1984/1985) 91
Bolsen, K.K., see Hinds, M.A. 12(1984/1985) 205
Bolt, M.J., see Slabbert, N. 21(1988)31
Borgida, L.P., see Szylit, O. 3(1978)73
Borhami, B.E.A. and Sundstøl, F.
Studies on ammonia-treated straw. I. The effects of type and level of ammonia, moisture content and treatment time on the digestibility in vitro and enzyme soluble organic matter of oat straw 7(1982)45
Borhami, B.E.A., Sundstøl, F. and Garmo, T.H.
Studies on ammonia-treated straw. II. Fixation of ammonia in treated straw by spraying with acids 7(1982)53
Bothast, R.J., see Montgomery, R.R. 5(1980) 337
Boucque, Ch.V., see Cottyn, B.G. 1(1976)167
Boucque, Ch.V., Cottyn, B.G. and Buysse, F.X.
Maize silage with or without NPN, dehydrated whole-crop maize pellets or high moisture maize grain for finishing bulls 1 (1976)347
Boucque, Ch.V., Cottyn, B.G., Aerts, J.V. and Buysse, F.X.
Dried suger beet pulp as a high energy feed for beef cattle 1(1976)643
Boucqué, Ch.V., see Fiems, L.O. 8(1983)237
Boucqué, Ch.V., see Fiems, L.O. 16(1986/ 1987)287
Bouqué, Ch.V., Fiems, L.O., Cottyn, B.G., Casteels, M. and Buysse, F.X.
Monensin-sodium as a performance-promoting additive for fattening bulls and its impact on carcass and meat quality characteristics 7(1982)401
Bourdon, D., see Aumaitre, A. 24(1989)275
Bowland, J.P., Anderson, D.M. and Thacker, P.A.
Chemical composition and nutritive value of Canadian winter wheats (*Triticum aestivum* L.) given to laboratory rats 12(1984/ 1985)75
Bowman, D.K., see Koeln, L.L. 11(1984)181
Bowman, J.G.P. and Asplund, J.M.
Evaluation of mixed lucerne and caucasian bluestem hay diets fed to sheep 20(1988)19
Bowman, J.G.P. and Asplund, J.M.
Nitrogen utilization, ruminal fermentation and abomasal nitrogen flow in sheep fed caucasian bluestem hay supplemented with lucerne or urea 20(1988)33
Boza, J., see Molina, E. 9(1983)1

Brahmakshatriva, R.D., see Singh, K.S. 6 (1981)63
Braude, R., see Barber, R.S. 2(1977)161
Braude, R., see Barber, R.S. 3(1978)163
Braude, R., see Barber, R.S. 4(1979)117
Braude, R., see Barber, R.S. 4(1979)233
Braude, R., see Barber, R.S. 5(1980)215
Braude, R., see Barber, R.S. 6(1981)35
Bremel, D.H., see Teh, T.H. 17(1987)257
Brethour, J., see Hinds, M.A. 12(1984/1985) 205
Brett, P.A., see Rooke, J.A. 13(1985)255
Brewer, A., see Williams, P.E.V. 11(1984)103
Brewer, A., see Williams, P.E.V. 11(1984)115
Brice, R.E., see Morrison, I.M. 10(1983/1984) 229
Brice, R.E., see Robinson, P.H. 18(1987)75
Brigstocke, T.D.A., see Wilson, P.N. 6(1981)1
Brophy, P.O., see Caffrey, P.J. 19(1988)231
Broster, W.H., see Smith, T. 10(1983/1984) 189
Broster, W.H., see Smith, T. 14(1986)29
Brotz, P.G., see Schaefer, D.M. 25(1989)23
Brown, D.L. and Chavalimu, E.
Effects of ensiling or drying on five forage species in western Kenya: *Zea mays* (maize stover), *Pennisetum purpureum* (Pakistan napier grass), *Pennisetum* sp. (bana grass), *Impomea batata* (sweet potato vines) and *Cajanus cajan* (pigeon pea leaves) 13(1985) 1
Brown, D.L., Barnes, D.A., Rezende, S.A. and Klasing, K.C.
Yield, composition and feeding value of irrigated *Sesbania sesban* var. *Nubia* leaves harvested at latitude 38°N during a Mediterranean summer 18(1987)247
Brown, D.L., see Shqueir, A.A. 25(1989)137
Brunnen, J.M., see Givens, D.I. 18(1987)83
Brunnen, J.M., see Clark, P. 18(1987)225
Bruyneel, B., see Vandevoorde, L. 21(1988)99
Bryant, K.L., see Harper, A.F. 8(1983)69
Buchanan-Smith, J.G., see Oji, U.I. 4(1979) 187
Bunting, L.D., McCarthy, D.B., Richardson, C.R. and Tock, R.W.
Pilot plant production of sulphur dioxide-treated mesquite wood. II. Digestibility and feedlot performance by lambs 22(1988)61
Burke, A.B., see Waldroup, P.W. 3(1978)329
Bussell, W.D., see Waldroup, P.W. 3(1978)329
Butler, E.J., see Fenwick, G.R. 6(1981)421
Butler, E.J., see Pearson, A.W. 8(1983)119
Butler, K.L., see Minson, D.J. 9(1983)221
Butterworth, M.H., see Mosi, A.K. 12(1984/ 1985)241
Butterworth, M.H. and Mosi, A.
The intake and digestibility by sheep of oat straw and maize stover offered with different levels of noug (*Guizotia abyssinica*) meal 16 (1986/1987)99
Buxton, D.R., see Russell, J.R. 13(1985)69
Buysse, F.X., see Cottyn, B.G. 1(1976)167
Buysse, F.X., see Boucque, Ch.V. 1(1976)347
Buysse, F.X., see Boucque, Ch.V. 1(1976)643
Buysse, F.X., see Aerts, J.V. 2(1977)337
Buysse, F.X., see Aerts, J.V. 3(1978)309
Buysse, F.X., see Bouqué, Ch.V. 7(1982)401
Buysse, F.X., see Fiems, L.O. 8(1983)237
Buysse, F.X., see Cottyn, B.G. 11(1984)137
Buysse, F.X., see Aerts, J.V. 12(1984/1985)47
Buysse, F.X., see De Boever, J.L. 14(1986)203
Buysse, F.X., see Fiems, L.O. 16(1986/1987) 287
Buysse, F.X., see Andries, J.I. 18(1987)169
Buysse, F.X., see De Boever, J.L. 19(1988)247

C

Caballero, R., see Treviño, J. 16(1986/1987) 305
Caffrey, P.J., Miller, C., Brophy, P.O. and Kelleher, D.L.
The effects of method of processing of starters, tallow inclusion and roughage supplementation on the performance of early-weaned calves 19(1988)231
Caja, G., Galvez, J.F., Argamenteria, A. and Ciria, J.
Inhibition of ruminal deamination in vitro by formaldehyde treatment of sunflower-seed, soya bean and fish meals: response curves to protective treatment 2(1977)267
Campbell, D.R., see Noland, P.R. 2(1977)219
Campbell, D.R., see Sharp, R.N. 2(1977)327
Campbell, D.R., see Noland, P.R. 5(1980)51
Campbell, G.L., Sosulski, F.W., Classen, H.L. and Ballance, G.M.
Nutritive value of irradiated and β-glucanase-treated wild oat groats (*Avena fatua* L.) for broiler chickens 16(1986/1987)243
Campbell, G.L., see Thacker, P.A. 19(1988) 191
Campbell, G.L., see Edney, M.J. 25(1989)193
Campbell, R.G., see King, R.H. 3(1978)191
Campher, J.P., see Slabbert, N. 21(1988)31
Campling, R.C., see Jewell, S.N. 14(1986)81
Capper, A.L., see Edwards, S.A. 15(1986)129
Capper, A.L.
Fungal contamination of hydroponic forage 20(1988)163
Capper, B.S., see Machin, D.H. 11(1984)247
Capper, B.S.
Genetic variation in the feeding value of cereal straw 21(1988)127
Carlaw, P.M., see Panigrahi, S. 22(1988)285
Carlier, L.A., Cottyn, B.G. and Aerts, J.V.
Apparent and true digestibility of the Weende components, cell content and cell wall of ryegrass 1(1976)607
Carlier, L.A., see Aerts, J.V. 3(1978)309
Carlos, G., see Zervas, G. 21(1988)23
Carlsson, R., Hanczakowski, P. and Kaptur, T.
The quality of the green fraction of leaf pro-

tein concentrate from *Chenopodium quinoa* Willd. grown at different levels of fertilizer nitrogen 11(1984)239
Carr, J.R., see Pearson, G. 1(1976)631
Carr, J.R., see Pearson, G. 2(1977)49
Carré, B., see Lacassagne, L. 20(1988)59
Carter, P.M., see Wood, J.F. 9(1983)307
Carter, P.M., see Machin, D.H. 11(1984)247
Carter, R.R., see Leibholz, J. 15(1986)141
Casper, H.H., see Sanderson, M.A. 14(1986) 221
Castaing, J., see Gatel, F. 20(1988)145
Casteels, M., see Bouqué, Ch.V. 7(1982)401
Castrillo, C., see Gasa, J. 25(1989)67
Catalano, A., see Russo, V. 1(1976)25
Cavani, C., see Manfredini, M. 5(1980)233
Cayley, G.R., see Lord, K.A. 6(1981)73
Cayley, G.R., see Lacey, J. 6(1981)323
Cayley, G.R., see Lord, K.A. 7(1982)287
Cayley, G.R., see Lacey, J. 8(1983)283
Centeno, C., see Treviño, J. 16(1986/1987)305
Cervera, C., Fernandez-Carmona, J. and Marti, J.
Effect of urea on the ensiling process of orange pulp 12(1984/1985)233
Chahal, D.S., see Moo-Young, M. 4(1979)199
Chahal, D.S., see Moo-Young, M. 5(1980)175
Champ, M., Barry, J.L., Hoebler, C. and Delort-Laval, J.
Digestion and fermentation pattern of various dietary fiber sources in the rat 23(1989) 195
Chan, C., see Farrell, D.J. 9(1983)49
Chandra, S., Prasad, D.A. and Krishna, N.
Effect of sodium hydroxide treatment and/or extrusion cooking on the nutritive value of peanut hulls 12(1984/1985)187
Chandrasekaran, D., see Subramanian, P.R. 20(1988)343
Chandrasekaran, D., Kadirvel, R. and Viswanathan, K.
Nutritive value of pungam (*Pongamia glabra* Vent) cake for sheep 22(1988)321
Chandrasekaran, D., see Viswanathan, K. 22 (1988)327
Charmley, E. and Greenhalgh, J.F.D.
Nutritive value of three cultivars of triticale for sheep, pigs and poultry 18(1987)19
Chase Jr., C.C. and Hibberd, C.A.
Effect of level and frequency of maize supplementation on the utilization of low-quality grass hay by beef cows 24(1989)129
Chavalimu, E., see Brown, D.L. 13(1985)1
Chavez, E.R., Touchburn, S.P. and Moo-Young, M.
Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. I. Composition and nutritive evaluation 22(1988)1
Chavez, E.R., Touchburn, S.P. and Moo-Young, M.
Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. II. Effect of method of drying and response to amino acid supplementation 22(1988)13
Chavez, E.R., Touchburn, S.P. and Moo-Young, M.
Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. III. Nutritive value for chicks and piglets 22(1988)23
Cheng, K.-J., Stewart, C.S., Dinsdale, D. and Costerton, J.W.
Electron microscopy of bacteria involved in the digestion of plant cell walls 10(1983/1984)93
Cherbut, C., Barry, J.L., Wyers, M. and Delort-Laval, J.
Effect of the nature of dietary fibre on transit time and faecal excretion in the growing pig 20(1988)327
Cherney, J.H., Johnson, K.D., Tuite, J. and Volenec, J.J.
Microfloral and compositional changes in alfalfa hay treated with sodium diacetate and stored at different moisture contents 17 (1987)45
Chesson, A., see Gordon, A.H. 8(1983)147
Chesson, A.
Lignin-polysaccharide complexes of the plant cell wall and their effect on microbial degradation in the rumen 21(1988)219
Chiericato, G.M., see Parigi-Bini, R. 1(1976) 301
Choice, A., see Farrell, D.J. 9(1983)99
Christensen, D.A., see Horton, G.M.J. 7(1982) 1
Christensen, D.A., see Mann, M.E. 21(1988) 57
Christison, G.I., see Mandell, I.B. 20(1988)111
Christon, R., see Le Dividich, J. 3(1978)15
Church, D.C., see Razzaque, M.A. 14(1986) 265
Church, D.C., see Aderibigbe, A.O. 18(1987) 209
Chvojka, R., see Batterham, E.S. 9(1983)169
Cieslak, D.G., Benevenga, N.J. and Grummer, R.H.
The evaluation of fresh sweet liquid whey as a protein supplement to maize for growing pigs 14(1986)171
Ciria, J., see Caja, G. 2(1977)267
Clark, C.F.S., see Ørskov, E.R. 7(1982)135
Clark, J. and Beard, J.
Prediction of the digestibility of ruminant feeds from their solubility in enzyme solutions 2(1977)153
Clark, P., Givens, D.I. and Brunnen, J.M.
The chemical composition, digestibility and energy value of fodder-beet roots 18(1987) 225
Classen, H.L., see Campbell, G.L. 16(1986/1987)243
Classen, H.L., see Thacker, P.A. 19(1988)191
Classen, H.L., see Edney, M.J. 25(1989)193
Cobby, J.M., see Mason, V.C. 19(1988)159
Cobby, J.M., see Givens, D.I. 19(1988)173
Cobic, T.
Effect of weight at slaughter on the efficiency of fattening young cattle 1(1976)505

Cockburn, J.E., see Mason, V.C. 24(1989)313
Cockburn, J.H.S., see Morgan, C.A. 11(1984) 11
Coelho, M., Hembry, F.G., Barton, F.E. and Saxton, A.M.
A comparison of microbial, enzymatic, chemical and near-infrared reflectance spectroscopy methods in forage evaluation 20 (1988)219
Cohen, O., see Silanikove, N. 20(1988)87
Cohen, R.D.H., see Mann, M.E. 21(1988)57
Cole, D.J.A., see Green, S. 21(1988)43
Collins, M.
Composition and fibre digestion in morphological components of an alfalfa–timothy sward 19(1988)135
Combellas, J., see Alibés, X. 8(1983)63
Combs, G.E., see Myer, R.O. 22(1988)217
Compte, X., see Valdivie, M. 7(1982)185
Connor, J.K., see Moir, K.W. 2(1977)197
Conrad, J.H., see Paz, E.A. 11(1984)75
Cook, D.K., see Schaefer, D.M. 25(1989)23
Cook, J.E., see Woolford, M.K. 3(1978)89
Cook, J.E., see Woolford, M.K. 8(1983)107
Cook, J.E., see Mason, V.C. 24(1989)299
Cook, J.E., see Mason, V.C. 24(1989)313
Cooper, E.M., see Mason, V.C. 24(1989)299
Cooper, E.M., see Mason, V.C. 24(1989)313
Coppock, C.E., Lanham, J.K. and Horner, J.I.
A review of the nutritive value and utilization of whole cottonseed, cottonseed meal and associated by-products by dairy cattle 18(1987)89
Corcuera, P., see Penning, P.D. 5(1980)321
Cornelissen, J.P., see Grimbergen, A.H.M. 13 (1985)117
Cornell, J.A., see Myer, R.O. 22(1988)217
Costerton, J.W., see Cheng, K.-J. 10(1983/ 1984)93
Cote, M., see Sibbald, I.R. 8(1983)191
Cottyn, B.G., Boucque, Ch.V. and Buysse, F.X.
Survey of the use of maize for livestock feeding in Belgium 1(1976)167
Cottyn, B.G., see Boucque, Ch.V. 1(1976)347
Cottyn, B.G., see Carlier, L.A. 1(1976)607
Cottyn, B.G., see Boucque, Ch.V. 1(1976)643
Cottyn, B.G., see Aerts, J.V. 2(1977)337
Cottyn, B.G., see Aerts, J.V. 3(1978)309
Cottyn, B.G., see Bouqué, Ch.V. 7(1982)401
Cottyn, B.G., see Fiems, L.O. 8(1983)237
Cottyn, B.G., Aerts, J.V., Vanacker, J.M., Moermans, R.J. and Buysse, F.X.
The prediction of the net energy value of compound feedstuffs for dairy cattle 11 (1984)137
Cottyn, B.G., see Aerts, J.V. 12(1984/1985) 47
Cottyn, B.G., see De Boever, J.L. 14(1986)203
Cottyn, B.G., see Fiems, L.O. 16(1986/1987) 287
Cottyn, B.G., see Andries, J.I. 18(1987)169
Cottyn, B.G., see De Boever, J.L. 19(1988)247
Cottyn, B.G. and de Boever, J.L.
Upgrading of straw by ammoniation 21 (1988)287
Courtin, M.G. and Spoelstra, S.F.
Counteracting structure loss in pressed sugar beet pulp silage 24(1989)97
Cowan, P.J. and Michie, W.
Turkey feeding regimes with different ages of introduction of a choice of foods 4(1979) 125
Coxworth, E.C., see Kernan, J.A. 6(1981)257
Coxworth, E.C., see Kernan, J.A. 11(1984)301
Coxworth, E.C., see Mandell, I.B. 20(1988)111
Crickenberger, R.G., see Johnson, T.O. 12 (1984/1985)65
Crickenberger, R.G., see Ortigues, I. 24(1989) 13
Crofts, F.C., see Kellaway, R.C. 3(1978)201
Crofts, R.M.J., see Livingstone, R.M. 4(1979) 295
Croke, B., see Lahore, R. 3(1978)1
Crooks, P., see Morgan, C.A. 17(1987)81
Cross, D.L., see Grotheer, M.D. 14(1986)55
Crowle, W.L., see Kernan, J.A. 11(1984)301
Čuperlović, M., see Jovanović, M. 2(1977)351
Curl, C.L., see Pearson, A.W. 8(1983)119
Curtis, R.F., see Fenwick, G.R. 5(1980)255
Cuthbert, N.H., see Wilson, P.N. 6(1981)1

D

da Costa, J.S.
Survey of the use of maize for livestock feeding in Portugal 1(1976)137
Damron, B.L., see Paz, E.A. 11(1984)75
Daniels, L.B., Smith, M.J., Stallcup, O.T. and Rakes, J.M.
Nutritive value of ensiled broiler litter for cattle 8(1983)19
Das, B., Arora, S.K. and Luthra, Y.P.
Variability in structural carbohydrates and in vitro digestibility of forages. 4. Fenugreek (*Trigonella foenum-graecum*) 4(1979)17
Datta, S.C.
Estimation of lysine in compounded poultry diets after enzymic digestion 3(1978)57
Davidson, J., McIntosh, A.D. and Milne, E.
Note on the lysinoalanine content of alkali-treated barley 7(1982)217
Davis, T.P., see Belyea, R.L. 17(1987)57
de Boer, F.
Use of maize for livestock feeding in the Netherlands 1(1976)177
De Boever, J.L., see Aerts, J.V. 12(1984/1985) 47
De Boever, J.L., Cottyn, B.G., Buysse, F.X., Wainman, F.W. and Vanacker, J.M.
The use of an enzymatic technique to predict digestibility, metabolizable and net energy of compound feedstuffs for ruminants 14(1986)203
De Boever, J.L., Cottyn, B.G., Andries, J.I., Buysse, F.X. and Vanacker, J.M.

The use of a cellulase technique to predict digestibility, metabolizable and net energy of forages 19(1988)247
de Boever, J.L., see Cottyn, B.G. 21(1988)287
De Brabander, D.L., see Aerts, J.V. 3(1978) 309
De Brabander, D.L., see Aerts, J.V. 12(1984/1985)47
de Brabander, D.L., see Aerts, J.V. 2(1977) 337
de Brabander, D.L., see Andries, J.I. 18(1987) 169
De Groote, G., see Huyghebaert, G. 20(1988) 45
de Groote, G., see Ketels, E. 22(1988)105
de la Cuesta, D., see Gómez, G. 11(1984)57
De Munter, G., see Huyghebaert, G. 20(1988) 45
de Paepe, M., see Eeckhout, W. 24(1989)1
Decuypere, J.A., see Dierick, N.A. 23(1989) 141
Decuypere, J.A., see Vervaeke, I.J. 23(1989) 169
Deinum, B., Steg, A. and Hof, G.
Measurement and prediction of digestibility of forage maize in The Netherlands 10 (1983/1984)301
Del Monte, P., see Russo, V. 1(1976)25
Delort-Laval, J., see Szylit, O. 3(1978)73
Delort-Laval, J., see Cherbut, C. 20(1988)327
Delort-Laval, J., see Champ, M. 23(1989)195
Deltoro López, J. and Martinez Pascual, J.L.
Influence of the drying process on the composition of brewer's dried grains 6(1981)163
Deltoro López, J., Fernandez Carmona, J. and Martinez Pascual, J.L.
Evaluation of brewer's dried grains in the diets of laying hens 6(1981)169
Deltoro López, J. and Fernandez Carmona, J.
Evaluation of brewer's dried grains in the diets of broiler chickens 6(1981)179
Demarquilly, C., see Robert, P. 16(1986/1987) 215
Demeyer, D.I., see Kayouli, C. 10(1983/1984) 165
Demeyer, D.I., see Jouany, J.P. 21(1988)229
Demeyer, D.I., see Dierick, N.A. 23(1989)141
Demeyer, D.I., see Vervaeke, I.J. 23(1989)169
Demeyer, D.I., see Henderson, C. 23(1989)227
Dendooven, R., see Kayouli, C. 10(1983/1984) 165
Deolankar, R.P. and Singh, K.S.
Trypsin inhibitor, mineral availability, and performance of broiler chickens fed on diets based on rice bran 4(1979)133
Deschard, G., see Tetlow, R.M. 18(1987)271
Deschard, G., Tetlow, R.M. and Mason, V.C.
Treatment of whole-crop cereals with alkali. 3. Voluntary intake and digestibility studies in sheep given immature wheat ensiled with sodium hydroxide, urea or ammonia 18(1987)283
Deschard, G., Mason, V.C. and Tetlow, R.M.
Treatment of whole-crop cereals with alkali. 4. Voluntary intake and growth in steers given wheat ensiled with sodium hydroxide, urea or ammonia 19(1988)55
Deyoe, C.W., see Skoch, E.R. 6(1981)83
Dias-da-Silva, A.A. and Sundstøl, F.
Urea as a source of ammonia for improving the nutritive value of wheat straw 14(1986) 67
Dias-da-Silva, A.A., Mascarenhas Ferreira, A. and Guedes, C.V.M.
Effects of moisture level, treatment time and soya bean addition on the nutritive value of urea-treated maize stover 19(1988) 67
Dias-da-Silva, A.A., see Mascarenhas-Ferreira, A. 25(1989)157
Dierick, N.A., Vervaeke, I.J., Demeyer, D.I. and Decuypere, J.A.
Approach to the energetic importance of fibre digestion in pigs. I. Importance of fermentation in the overall energy supply 23 (1989)141
Dierick, N.A., see Vervaeke, I.J. 23(1989)169
Dinsdale, D., see Cheng, K.-J. 10(1983/1984) 93
Dinter, B., see Weinberg, Z.G. 20(1988)335
Dixon, R.M.
Effects of dietary concentrates on rumen digestion of fibrous feedstuffs 14(1986)193
Djajanegara, A., Molina, B.T. and Doyle, P.T.
The utilization of untreated and calcium hydroxide treated wheat straw by sheep 12 (1984/1985)141
Djajanegara, A. and Doyle, P.T.
Digestion rates in and outflow rates from the rumen of sheep fed untreated or calcium hydroxide-treated wheat straw 25(1989)179
D'Mello, J.P.F.
Amino acid supplementation of methanol-grown dried microbial cells in diets for young chicks 4(1979)63
D'Mello, J.P.F.
Limiting amino acids in dried microbial cells given to young turkeys 4(1979)209
D'Mello, J.P.F. and Acamovic, T.
Growth performance of, and mimosine excretion by, young chicks fed on *Leucaena leucocephala* 7(1982)247
D'Mello, J.P.F.
Utilisation of L- and DL-amino acids by broiler chicks fed diets containing Pruteen 17(1987)1
Donkoh, A., see Atuahene, C.C. 17(1987)157
Donkoh, A., Atuahene, C.C., Kese, A.G. and Mensah-Asante, B.
The nutritional value of dried coffee pulp (DCP) in broiler chickens's diets 22(1988) 139
Doran, B.E., see Goetsch, A.L. 18(1987)151
Dove, H., see Freer, M. 11(1984)87
Doyle, P.T., see Djajanegara, A. 12(1984/1985)141
Doyle, P.T., see Djajanegara, A. 25(1989)179
Draper, S.R., see El-Sherbiny, A.E. 5(1980)33

Draper, S.R., see El-Sherbiny, A.E. 5(1980)43
Dryden, G.McL. and Kempton, T.J.
Digestion of organic matter and nitrogen in ammoniated barley straw 10(1983/1984)65
Dryden, G.McL. and Leng, R.A.
Treatment of barley straw with ammonia and sulphur dioxide gases under laboratory conditions 14(1986)41
Dryden, G.McL. and Leng, R.A.
Effects of ammonia and sulphur dioxide gases on the composition and digestion of barley straw 19(1988)121
Dumay, C., see Durand, M. 21(1988)197
Durand, M., see Szylit, O. 3(1978)73
Durand, M., Dumay, C., Beaumatin, P. and Morel, M.T.
Use of the rumen simulation technique (RUSITEC) to compare microbial digestion of various by-products 21(1988)197
Dutton, M.F., see Westlake, K. 25(1989)169

E

Eddebbarh, A., see Garrett, J.E. 25(1989)11
Edney, M.J., Campbell, G.L. and Classen, H.L.
The effect of β-glucanase supplementation on nutrient digestibility and growth in broilers given diets containing barley, oat groats or wheat 25(1989)193
Edwards, R.A., see Heron, S.J.E. 19(1988)85
Edwards, S.A., Fairbairn, C.B. and Capper, A.L.
Liquid potato feed for finishing pigs: feeding value, inclusion rate and storage properties 15(1986)129
Eeckhout, W.
The energy systems used in the EEC countries: their application to 24 commercial pig-feeds and relationships between them 6 (1981)211
Eeckhout, W. and Moermans, R.
Prediction of the energy value of commercial pig feeds, according to the various energy systems used in the European Community, that are based on simple analytical parameters of the feed 6(1981)367
Eeckhout, W. and de Paepe, M.
Alkali-treated soya flour for piglets 24(1989) 1
Ehle, F.R., see Hannah, S.M. 16(1986/1987) 51
Ekpenyong, T.E.
Nutrient and amino acid composition of *Leucaena leucocephala* (Lam.) de Wit 15 (1986)183
Ekström, N., see Kaspersson, A. 19(1988)299
El-Alaily, H., see Mohamed, K. 11(1984)199
El Boushy, A.R., see Papadopoulos, M.C. 14 (1986)279
El-Din, T., see Abel, Hj. 19(1988)111
El Hag, H.M.A. and Greenhalgh, J.F.D.
Intake and digestibility in sheep fed on alkali-treated straw and concentrates, as complete diets or separate feeds 7(1982)271
El Hag, M.G., Kurdi, O.I. and Maghoub, S.O.
Performance and carcass characteristics of Sudan desert sheep and goats on high roughage diets with added fat 13(1985)147
El Khidir, O.A. and Vestergaard Thomsen, K.
The effect of high levels of molasses in combinations with hay on digestibility of organic matter, microbial protein synthesis and volatile fatty acid production in vitro 7 (1982)277
El Khidir, O.A. and Vestergaard Thomsen, K.
Effect of intake on digestibility of plant cell walls and cell contents of complete diets for ruminants 9(1983)197
El Khidir, O.A., Gumaa, A.Y., Fangali, O.A.I. and Badir, N.A.
The use of Balanites kernel cake in a diet for fattening sheep 9(1983)301
El-Negoumy, A.M., see Moss, B.R. 8(1983)25
El Shazly, K., see Abel, Hj. 19(1988)111
El-Sherbiny, A.E., Draper, S.R. and Topps, J.H.
The determination of available lysine in barley using *Tetrahymena pyriformis* W. 5 (1980)33
El-Sherbiny, A.E., Draper, S.R. and Topps, J.H.
A comparison of three methods for the determination of available lysine in barley 5 (1980)43
El Tayeb, A.E., see Osman, A.G. 19(1988)351
El-Tayeb, A.E., Abdel-Raheim, F.M. and Awadel-Kareim, A.M.
Comparison of cottonseed cake and dried blood as nitrogen sources for growing Sudan Nubian goats 22(1988)333
Elliot, J.I., see Sibbald, I.R. 8(1983)191
Elliot, J.I., Modler, H.W. and Timbers, G.E.
A continuous process for the production of purified porcine γ-globulin for use in pig-milk replacer supplements 17(1987)213
Elliott, R., see McMeniman, N.P. 19(1988)43
Elliott, R., see Alcántara, E. 23(1989)323
Elliott, R., see Tangtaweewipat, S. 25(1989) 123
Ellis, W.C., see Schake, L.M. 7(1982)257
Elwinger, K., see Hesselman, K. 7(1982)351
Ely, D.G., see Pendlum, L.C. 2(1977)225
Englyst, H.
Classification and measurement of plant polysaccharides 23(1989)27
Est, Y., see Ben-Ghedalia, D. 19(1988)219
Est, Y., see Ben-Ghedalia, D. 24(1989)289
Estergaard Thomsen, K., see Israelsen, M. 3 (1978)227
Evans, D.F. and Leibholz, J.
Meat meal in the diet of the early-weaned pig. I. A comparison of meat meal and soya bean meal 4(1979)33
Evans, D.F. and Leibholz, J.
Meat meal in the diet of the early-weaned

pig. II. Amino acid supplementation 4 (1979)43
Everington, J.M., see Givens, D.I. 24(1989)27

F

Faci, R., see Alibés, X. 10(1983/1984)239
Fadel, J.G., see Robinson, P.H. 15(1986)249
Fahey Jr., G.C., see Guardiola, C.M. 8(1983) 129
Fahey Jr., G.C., see Mora, M.I. 9(1983)205
Fahey Jr., G.C., see Lynch, G.L. 14(1986)127
Fahey Jr., G.C., see Lewis, S.M. 17(1987)179
Fairbairn, C.B., see Edwards, S.A. 15(1986) 129
Fairbairn, R., see Alli, I. 9(1983)291
Fallon, R.J., Williams, P.E.V. and Innes, G.M.
The effects on feed intake, growth and digestibility of nutrients of including calcium soaps of fat in diets for young calves 14 (1986)103
Fangali, O.A.I., see El Khidir, O.A. 9(1983) 301
Fanta, G.F., see Lewis, S.M. 17(1987)179
Farrell, D.J.
A nutritional evaluation of buckwheat (*Fagopyrum esculentum*) 3(1978)95
Farrell, D.J., Chan, C., McCrae, F. and McKenzie, R.J.
A nutritional evaluation of triticale with pigs 9(1983)49
Farrell, D.J., Thomson, E., Choice, A., Ashes, J.R., Peck, N.J. and Hogan, J.P.
Effects of milling and pelleting of maize, barley and wheat on their metabolizable energy value for cockerels and chicks 9(1983) 99
Farrell, D.J., see Raharjo, Y. 12(1984/1985)29
Faulkner, D.B., Llamas, G.L., Ward, J.K. and Klopfenstein, T.J.
Improving the intake and nutritive value of wheat straw for beef cows 12(1984/1985)125
Fay, J.P. and Ovejero, F.M.A.
Effect of lactate on the in vitro digestion of *Agropyron elongatum* by rumen microorganisms 16(1986/1987)161
Fegeros, K., see Papadopoulos, G. 18(1987) 303
Fenwick, G.R. and Curtis, R.F.
Rapeseed meal and its use in poultry diets. A review 5(1980)255
Fenwick, G.R., Pearson, A.W., Greenwood, N.M. and Butler, E.J.
Rapeseed meal tannins and egg taint 6 (1981)421
Fenwick, G.R., see Pearson, A.W. 8(1983)119
Fernandez Carmona, J., see Deltoro López, J. 6(1981)169
Fernandez Carmona, J., see Deltoro López, J. 6(1981)179
Fernandez-Carmona, J., see Cervera, C. 12 (1984/1985)233
Fernández Carmona, J., see Martinez Pascual, J. 5(1980)1
Fernández Carmona, J., see Martinez Pascual, J. 5(1980)11
Fernández Carmona, J., see Martinez Pascual, J. 5(1980)23
Fetuga, B.L., see Babatunde, G.M. 4(1979)23
Fetuga, B.L., see Balogun, O.O. 7(1982)233
Fetuga, B.L.A., see Balogun, O.O. 8(1983)197
Fetuga, B.L.A., see Ikurior, S.A. 20(1988)251
Fiallo, J.F., see Wohlt, J.E. 6(1981)115
Fiems, L.O., see Bouqué, Ch.V. 7(1982)401
Fiems, L.O., Boucqué, Ch.V., Cottyn, B.G. and Buysse, F.X.
Effect of avoparcin on performance, body composition and meat quality of finishing bulls 8(1983)237
Fiems, L.O., Cottyn, B.G., Boucqué, Ch.V. and Buysse, F.X.
Effect of formaldehyde-treated soya bean meal and urea in starters on nitrogen quality, degradibility in sacco, sheep digestibility and calf performance 16(1986/1987)287
Findlay, M., see Partridge, G.G. 13(1985)299
Findlay, M., see Partridge, G.G. 16(1986/ 1987)109
Finley, J.W., see Nelson, M.L. 24(1989)141
Fischer, J.R., see Goedeken, F.K. 13(1985)47
Fishwick, G., see Bass, J.M. 7(1982)93
Fishwick, G., see Alawa, J.P. 19(1988)33
Flachowsky, G., Wolfram, D., Wolf, I. and Hennig, A.
Digestibility and beef performance after feeding NaOH- or KOH-treated pig wastes 17(1987)235
Flám, F., see Vencl, B. 6(1981)235
Flipot, P., Mason, W. and Lalande, G.
Chemical composition and animal performance of grass forage of varying maturity stored as hay or silage 11(1984)35
Foale, M.A., see Kondos, A.C. 8(1983)85
Folman, Y., see Holzer, Z. 5(1980)299
Fonnesbeck, P.V., Garcia de Hernandez, M.M., Kaykay, J.M. and Saiady, M.Y.
Estimating yield and nutrient losses due to rainfall on field-drying alfalfa hay 16(1986/ 1987)7
Fonolla, J., Prieto, C. and Sanz, R.
Influence of age on the nutrient utilization of diets for broilers 6(1981)405
Fonolla, J., see Aguilera, J.F. 15(1986)33
Fontenot, J.P., see Rao, N.M. 18(1987)67
Fontenot, J.P., see Ortigues, I. 24(1989)13
Fonty, G., see Stewart, C.S. 21(1988)69
Fonty, G. and Gouet, Ph.
Fibre-degrading microorganisms in the monogastric digestive tract 23(1989)91
Fordyce, R.A., see Partridge, G.G. 16(1986/ 1987)109
Francesch, M., see Lacassagne, L. 20(1988)59
Franci, O., see Antongiovanni, M. 5(1980)169
Fraser, J., see Hemingway, R.G. 15(1986)123
Frederiksen, J.H., see Kategile, J.A. 4(1979)1
Freer, M. and Dove, H.

Rumen degradation of protein in sunflower meal, rapeseed meal and lupin seed placed in nylon bags 11(1984)87
Froněk, P., see Heger, J. 8(1983)163
Frydrych, Z. and Heger, J.
Nutritional evaluation, using rats, of diets based on cereals and ethanol-grown yeast 6 (1981)309
Frydrych, Z., see Heger, J. 8(1983)163
Frydrych, Z., see Heger, J. 15(1986)21
Fundora, O., see Valdivie, M. 7(1982)185
Funk, M.A., see Goetsch, A.L. 18(1987)151

G

Gabir, S., see Maglad, M.A. 15(1986)69
Galvez, J.F., see Caja, G. 2(1977)267
Garcia, G., see Alli, I. 7(1982)411
Garcia, G., see Alli, I. 9(1983)291
Garcia de Hernandez, M.M., see Fonnesbeck, P.V. 16(1986/1987)7
Garmo, T.H., see Borhami, B.E.A. 7(1982)53
Garmo, T.H., see Wanapat, M. 12(1984/1985) 295
Garrett, J.E., Guessous, F. and Eddebbarh, A.
Utilization of sugar beet molasses and monensin for finishing dairy bullocks 25(1989) 11
Garrigus, U.S., see Guardiola, C.M. 8(1983) 129
Gasa, J., Castrillo, C., Baucells, M.D. and Guada, J.A.
By-products from the canning industry as feedstuff for ruminants: digestibility and its prediction from chemical composition and laboratory bioassays 25(1989)67
Gascoyne, D.J. and Theodorou, M.K.
Consecutive batch culture – a novel technique for the in vitro study of mixed microbial populations from the rumen 21(1988) 183
Gaspari, F., see Giardini, A. 1(1976)313
Gaspari, F., see Giardini, A. 1(1976)465
Gast, C.H., see Mascarenhas Ferreira, A. 9 (1983)19
Gatel, F., Grosjean, F. and Castaing, J.
Feeding value of ensiled high-moisture maize grain with cob for growing-finishing pigs 20 (1988)145
Gatel, F., Grosjean, F. and Leuillet, M.
Utilization of white-flowered smooth-seeded spring peas (*Pisum sativum hortense*, CV Amino) by the breeding sow 22(1988)91
Gauld, S.A., see Mould, F.L. 10(1983/1984)31
Gauld, S.A., see Williams, P.E.V. 10(1983/ 1984)247
Geay, Y. and Malterre, C.
Nutritive value of dehydrated whole-crop maize 1(1976)429
Genizi, A., see Goldman, A. 18(1987)233
Giardini, A., Gaspari, F., Vechiettini, M. and Schenoni, P.
Effect of maize silage harvest stage on yield, plant composition and fermentation losses 1 (1976)313
Giardini, A., Vecchiettini, M. and Lo Bruno, A.
Energy supplementation of maize silage harvested at different maturity stages 1(1976) 369
Giardini, A., Lambertini, F., Gaspari, F. and Lo Bruno, A.
Levels, sources and methods of nitrogen supplementation of maize silage for beef production 1(1976)465
Gibson, D.M., Kennelly, J.J., Aherne, F.X. and Mathison, G.W.
Efficacy of sulfur dioxide as a grain preservative 19(1988)203
Giger, S., see Sauvant, D. 13(1985)7
Gil, F., see Aguilera, J.F. 15(1986)33
Giorgetti, A., see Antongiovanni, M. 5(1980) 169
Giouzeljannis, A., see Karalazos, A. 20(1988) 13
Girardeau, J.P., see Gouet, P. 2(1977)297
Givens, D.I. and Barber, W.P.
Nutritive value of apple pomace for ruminants 16(1986/1987)311
Givens, D.I. and Brunnen, J.M.
Nutritive value of naked oats for ruminants 18(1987)83
Givens, D.I., see Clark, P. 18(1987)225
Givens, D.I., Adamson, A.H. and Cobby, J.M.
The effect of ammoniation on the nutritive value of wheat, barley and oat straws. II. Digestibility and energy value measurements in vivo and their prediction from laboratory measurements 19(1988)173
Givens, D.I., Everington, J.M. and Adamson, A.H.
The digestibility and metabolisable energy content of grass silage and their prediction from laboratory measurements 24(1989)27
Godeau, J.-M., see Teller, E. 13(1985)25
Goedeken, F.K., Paterson, J.A., Koeln, L.L., Fischer, J.R. and Williams, J.E.
Nitrogen balance and abomasal protein-nitrogen flow in growing ruminants fed methane digester effluent in combination with distillers dried grains 13(1985)47
Goetsch, A.L., Owens, F.N., Funk, M.A. and Doran, B.E.
Effects of whole or ground corn with different forms of hay in 85% concentrate diets on digestion and passage rate in beef heifers 18 (1987)151
Goetsch, A.L., see Hardin, A.C. 25(1989)99
Goldman, A., Genizi, A., Yulzari, A. and Seligman, N.G.
Improving the reliability of the two-stage in vitro assay for ruminant feed digestibility by calibration against in vivo data from a wide range of sources 18(1987)233
Gómez, G., see Santos, J. 8(1983)313

Gómez, G. and Valdivieso, M.
Cassava for animal feeding: effect of variety and plant age on production of leaves and roots 11(1984)49
Gómez, G., Valdivieso, M., de la Cuesta, D. and Salcedo, T.S.
Effect of variety and plant age on the cyanide content of whole-root cassava chips and its reduction by sun-drying 11(1984)57
Gomez, M., see Abate, A.N. 10(1983/1984)291
Gonzalez, G. and Monteagudo, A.
Survey of the use of maize for livestock feeding in Spain 1(1976)129
Gonzalez, G., see Alzueta, C. 8(1983)155
González, G., Alzueta, C., Barro, C. and Salvador, A.
Yield and composition of protein concentrate, press cake, green juice and solubles concentrate from wet fractionation of *Sophora japonica* L. foliage 20(1988)177
González, G., see Rebolé, A. 22(1988)173
Gonzalez Santillana, R., see Wilkinson, J.M. 3(1978)117
Gonzalez Santillana, R., see Wilkinson, J.M. 3(1978)133
Gonzalez Santillana, R. and Wilkinson, J.M.
Effect of time of storage and method of drying on the digestibility in vitro of alkali-treated barley straw 3(1978)381
Goode, L., see Johnson, T.O. 12(1984/1985) 65
Göransson, L., Olsson, O. and Larsson, K.
Effects of different handling systems on pig feed quality 7(1982)11
Göransson, L. and Ogle, R.B.
Anaerobically stored high moisture cereals for growing pigs 12(1984/1985)159
Gorbet, D.W., see Myer, R.O. 12(1984/1985) 179
Gordon, A.H. and Chesson, A.
The effect of prolonged storage on the digestibility and nitrogen content of ammonia-treated barley straw 8(1983)147
Gordon, G.L.R., see Akin, D.E. 23(1989)305
Gorman, T.B.S., see Batterham, E.S. 9(1983) 169
Goto, I. and Minson, D.J.
Prediction of the dry matter digestibility of tropical grasses using a pepsin-cellulase assay 2(1977)247
Goto, I., see Shimojo, M. 24(1989)173
Gouet, P., Girardeau, J.P. and Riou, Y.
Inhibition of *Listeria monocytogenes* by defined lactic microflora in gnotobiotic silages of lucerne, fescue, ryegrass and maize – influence of dry matter and temperature 2 (1977)297
Gouet, Ph., see Stewart, C.S. 21(1988)69
Gouet, Ph., see Fonty, G. 23(1989)91
Gould, J.M., see Lewis, S.M. 17(1987)179
Goulden, K.M., see Whitehead, D.C. 14(1986) 231
Graham, C.A., see Kellaway, R.C. 3(1978)201
Graham, H. and Åman, P.
A comparison between degradation in vitro and in sacco of constituents of untreated and ammonia-treated barley straw 10(1983/1984)199
Graham, H., Åman, P., Theander, O., Kolankaya, N. and Stewart, C.S.
Influence of heat sterilization and ammoniation on straw composition and degradation by pure cultures of cellulolytic rumen bacteria 12(1984/1985)195
Graham, H., see Åman, P. 17(1987)15
Graham, H. and Åman, P.
Whole-crop peas. II. Digestion of early- and late-harvested crops in the gastrointestinal tract of pigs 17(1987)33
Graham, H., see Nordkvist, E. 17(1987)295
Graham, H., see Theander, O. 23(1989)205
Graham, H., see Narasimhalu, P. 24(1989)159
Grain, J., see Jouany, J.P. 21(1988)229
Green, A.G. and Oram, R.N.
Variability for protein and oil quality in *Lupinus albus* 9(1983)271
Green, S., Wiseman, J. and Cole, D.J.A.
Examination of stability, and its effect on the nutritive value, of fish silage in diets for growing pigs 21(1988)43
Greenhalgh, J.F.D.
The dilemma of animal feeds and nutrition 1(1976)1
Greenhalgh, J.F.D., see Manten, A.A. 2(1977) 1
Greenhalgh, J.F.D., see Pirie, R. 3(1978)143
Greenhalgh, J.F.D., Pirie, R., Shin, H.T. and Stewart, C.S.
Akali treatment of straw for ruminants. II. Nutritive value of straw ensiled after alkali treatment. 3(1978)289
Greenhalgh, J.F.D., see El Hag, H.M.A. 7 (1982)271
Greenhalgh, J.F.D., see Charmley, E. 18(1987) 19
Greenwood, N.M., see Fenwick, G.R. 6(1981) 421
Greenwood, N.M., see Pearson, A.W. 8(1983) 119
Gregory, K.F., Reade, A.E., Santos-Nunez, J., Alexander, J.C., Smith, R.E. and MacLean, S.J.
Further thermotolerant fungi for the conversion of cassava starch to protein 2(1977) 7
Greife, H.A., Molnar, S. and Günther, K.-D.
Nitrogen metabolism in broiler chickens consuming the bacterial strain *Alcaligenes eutrophus* 5(1980)241
Grenet, E. and Barry, P.
Colonization of thick-walled plant tissues by anaerobic fungi 19(1988)25
Griffith, E.S., see McAllan, A.B. 17(1987)65
Grigera-Naón, J.J., see Smith, T. 10(1983/1984)189
Grigera-Naón, J.J., see Smith, T. 14(1986)29

Grimbergen, A.H.M., Cornelissen, J.P. and Stappers, H.P.
The relative availability of phosphorus in inorganic feed phosphates for young turkeys and pigs 13(1985)117
Grimes, L.W., see Grotheer, M.D. 14(1986)55
Grosjean, F., see Gatel, F. 20(1988)145
Grosjean, F., see Gatel, F. 22(1988)91
Grotheer, M.D., Cross, D.L. and Grimes, L.W.
Effect of ammonia level and time of exposure to ammonia on nutritional and preservatory characteristics of dry and high-moisture coastal bermuda grass hay 14 (1986)55
Grubb, D.A., see Ørskov, E.R. 2(1977)307
Grubb, D.A., see Ørskov, E.R. 6(1981)273
Grummer, R.H., see Cieslak, D.G. 14(1986) 171
Grummitt, N., see Minson, D.J. 9(1983)221
Guada, J.A., see Gasa, J. 25(1989)67
Guardiola, C.M., Fahey Jr., G.C., Spears, J.W., Garrigus, U.S., Izquierdo, O.A. and Pedroza, C.
The effects of sulphur supplementation on cellulose digestion in vitro and on nutrient digestion, nitrogen metabolism and rumen characteristics of lambs fed on good quality fescue and tropical star grass hays 8(1983) 129
Guedes, C.V.M., see Dias-da-Silva, A.A. 19 (1988)67
Guedes, C.V.M., see Mascarenhas-Ferreira, A. 25(1989)157
Guenter, W., see Al-Athari, A.K. 22(1988)119
Guenter, W., see Al-Athari, A.K. 22(1988)273
Guenter, W., see Rotter, B.A. 24(1989)233
Guessous, F., see Garrett, J.E. 25(1989)11
Gumaa, A.Y., see El Khidir, O.A. 9(1983)301
Günther, K.-D., see Greife, H.A. 5(1980)241

H

Ha, J.K., Kennelly, J.J. and Berzins, R.
Effect of dietary nitrogen source on microbial protein synthesis, dietary protein degradation and nutrient digestion in steers 14(1986)117
Hadjipanayiotou, M.
The value of urea-treated straw in diets of lactating goats 11(1984)67
Hadobás, K., see Hegedüs, M. 9(1983)79
Hahn-Hägerdal, B., Tjerneld, F. and Zacchi, G.
Production of ethanol from lignocellulosic materials 21(1988)175
Håkansson, J., see Lund, D. 16(1986/1987)119
Hall, J.M.R., see Wanapat, M. 14(1986)215
Hall, K.L., see Hardin, A.C. 25(1989)99
Hamilton, R.M.G. and Trenholm, H.L.
Observations on the chemical and nutritive content of white winter and spring wheats contaminated with deoxynivalenol (vomitoxin) 11(1984)293
Hanczakowska, E., see Hanczakowski, P. 6 (1981)189
Hanczakowski, P. and Petzel, M.
Nutritive value of leaf protein coagulum and cereals mixtures 4(1979)91
Hanczakowski, P., Skraba, B. and Hanczakowska, E.
Nutritive value of lupin-seed protein and leaf-protein concentrates supplemented with various sulphur sources 6(1981)189
Hanczakowski, P., Skraba, B. and Mlodkowski, M.
Nutritive value of leaf-protein concentrate from potato haulm for rats and chicks 6 (1981)413
Hanczakowski, P., see Carlsson, R. 11(1984) 239
Hanczakowski, P. and Skraba, B.
The effect of different precipitating agents on quality of leaf protein concentrate from lucerne 12(1984/1985)11
Hanczakowski, P. and Skraba, B.
The effect of toxic substances present in some leaf protein concentrates on their nutritive value 24(1989)151
Hannah, S.M., Stern, M.D. and Ehle, F.R.
Evaluation of a dual flow continuous culture system for estimating bacterial fermentation in vivo of mixed diets containing various soya bean products 16(1986/1987)51
Hanrahan, T.J.
Effect of pellet size and pellet quality on pig performance 10(1983/1984)277
Hansen, P., see Opstvedt, J. 18(1987)181
Hanssen, J.F., see Muindi, P.J. 6(1981)197
Hardin, A.C., Goetsch, A.L., Landis, K.M., Murphy, G.E., Johnson, Z.B. and Hall, K.L.
Intake, digestion and daily gain by cattle consuming bermudagrass (*Cynodon dactylon*) and supplemented with different combinations of ground corn, vegetable oil, urea and corn gluten and blood meals 25(1989) 99
Hargreave, G.T., see Leibholz, J. 5(1980)309
Harmon, R.J., see Teh, T.H. 17(1987)257
Harper, A.F., Kornegay, E.T., Bryant, K.L. and Thomas, H.R.
Efficacy of virginiamycin and a commercially-available lactobacillus probiotic in swine diets 8(1983)69
Harte, F.J.
The current situation regarding the use of maize in Ireland 1(1976)195
Hartley, R.D., see Theander, O. 10(1983/ 1984)89
Hartley, R.D., see Van Soest, P.J. 10(1983/ 1984)155
Hartley, R.D., see Whitehead, D.C. 14(1986) 231
Hartley, R.D., see Mason, V.C. 19(1988)159
Hartley, R.D., see Akin, D.E. 19(1988)261
Hartley, R.D., see Mason, V.C. 24(1989)299
Harvey, R.W., see Johnson, T.O. 12(1984/ 1985)65

Harvey, R.W., see Ortigues, I. 24(1989)13
Harvey, R.W. and Spears, J.W.
Effect of blood meal processing method on performance and nitrogen utilization by growing cattle 24(1989)337
Hasimoglu, S., see Umunna, N.N. 7(1982)375
Hazlewood, G.P., see Orpin, C.G. 21(1988)161
Heath, M.E., see Waterworth, D.G. 6(1981) 297
Hector, D.A., see Machin, D.H. 11(1984)247
Hector, D.A., see Machin, D.H. 15(1986)273
Hegedüs, M., Bokori, J. and Hadobás, K.
Protein quality of mixtures of animal protein meals 9(1983)79
Heger, J., see Frydrych, Z. 6(1981)309
Heger, J., Frydrych, Z. and Froněk, P.
Evaluation of optimum lysine and threonine supplements to a wheat and barley-based diet in rats 8(1983)163
Heger, J. and Frydrych, Z.
Estimation of optimum supplements of essential amino acids to a wheat and barley-based diet in rats using a nitrogen balance technique 15(1986)21
Heikonen, M., see Moisio, T. 22(1988)341
Hembry, F.G., see Coelho, M. 20(1988)219
Hemingway, R.G., Parkins, J.J. and Fraser, J.
Sugar beet pulp products for dairy cows 15 (1986)123
Hemingway, R.G., see Alawa, J.P. 19(1988)33
Hemken, R.W., see Teh, T.H. 17(1987)257
Henderson, A.R., McDonald, P. and Anderson, D.H.
The effect of silage additives containing formaldehyde on the fermentation of ryegrass ensiled at different dry matter levels and on the nutritive value of direct-cut silage 7(1982)303
Henderson, A.R., see McDonald, P. 8(1983) 259
Henderson, C. and Demeyer, D.I.
The rumen as a model of the microbiology of fibre digestion in the human intestine 23 (1989)227
Hendrosoekarjo, S. and Pearce, G.R.
Utilization by sheep of dried pig faeces containing a high concentration of copper 3 (1978)31
Hennig, A., see Flachowsky, G. 17(1987)235
Hennig, A., see Schöne, F. 22(1988)45
Henry, P.R., see Potter, S.G. 12(1984/1985) 275
Hermansen, J.E.
Feed intake and milk yield at increasing supplement of a palmitic and stearic acid-rich type of fat in comparison with animal fat 22(1988)179
Hermansen, J.E.
Feed intake, milk yield and milk composition by replacing unprotected fat by Ca-soaps for dairy cows 22(1988)193
Heron, S.J.E., Edwards, R.A. and McDonald, P.
The effects of inoculation, addition of glucose and mincing on fermentation and proteolysis in ryegrass ensiled in laboratory silos 19(1988)85
Hesselman, K., Elwinger, K. and Thomke, S.
Influence of increasing levels of β-glucanase on the productive value of barley diets for broiler chickens 7(1982)351
Hesselman, K. and Åman, P.
The effect of β-glucanase on the utilization of starch and nitrogen by broiler chickens fed on barley of low- or high-viscosity 15 (1986)83
Hesselman, K., see Pettersson, D. 17(1987)145
Hibberd, C.A., Wagner, D.G., Schemm, R.L., Mitchell Jr., E.D., Weibel, D.E. and Hintz, R.L.
Response of different sorghum grain and maize varieties to reconstitution 15(1986) 231
Hibberd, C.A., see Chase Jr., C.C. 24(1989) 129
Higgins, T.R. and Spooner, A.E.
Microwave drying of alfalfa compared to field- and oven-drying: effects on forage quality 16(1986/1987)1
Hinds, M.A., Bolsen, K.K., Brethour, J., Milliken, G. and Hoover, J.
Effects of molasses/urea and bacterial inoculant additives on silage quality, dry matter recovery, and feeding value for cattle 12 (1984/1985)205
Hinks, C.E.
The replacement of skim milk by dried microbial cells and whey in milk diets for calves 2(1977)85
Hintz, R.L., see Hibberd, C.A. 15(1986)231
Hlödversson, R. and Kaspersson, A.
Nutrient losses during deterioration of hay in relation to changes in biochemical composition and microbial growth 15(1986)149
Hlödversson, R.
The nutritive value of white- and dark-flowered cultivars of pea for growing-finishing pigs 17(1987)245
Hlödversson, R.
The effect of mechanical treatment on the nutritive value of high-moisture barley and the performance of growing-finishing pigs 24 (1989)121
Ho, Y.W., Abdullah, N. and Jalaludin, S.
Colonization of guinea grass by anaerobic rumen fungi in swamp buffalo and cattle 22 (1988)161
Hoadley, C.J., see Mason, V.C. 24(1989)313
Hoebler, C., see Champ, M. 23(1989)195
Hof, G., see Deinum, B. 10(1983/1984)301
Hogan, J.P., see Farrell, D.J. 9(1983)99
Holland, S.M., see Ørskov, E.R. 8(1983)247
Holzer, Z., Levy, D. and Folman, Y.
Performance of fattening cattle and lactating beef cows on diets high in chemically treated wheat straw and poultry litter 5 (1980)299

Holzer, Z., Levy, D., Samuel, V., Angemar, Y. and Ilan, D.
Wheat silage for fattening young male cattle 12(1984/1985)253
Holzer, Z., see Levy, D. 16(1986/1987)63
Holzgraefe, D.P., see Lewis, S.M. 17(1987)179
Hoover, J., see Hinds, M.A. 12(1984/1985)205
Horn, G.W., see Streeter, C.L. 7(1982)325
Horn, G.W., Batchelder, D.G., Manor, G., Streeter, C.L. and McLaughlin, G.L.
Ammoniation of wheat straw and native grass hay during baling of large round bales 8(1983)35
Horn, G.W., see Zorrilla-Rios, J. 22(1988)305
Horn, G.W., Zorilla-Rios, J. and Akin, D.E.
Influence of stage of forage maturity and ammoniation of wheat straw on ruminal degradation of wheat forage tissues 24 (1989)201
Horner, J.I., see Coppock, C.E. 18(1987)89
Horton, G.M.J., Nicholson, H.H. and Christensen, D.A.
Ammonia and sodium hydroxide treatment of wheat straw in diets for fattening steers 7(1982)1
Horvath, D.J., see Reid, R.L. 5(1980)95
Horvath, P.J., see Jeraci, J.L. 23(1989)121
Hosking, Z.D., see Barber, R.S. 4(1979)117
Hove, E.L., see Ruiz Jr., L.P. 2(1977)59
Huhtanen, P.
The effects of barley, unmolassed sugar-beet pulp and molasses supplements on organic matter, nitrogen and fibre digestion in the rumen of cattle given a silage diet 20(1988) 259
Huhtanen, P. and Robertson, S.
The effect of dietary inclusion of starch, surcrose and xylose on the utilization of dietary energy in sheep 21(1988)11
Huisman, J., see Van Weerden, E.J. 2(1977) 377
Hulan, H.W., see Zarkadas, C.G. 14(1986)291
Hulan, H.W., Proudfoot, F.G. and Zarkadas, C.G.
The effect of adding white fish meal containing enzyme digested or untreated stickwater solids to diets for broiler chickens 16(1986/ 1987)253
Hunt, C.W., see Koeln, L.L. 11(1984)181
Hunt, J.R. and Barton, G.M.
Nutritive value of spruce muka (foliage) for the growing chick 3(1978)63
Husain, R.A.K. and Offer, N.W.
Effect of formaldehyde treatment on the degradation of acid-preserved fish silage protein in vitro 16(1986/1987)297
Husain, R.A.K., see Offer, N.W. 17(1987)165
Hussar, N., see Leeson, S. 19(1988)313
Hutagalung, R.I., see Abdullah, N. 20(1988) 79
Huyghebaert, G., De Munter, G. and De Groote, G.
The metabolisable energy (AME_n) of fats for broilers in relation to their chemical composition 20(1988)45
Hyslop, J.J., Weir, J., Offer, N.W., Reid, S.J. and Wilcock, J.M.
Effect of formaldehyde treatment on the rumen digestion of barley and wheat grain and its influence on hay intake and digestion by sheep 25(1989)55

I

Ibbotson, C.F., Mansbridge, R. and Adamson, A.H.
Commercial experience of treating straw with ammonia 10(1983/1984)223
Ibraham, M.N.M., Tamminga, S. and Zemmelink, G.
Effect of urea treatment on rumen degradation characteristics of rice straws 24(1989) 83
Igwe, F.O., see Udedibie, A.B.I. 24(1989)111
Ihnat, M., see Ivan, M. 9(1983)131
Ikurior, S.A. and Fetuga, B.L.A.
Chemical and biological evaluation of Nigerian cottonseed meal protein quality 20 (1988)251
Ilan, D., see Holzer, Z. 12(1984/1985)253
Ilan, D., see Levy, D. 16(1986/1987)63
Ilian, M.A., Razzaque, M.A., Al-Awadi, A. and Salman, A.J.
Use of fat in diets of sheep in hot environments. I. Effects on performance, carcass characteristics and lipid composition of plasma 19(1988)327
Ilian, M.A., Razzaque, M.A. and Salman, A.J.
Use of fat in diets of sheep in hot environments. II. Effect on rumen metabolism 19 (1988)343
Ilori, J.O., see Smith, O.B. 11(1984)231
Ingalls, J.R., see Sharma, H.R. 10(1983/1984) 77
Ingalls, J.R., see Sharma, H.R. 15(1986)101
Ingalls, J.R., see Mbugi, P.K. 24(1989)267
Innes, G.M., see Ørskov, E.R. 6(1981)273
Innes, G.M., see Ørskov, E.R. 6(1981)355
Innes, G.M., see Williams, P.E.V. 10(1983/ 1984)247
Innes, G.M., see Williams, P.E.V. 11(1984)103
Innes, G.M., see Williams, P.E.V. 11(1984)115
Innes, G.M., see Williams, P.E.V. 12(1984/ 1985)1
Innes, G.M., see Fallon, R.J. 14(1986)103
Israelsen, M., Rexen, B. and Estergaard Thomsen, K.
Cellulase insoluble fibre as a measure of unavailable organic matter in cattle compounds containing alkali-treated straw 3 (1978)227
Istasse, L., Reid, G.W., Tait, C.A.G. and Ørskov, E.R.
Concentrates for dairy cows: effects of feeding method, proportion in diet and type 15 (1986)167
Istasse, L., Smart, R.I. and Ørskov, E.R.

Comparison between two methods of feeding concentrate to sheep given a diet high or low in concentrate with or without buffering substances 16(1986/1987)37
Istasse, L., Rollin, F., Kameni Djiele, N., Van Eenaeme, C., Biston, L. and Bienfait, J.M. Discriminating power of the nylon bag technique and the pepsin-cellulase method to detect less commonly used components included in concentrates for dairy cows 25 (1989)1
Ivan, M., Ihnat, M. and Veira, D.M. Solubility and flow of calcium, magnesium and phosphorus in the digestive tract of sheep given maize or alfalfa silages 9(1983) 131
Izquierdo, O.A., see Guardiola, C.M. 8(1983) 129

J

Jackson, A.J., see Anglesea, J.D. 13(1985)39
Jackson, M.G. Review article: the alkali treatment of straws 2(1977)105
Jackson, M.G. Who needs feeding standards? 6(1981)101
Jahreis, G., see Schöne, F. 22(1988)45
Jain, V.K., Jayal, M.M. and Pathak, N.N. Nutritional evaluation of gram (*Cicer arietinum*) and arhar (*Cajanus cajan*) chunies for cattle 5(1980)315
Jakhmola, R.C., see Kamra, D.N. 9(1983)185
Jakhmola, R.C., see Singh, R. 12(1984/1985) 133
Jakhmola, R.C., Kundu, S.S., Punj, M.L., Singh, K., Kamra, D.N. and Singh, R. Animal excreta as ruminant feed – scope and limitations under Indian conditions 19 (1988)1
Jalaludin, S., see Ho, Y.W. 22(1988)161
James, K.A.C., see Smith, W.C. 18(1987)143
Janssen, W.M.M.A., see Kan, C.A. 19(1988) 97
Jayal, M.M., see Jain, V.K. 5(1980)315
Jayasuriya, M.C.N. and Panditharatne, S. Spent tea leaf as a ruminant feed 3(1978) 219
Jayasuriya, M.C.N., Perdok, H.B., Ross-Parker, H.M. and van Houtert, M.F.J. Effect of alkali-treated rice straw supplemented with spent tea leaf and thyroprotein on milk yield, milk composition and certain physiological parameters of dairy cows 7 (1982)201
Jayasuriya, M.C.N., Wijeyatunge, C. and Perera, H.G.D. Rumen and post-rumen fermentation of spent tea leaf protein and other protein sources studied by the nylon bag method 7(1982)221
Jayasuriya, M.C.N. and Pearce, G.R. The effect of urease enzyme on treatment time and the nutritive value of straw treated with ammonia as urea 8(1983)271
Jelev, K., see Peteva-Vantcheva, Z. 3(1978)23
Jensen, A.H., see Adams, K.L. 12(1984/1985) 267
Jensen, A.H., see Adams, K.L. 13(1985)159
Jensen, A.H., see Adams, K.L. 17(1987)201
Jeraci, J.L. and Lewis, B.A. Determination of soluble fiber components: (1-<3; 1-<4)-β-D-glucans and pectins 23 (1989)15
Jeraci, J.L. and Horvath, P.J. In vitro fermentation of dietary fiber by human fecal organisms 23(1989)121
Jewell, S.N. and Campling, R.C. Aqueous ammonia treatment of wheat straw: voluntary intake and digestibility in cattle 14(1986)81
Job, T.A., see Aregheore, E.M. 20(1988)233
Johnson, K.D., see Cherney, J.H. 17(1987)45
Johnson, T.O., Harvey, R.W., Goode, L., Linnerud, A.C. and Crickenberger, R.G. Effects of stage of maturity and addition of molasses on nutritive value of maize stover silage 12(1984/1985)65
Johnson, W.L., see Van Eys, J.E. 18(1987)197
Johnson, Z.B., see Noland, P.R. 2(1977)219
Johnson, Z.B., see Waldroup, P.W. 3(1978)329
Johnson, Z.B., see Noland, P.R. 5(1980)51
Johnson, Z.B., see Waldroup, P.W. 7(1982)419
Johnson, Z.B., see Hardin, A.C. 25(1989)99
Jones, A.S., see Livingstone, R.M. 2(1977)31
Jones, A.S., see Livingstone, R.M. 2(1977)277
Joshi, A.L., see Rangnekar, D.V. 7(1982)61
Jouany, J.P. and Thivend, P. In vitro effects of avoparcin on protein degradability and rumen fermentation 15 (1986)215
Jouany, J.P., Demeyer, D.I. and Grain, J. Effect of defaunating the rumen 21(1988) 229
Jovanović, M. and Čuperlović, M. Nutritive value of rumen contents for monogastric animals 2(1977)351
Jovanović, M., see Nikolić, J.A. 15(1986)57

K

Kabaija, E. and Smith, O.B. The effect of age of regrowth on content and release of manganese, iron, zinc and copper from four tropical forages incubated in sacco in rumen of sheep 20(1988)171
Kadirvel, R., see Thanu, K. 9(1983)263
Kadirvel, R., see Poyyamozhi, V.S. 15(1986) 95
Kadirvel, R., see Subramanian, P.R. 20(1988) 343
Kadirvel, R., see Chandrasekaran, D. 22(1988) 321
Kadirvel, R., see Viswanathan, K. 22(1988) 327

Kadirvel, R., see Natanam, R. 25(1989)201
Kalač, P. and Kyzlink, V.
Losses of beta-carotene in red clover in an acid medium during ensiling 4(1979)81
Kalač, P. and Kyzlink, V.
The enzymic nature of the degradation of beta-carotene in red clover and in other forage crops during silage-making with acid additives 5(1980)59
Kalač, P.
Losses of beta-carotene in unwilted forage crops during silage-making and feeding 9 (1983)63
Kali, J., see Levy, D. 16(1986/1987)63
Kameni Djiele, N., see Istasse, L. 25(1989)1
Kamra, D.N., Singh, R., Jakhmola, R.C. and Srivastava, R.V.N.
Effect of wilting and the additives, straw, molasses and urea on the fermentation pattern of maize silage 9(1983)185
Kamra, D.N., see Singh, R. 12(1984/1985)133
Kamra, D.N., see Jakhmola, R.C. 19(1988)1
Kamra, D.N., Lall, D., Kewalramant, N. and Pathak, N.N.
Ensiling characteristics of cattle waste fermented with wheat straw and green berseem (*Trifolium alexandrium*) 25(1989)149
Kan, C.A., Scheele, C.W. and Janssen, W.M.M.A.
The energy content of full-fat soya beans in meal and pelleted feeds for adult cocks and broilers 19(1988)97
Kaptur, T., see Carlsson, R. 11(1984)239
Karalazos, A. and Swan, H.
The nutritional value for sheep of molasses and condensed molasses solubles 2(1977)143
Karalazos, A. and Giouzeljannis, A.
A note on the use of sugar-beet pulp silage and molasses in the diet of lactating dairy cows 20(1988)13
Karamitros, D.
Sugar beet molasses for growing and fattening pigs 18(1987)131
Karue, C.N., see Abate, A. 6(1981)15
Karunajeewa, H. and Tham, S.H.
The influence of oat groats and dietary level of lysine on the laying performance of crossbred hens 17(1987)271
Kaspersson, A., see Hlödversson, R. 15(1986) 149
Kaspersson, A., Lindgren, S. and Ekström, N.
Microbial dynamics in barley grain stored under controlled atmosphere 19(1988)299
Kategile, J.A. and Frederiksen, J.H.
Effect of level of sodium hydroxide treatment and volume of solution on the nutritive value of maize cobs 4(1979)1
Kategile, J.A.
Performance of heifers fed on diets based on NaOH-treated maize cobs and the effect of supplementary urea and source of carbohydrates 4(1979)97
Kategile, J.A., Urio, N.A., Sundstøl, F. and Mzihirwa, Y.G.
Simplified method for alkali treatment of low-quality roughages for use by smallholders in developing countries 6(1981)133
Kategile, J.A., see Kiangi, E.M.I. 6(1981)377
Kategile, J.A.
Feed value of Tanzanian sisal wastes: chemical composition and digestibility in vitro 14 (1986)255
Kategile, J.A., Mngulwi, J.G.J. and Abate, A.
Net energy for gain (NE_{gain}) of cassava tops 20(1988)97
Kategile, J.A., see Sarwatt, S.V. 22(1988)237
Katz, R.J., see Thomas, V.M. 11(1984)269
Kaufman, W.
Discussion on session 4 1(1976)485
Kay, M., see Mbatya, P.B.A. 8(1983)221
Kay, M., see Mbatya, P.B.A. 8(1983)229
Kay, M., see Mbatya, P.B.A. 13(1985)281
Kay, M., see Mbatya, P.B.A. 13(1985)293
Kaykay, J.M., see Fonnesbeck, P.V. 16(1986/ 1987)7
Kayongo-Male, H., see Abate, A. 6(1981)15
Kayongo-Male, H., see Nangole, F.N. 9(1983) 121
Kayouli, C., Demeyer, D.I., Van Nevel, C.J. and Dendooven, R.
Effect of defaunation on straw digestion in sacco and on particle retention in the rumen 10(1983/1984)165
Keene, A.S., see Mason, V.C. 19(1988)159
Keene, A.S., see Mason, V.C. 24(1989)299
Keith, M.O., see Bell, J.M. 24(1989)253
Kellaway, R.C., Crofts, F.C., Thiago, L.R.L., Redman, R.G., Leibholz, J.M.L. and Graham, C.A.
A new technique for upgrading the nutritive value of roughages under field conditions 3 (1978)201
Kellaway, R.C., see Leibholz, J. 5(1980)309
Kellaway, R.C., see Thiago, L.R.L. de S. 7 (1982)71
Kellaway, R.C., see Amaning-Kwarteng, K. 16 (1986/1987)75
Kelleher, D.L., see Caffrey, P.J. 19(1988)231
Kelly, P.
Sugar beet pulp – a review 8(1983)1
Kempton, T.J., see Abidin, Z. 6(1981)145
Kempton, T.J., see Dryden, G.McL. 10(1983/ 1984)65
Kennedy, S., see Mbatya, P.B.A. 13(1985)281
Kennedy, S., see Mbatya, P.B.A. 13(1985)293
Kennelly, J.J., see Ha, J.K. 14(1986)117
Kennelly, J.J., see Gibson, D.M. 19(1988)203
Kernan, J.A., Coxworth, E.C. and Spurr, D.T.
New crop residues and forages for Western Canada: assessment of feeding value in vitro and response to ammonia treatment 6(1981) 257
Kernan, J.A., Crowle, W.L., Spurr, D.T. and Coxworth, E.C.
The nutritional value of crop residue components from several wheat cultivars grown at different fertilizer levels 11(1984)301
Kernan, J.A., see Mann, M.E. 21(1988)57
Kerstens, J., see Mascarenhas Ferreira, A. 9

(1983)19
Kese, A.G., see Donkoh, A. 22(1988)139
Ketelaars, E.H., see Papadopoulos, M.C. 14 (1986)279
Ketels, E. and de Groote, G.
The nutritional value for broilers of fats characterized by short-chain fatty acids as affected by level of inclusion and age 22 (1988)105
Kewalramant, N., see Kamra, D.N. 25(1989) 149
Kharat, S.T., see Rangnekar, D.V. 7(1982)61
Kiangi, E.M.I. and Kategile, J.A.
Different sources of ammonia for improving the nutritive value of low quality roughages 6(1981)377
Kies, A.K., Moughan, P.J. and Smith, W.C.
The apparent and true ileal digestibility of nitrogen and amino acids in lactic casein for the growing pig 16(1986/1987)169
Kilkenny, J.B.
The use of maize for livestock feeding in the United Kingdom 1(1976)199
King, R.H. and Campbell, R.G.
Blood meal as a source of protein for grower/finisher pigs 3(1978)191
King, R.H.
Lupin-seed meal (*Lupinus albus* cv. Hamburg) as a source of protein for growing pigs 6(1981)285
Kipnis, T., see Silanikove, N. 20(1988)87
Kirby, A.C., see Amaning-Kwarteng, K. 16 (1986/1987)75
Kirchgessner, M., see Roth, F.X. 1(1976)33
Kirchgessner, M., see Stehr, W. 1(1976)53
Kirchgessner, M., see Weigand, E. 5(1980)221
Kirchgessner, M., Schwarz, F.J. and Lindner, H.P.
Feed consumption and performance of dairy cows with alternate feedings of grass silage and maize silage 6(1981)337
Kirschgessner, M., see Kreuzer, M. 16(1986/1987)233
Kirton, A.H., see Wright, D.E. 2(1977)93
Klasing, K.C., see Brown, D.L. 18(1987)247
Klasing, K.C., see Shqueir, A.A. 25(1989)137
Klopfenstein, T.J., see Umunna, N.N. 7(1982) 375
Klopfenstein, T.J., see Faulkner, D.B. 12 (1984/1985)125
Klopfenstein, T.J., see Umunna, N.N. 12 (1984/1985)151
Klopfenstein, T.J., see Umunna, N.N. 14 (1986)161
Koeln, L.L., Hunt, C.W., Bowman, D.K. and Paterson, J.A.
Maize whole stillage as a supplemental protein source for growing lambs and finishing steers 11(1984)181
Koeln, L.L., see Goedeken, F.K. 13(1985)47
Koes, R.M., see Rule, D.C. 15(1986)113
Kolankaya, N., see Graham, H. 12(1984/1985) 195
Kondos, A.C. and Foale, M.A.
Comparison of the nutritional value of low and medium tannin sorghum grains for pigs 8(1983)85
Koong, L.J., see Sidahmed, A.E. 8(1983)47
Kornegay, E.T.
Protein digestibility of hydrolyzed hog hair meal for pigs 3(1978)323
Kornegay, E.T., see Harper, A.F. 8(1983)69
Kornegay, E.T., see Ravindran, V. 17(1987) 227
Kothandaraman, P., see Narahari, D. 9(1983) 319
Kothandaraman, P., see Narahari, D. 10 (1983/1984)257
Koukoura, Z.
Composition of kermes oak browse as affected by shade and stage of maturity 21 (1988)1
Kowalczyk, J., Robinson, J.J. and Otwinowska, A.
The digestion in the small intestine of young bulls of the protein of rapeseed meal treated or untreated with formaldehyde 7(1982)225
Kreuzer, M., Kirschgessner, M. and Müller, H.L.
Effect of defaunation on the loss of energy in wethers fed different quantities of cellulose and normal or steamflaked maize starch 16 (1986/1987)233
Krishna, N., see Chandra, S. 12(1984/1985) 187
Krishna Reddy, G.V. and Raj Reddy, M.
Nutritive value of rice straw (*Oryza sativa*) ensiled with animal excreta and rumen digesta 24(1989)69
Kromann, R.P., see Murdock, F.R. 2(1977)287
Kugenheim, Y., see Silanikove, N. 20(1988)87
Kumaresan, A., Mshelia, T.A. and Aliu, Y.O.
Biochemical evaluation of bagaruwa seeds (*Acacia nilotica*) for use as livestock feed 11 (1984)45
Kundu, S.S., see Jakhmola, R.C. 19(1988)1
Kurdi, O.I., see El Hag, M.G. 13(1985)147
Kyzlink, V., see Kalač, P. 4(1979)81
Kyzlink, V., see Kalač, P. 5(1980)59

L

Laber, B.
Survey of the use of maize for livestock feeding in Austria 1(1976)153
Lacassagne, L., Francesch, M., Carré, B. and Melcion, J.P.
Utilization of tannin-containing and tannin-free faba beans (*Vicia faba*) by young chicks: effects of pelleting feeds on energy, protein and starch digestibility 20(1988)59
Lacey, J., see Lord, K.A. 6(1981)73
Lacey, J., Lord, K.A. and Cayley, G.R.
Chemicals for preventing moulding in damp hay 6(1981)323
Lacey, J., see Lord, K.A. 7(1982)287
Lacey, J., Lord, K.A. and Cayley, G.R.

Problems of testing novel chemicals for the preservation of damp hay 8(1983)283
Lahore, R. and Croke, B.
Energetics of stockfeed production 3(1978) 1
Laksevela, B.
Adjustments of the protein level for slaughter pigs and supplementation with lysine and methionine 1(1976)619
Lalande, G., see Flipot, P. 11(1984)35
Lall, D., Lohan, O.P., Vaid, J., Sharma, O.P. and Negi, S.S.
Can lantana seeds be utilized in lamb rations? 9(1983)29
Lall, D., see Makkar, H.P.S. 20(1988)1
Lall, D. and Prasad, T.
Compositional quality of certain uncoventional calcium and phosphorus sources in India for use as mineral supplements for livestock 23(1989)343
Lall, D., see Kamra, D.N. 25(1989)149
Lambertini, F., see Giardini, A. 1(1976)465
Lanari, D. and Rioni, M.
Rate of growth, feed utilisation and digestibility observed with diets containing ensiled maize grain or maize ears fed alone or in mixtures 1(1976)401
Lanari, D., see Rioni, M. 1(1976)531
Lanari, D.
Discussion on session 5 1(1976)545
Landis, K.M., see Hardin, A.C. 25(1989)99
Langar, P.N., Bakshi, M.P.S. and Rana, V.K.
Nutritive evaluation of uromol–wheat straw as a basal ration for buffaloes 13(1985)103
Lanham, J.K., see Coppock, C.E. 18(1987)89
Larsson, K., see Göransson, L. 7(1982)11
Lau, M.M. and van Soest, P.J.
Titratable groups and soluble phenolic compounds as indicators of the digestibility of chemically treated roughages 6(1981)123
Laurent, F., Blanchart, G. and Vignon, B.
Effects of ethanolamine and ethylenediamine treatments on the composition and degradability of wheat straw 13(1985)131
Law, D.P., see Minson, D.J. 9(1983)221
Lawlor, M.J. and O'Shea, J.
The effect of ammoniation on the intake and nutritive value of straw 4(1979)169
Lawrence, T.L.J.
Effects of micronization on the digestibility of whole soya beans and rapeseeds for the growing pig 3(1978)179
Lawrence, T.L.J.
The effects of cereal particle size and pelleting on the nutritive value of oat-based diets for the growing pig 8(1983)91
Le Dividich, J., Christon, R., Peiniau, J. and Aumaitre, A.
Proximate chemical analysis of final cane molasses and effect of feeding 30% molasses on intestinal sucrase and maltase activities in the rat 3(1978)15
Lechtenberg, V.L., see Moore, K.J. 17(1987) 109
Leclercq, B., see Lessire, M. 7(1982)365
Leclercq, B., see Mohamed, K. 11(1984)199
Lee, B.D., see Leeson, S. 11(1984)211
Lee, N.-H., Rooke, J.A. and Armstrong, D.G.
The digestion by sheep of barley and maize-based diets containing either meat and bone meal or soya bean meal 15(1986)301
Lee, N.H., see Ørskov, E.R. 8(1983)247
Leeson, S., see Summers, J.D. 3(1978)243
Leeson, S., see Slinger, S.J. 3(1978)251
Leeson, S., see Shen, H. 8(1983)303
Leeson, S., Summers, J.D. and Lee, B.D.
Nutritive value of single-cell protein produced by *Chaetomium cellulolyticum* grown on maize stover and pulp-mill sludge 11 (1984)211
Leeson, S., see Shen, H. 13(1985)57
Leeson, S., see Peer, D.J. 13(1985)183
Leeson, S., see Peer, D.J. 13(1985)191
Leeson, S., see Peer, D.J. 13(1985)203
Leeson, S., see Summers, J.D. 16(1986/1987) 129
Leeson, S., Hussar, N. and Summers, J.D.
Feeding and nutritive value of hominy and corn grits for poultry 19(1988)313
Leibholz, J., see Evans, D.F. 4(1979)33
Leibholz, J., see Evans, D.F. 4(1979)43
Leibholz, J.
Meat meal in the diet of the early-weaned pig. III. Meat meal quality and the processing of meat meals 4(1979)53
Leibholz, J., Kellaway, R.C. and Hargreave, G.T.
Effects of sodium chloride and sodium bicarbonate in the diet on the performance of calves 5(1980)309
Leibholz, J.
Meat meal in the diet of the early-weaned pig. IV. The supplementation of diets with tryptophane, lysine and methionine 7(1982) 27
Leibholz, J., Love, R.J., Mollah, Y. and Carter, R.R.
The absorption of dietary L-lysine and extruded L-lysine in pigs 15(1986)141
Leibholz, J.M.L., see Kellaway, R.C. 3(1978) 201
Lelong, C.
Maize and animal production in France 1 (1976)141
Lelong, C., see Malterre, C. 1(1976)417
Lelong, C.
For beef cattle, all forms of maize can be used 1(1976)521
Lemerle, C., see Moran, J.B. 20(1988)299
Leng, R.A., see Dryden, G.McL. 14(1986)41
Leng, R.A., see Sudana, I.B. 16(1986/1987)25
Leng, R.A., see Perdok, H.B. 17(1987)121
Leng, R.A., see Dryden, G.McL. 19(1988)121
Lessire, M. and Leclercq, B.
Metabolisable energy value of fats in chicks and adult cockerels 7(1982)365
Letcher, A.J., see Orpin, C.G. 10(1983/1984) 145

Leuillet, M., see Gatel, F. 22(1988)91
Levanon, D., see Silanikove, N. 20(1988)87
Levy, D., see Holzer, Z. 5(1980)299
Levy, D., see Holzer, Z. 12(1984/1985)253
Levy, D., Bar-Tsur, A., Holzer, Z., Samuel, V., Ashbell, G., Lisker, N., Ilan, D., Berkovitz, S. and Kali, J.
High grain content maize silage in fattening diets of young male cattle 16(1986/1987)63
Lewis, A.J., see Aherne, F.X. 3(1978)235
Lewis, B.A., see Jeraci, J.L. 23(1989)15
Lewis, D., see Oguntona, T. 12(1984/1985)285
Lewis, S.M., Holzgraefe, D.P., Berger, L.L., Fahey Jr., G.C., Gould, J.M. and Fanta, G.F.
Alkaline hydrogen peroxide treatments of crop residues to increase ruminal dry matter disappearance in sacco 17(1987)179
Lindberg, J.E., Malmberg, A. and Theander, O.
The chemical composition and nutritive value for ruminants of four possible energy crops and their residues of anaerobic fermentation 15(1986)197
Lindgren, S., see Kaspersson, A. 19(1988)299
Lindner, H.P., see Kirchgessner, M. 6(1981) 337
Linnerud, A.C., see Johnson, T.O. 12(1984/1985)65
Linnerud, A.C., see Ortigues, I. 24(1989)13
Lisker, N., see Levy, D. 16(1986/1987)63
Litsky, J., see Pringle, D. 9(1983)89
Little, D.A., see Mahyuddin, P. 22(1988)69
Livingstone, R.M., Jones, A.S. and Mennie, I.
Swedes (*Brassica napus*) for growing pigs: chemical composition and use as a replacement for barley in the diet 2(1977)31
Livingstone, R.M. and Jones, A.S.
The effect of supplementary protein on the performance of pigs given swedes (*Brassica napus*) as a replacement for barley in the diet 2(1977)277
Livingstone, R.M., Baird, B.A., Atkinson, T. and Crofts, R.M.J.
The effect of different patterns of thermal processing of potatoes on their digestibility by growing pigs 4(1979)295
Livingstone, R.M., Baird, B.A. and Atkinson, T.
Cabbage (*Brassica oleracea*) in the diet of growing–finishing pigs 5(1980)69
Lizama, L.C., Marion, J.E. and McDowell, L.R.
Utilization of aquatic plants *Elodea canadensis* and *Hydrilla verticillata* in broiler chick diets 20(1988)155
Llamas, G.L., see Faulkner, D.B. 12(1984/1985)125
Lo Bruno, A., see Giardini, A. 1(1976)369
Lo Bruno, A., see Giardini, A. 1(1976)465
Lohan, O.P., see Lall, D. 9(1983)29
Longland, A.C. and Low, A.G.
Digestion of diets containing molassed or plain sugar-beet pulp by growing pigs 23 (1989)67
Lord, K.A., Cayley, G.R. and Lacey, J.
Laboratory application of preservatives to hay and the effects of irregular distribution on mould development 6(1981)73
Lord, K.A., see Lacey, J. 6(1981)323
Lord, K.A., Lacey, J. and Cayley, G.R.
Laboratory tests of 8-quinolinol as an additive to volatile fatty acids for preserving hay and other feed-stuffs 7(1982)287
Lord, K.A., see Lacey, J. 8(1983)283
Lough, D.S., see Staples, C.R. 23(1989)277
Love, R.J., see Leibholz, J. 15(1986)141
Low, A.G.
Secretory response of the pig gut to non-starch polysaccharides 23(1989)55
Low, A.G., see Longland, A.C. 23(1989)67
Lowry, J.B., see Mahyuddin, P. 22(1988)69
Lüdke, H. and Schöne, F.
Copper and iodine in pig diets with high glucosinolate rapeseed meal. I. Performance and thyroid hormone status of growing pigs fed on a diet with rapeseed meal treated with copper sulphate solution or untreated and supplements of iodine, copper or a quinoxaline derivative 22(1988)33
Lüdke, H., see Schöne, F. 22(1988)45
Lund, D. and Håkansson, J.
Nutritional and growth studies with pea-crop meals and peas for growing–finishing pigs 16(1986/1987)119
Lutfi, A.A.A., see Maglad, M.A. 15(1986)69
Luthra, Y.P., see Das, B. 4(1979)17
Lynch, G.L., Berger, L.L., Fahey Jr., G.C. and Baker, E.C.
Effects of isopropyl alcohol treatment of soya bean meal on nitrogen utilization by ruminants 14(1986)127

M

Macdearmid, A., see Ørskov, E.R. 6(1981)273
Macdearmid, A., see Ørskov, E.R. 6(1981)355
Macdearmid, A., see Williams, P.E.V. 10 (1983/1984)247
Macdearmid, A., see Williams, P.E.V. 12 (1984/1985)1
Machin, D.H., Hector, D.A., Capper, B.S. and Carter, P.M.
The utilization by broiler chickens of poultry offal hydrolysed in formic acid 11(1984) 247
Machin, D.H., Silverside, D.E., Hector, D.A. and Parr, W.H.
The utilisation by growing pigs of ruminant offal hydrolysed in formic acid 15(1986)273
Mackay, B.E., see Walker, H.G. 9(1983)283
Mackie, I.M.
General review of fish protein hydrolysates 7(1982)113
Mackie, I.M., see Ritchie, A.H. 7(1982)125
Mackie, R.I., see Westlake, K. 25(1989)169

MacLean, S.J., see Gregory, K.F. 2(1977)7
Macleod, G.K., see Smith, O.B. 5(1980)77
Macleod, G.K., see Summers, J.D. 5(1980)205
Madasci, J.P., Parrish, F.W. and McNaughton, J.L.
Treatment of low-tannin sorghum grain for broiler feed 20(1988)69
Madhisetty, M., see Belyea, R.L. 17(1987)57
Maestre, M.R., see Alibés, X. 8(1983)63
Maghoub, S.O., see El Hag, M.G. 13(1985)147
Maglad, M.A., Lutfi, A.A.A. and Gabir, S.
The effect of grinding groundnut hulls either with or without alkali treatment on digestibility of diet and on ruminal and blood components 15(1986)69
Mahyuddin, P., Little, D.A. and Lowry, J.B.
Drying treatment drastically affects feed evaluation and feed quality with certain tropical forage species 22(1988)69
Makkar, H.P.S., Lall, D. and Negi, S.S.
Complexes of urea and formaldehyde as non-protein nitrogen compounds in ruminant rations: a review 20(1988)1
Makkar, H.P.S., see Negi, S.S. 22(1988)79
Malmberg, A., see Lindberg, J.E. 15(1986)197
Malossini, F., see Pace, V. 11(1984)125
Malterre, C. and Lelong, C.
Utilising the maize crop in various forms for beef production 1(1976)417
Malterre, C., see Geay, Y. 1(1976)429
Mandel, E., see Alumot, E. 2(1977)77
Mandell, I.B., Christison, G.I., Nicholson, H.H. and Coxworth, E.C.
The effect of variation in the water content of wheat straw before ammoniation on its nutritive value for beef cattle 20(1988)111
Manfredini, M. and Cavani, C.
Distillery effluents as animal feed: the use of condensed beet molasses stillage (CBMS) in broiler feeding 5(1980)233
Mann, M.E., Cohen, R.D.H., Kernan, J.A., Nicholson, H.H., Christensen, D.A. and Smart, M.E.
The feeding value of ammoniated flax straw, wheat straw and wheat chaff for beef cattle 21(1988)57
Mann, S.O., see Mould, F.L. 10(1983/1984)15
Mann, S.P., see Orpin, C.G. 21(1988)161
Manor, G., see Horn, G.W. 8(1983)35
Mansbridge, R., see Ibbotson, C.F. 10(1983/1984)223
Manten, A.A. and Greenhalgh, J.F.D.
Titles of scientific papers 2(1977)1
Manten, A.A.
Citation decay in animal science 2(1977)195
Mariani, P., see Russo, V. 1(1976)25
Marion, J.E., see Lizama, L.C. 20(1988)155
Marlett, J.A.
Measuring dietary fiber 23(1989)1
Marquardt, R.R., see Rotter, B.A. 24(1989)233
Marti, J., see Cervera, C. 12(1984/1985)233
Martinez Pascual, J. and Fernández Carmona, J.
Composition of citrus pulp 5(1980)1
Martinez Pascual, J. and Fernández Carmona, J.
Citrus pulp in diets for fattening lambs 5 (1980)11
Martinez Pascual, J. and Fernández Carmona, J.
Citrus pulp in diets for fattening rabbits 5 (1980)23
Martinez Pascual, J.L., see Deltoro López, J. 6(1981)163
Martinez Pascual, J.L., see Deltoro López, J. 6(1981)169
Martz, F.A., see Belyea, R.L. 17(1987)57
Mascarenhas Ferreira, A., Kerstens, J. and Gast, C.H.
The study of several modifications of the neutral detergent fibre procedure 9(1983)19
Mascarenhas Ferreira, A., see Van Soest, P.J. 10(1983/1984)155
Mascarenhas Ferreira, A., see Dias-da-Silva, A.A. 19(1988)67
Mascarenhas-Ferreira, A., Guedes, C.V.M. and Dias-da-Silva, A.A.
Effects of urea treatment on chemical composition and in vitro digestibility on meadow hays of Northern Portugal 25(1989) 157
Maslowska, J. and Bazylak, G.
Thermofractographic determination of monocarbonyl compounds in animal waste fats used as feed fat 13(1985)227
Mason, V.C., see Tetlow, R.M. 18(1987)257
Mason, V.C., see Tetlow, R.M. 18(1987)271
Mason, V.C., see Deschard, G. 18(1987)283
Mason, V.C., see Deschard, G. 19(1988)55
Mason, V.C., Hartley, R.D., Keene, A.S. and Cobby, J.M.
The effect of ammoniation on the nutritive value of wheat, barley and oat straws. I. Changes in chemical composition in relation to digestibility in vitro and cell wall degradability 19(1988)159
Mason, V.C., Keene, A.S., Cook, J.E., Cooper, E.M. and Hartley, R.D.
Oven and stack ammoniation of grass hays. 1. Changes in chemical composition in relation to digestibility in vitro and cell-wall degradability 24(1989)299
Mason, V.C., Cook, J.E., Cooper, E.M., Hoadley, C.J. and Cockburn, J.E.
Oven and stack ammoniation of grass hays. 2. Voluntary intake, digestibility and faecal nitrogen excretion in sheep in relation to hay composition 24(1989)313
Mason, W., see Flipot, P. 11(1984)35
Masoud, S., see Abouheif, M.A. 13(1985)215
Mathison, G.W., see Gibson, D.M. 19(1988) 203
May, P.J. and Barker, D.J.
Milling barley and lupin grain in diets for cattle 12(1984/1985)57
Mbatya, P.B.A., see Petchey, A.M. 2(1977)315
Mbatya, P.B.A., Kay, M. and Smart, R.I.

Methods of improving the utilization of cereal straw by ruminants. I. Supplements of urea, molasses and dried grass and treatment with sodium hydroxide 8(1983)221
Mbatya, P.B.A., Kay, M., Smart, R.I. and Milne, J.
Methods of improving the utilization of cereal straw by ruminants. II. Further studies on dried grass as a supplement for lambs 8 (1983)229
Mbatya, P.B.A.
Methods of improving the utilization of cereal straw by ruminants. III. A note on the effect of ensiling straw treated with urea 9 (1983)181
Mbatya, P.B.A., Kay, M., Smart, R.I. and Kennedy, S.
Methods of improving the utilization of cereal straw by ruminants. IV. The effect on intake of level of supplementation with urea and dried grass 13(1985)281
Mbatya, P.B.A., Kay, M., Smart, R.I. and Kennedy, S.
Methods of improving the utilization of cereal straw by ruminants. V. The effect of molasses and dried grass 13(1985)293
Mbugi, P.K., Ingalls, J.R. and Sharma, H.R.
Evaluation of pea protein concentrate as a source of protein in milk replacers for Holstein calves 24(1989)267
McAllan, A.B. and Griffith, E.S.
The effects of different sources of nitrogen supplementation on the digestion of fibre components in the rumen of steers 17(1987) 65
McBurney, M.I. and Thompson, L.U.
Dietary fiber and energy balance: integration of the human ileostomy and in vitro fermentation models 23(1989)261
McCarthy, D.B., see Bunting, L.D. 22(1988) 61
McCarthy, R.E., see Salyers, A.A. 23(1989) 109
McClure, W.H., see Ortigues, I. 24(1989)13
McCrae, F., see Farrell, D.J. 9(1983)49
McDonald, P., see Henderson, A.R. 7(1982) 303
McDonald, P., Proven, M.J. and Henderson, A.R.
The effect of some pre-ensiling treatments on silage composition and nitrogen disappearance in the rumen 8(1983)259
McDonald, P., see Heron, S.J.E. 19(1988)85
McDowell, L.R., see Paz, E.A. 11(1984)75
McDowell, L.R., see Lizama, L.C. 20(1988)155
McEllhiney, R.R. and Tangprasertchai, P.
The effect of dilution levels in premixes on micro-ingredient dispersion in animal feeds 8(1983)139
McGilliard, A.D., see Adamu, A.M. 22(1988) 227
McGinn, R., see McLellan, A.R. 6(1981)245
McGinn, R., see McLellan, A.R. 9(1983)107
McGinn, R. and McLellan, A.R.
Energy metabolism in lambs fed silage diets with sucrose or a growth promoter 16(1986/1987)203
McIntosh, A.D., see Davidson, J. 7(1982)217
McKenzie, R.J., see Farrell, D.J. 9(1983)49
McKirdy, J.A., see Sharma, H.R. 10(1983/1984)77
McLaughlin, G.L., see Horn, G.W. 8(1983)35
McLellan, A.R. and McGinn, R.
A calorimetric investigation of the nutritive value of formic acid- and formaldehyde–formic acid-treated grass silages 6(1981)245
McLellan, A.R. and McGinn, R.
Energy and nitrogen balance, intake and growth rate of lambs fed on unwilted grass silages treated with a formaldehyde–acid mixture or untreated 9(1983)107
McLellan, A.R., see McGinn, R. 16(1986/1987)203
McLeod, M.N. and Minson, D.J.
The analytical and biological accuracy of estimating the dry matter digestibility of different legume species 1(1976)61
McLeod, M.N. and Minson, D.J.
The accuracy of the pepsin–cellulase technique for estimating the dry matter digestibility in vivo of grasses and legumes 3 (1978)277
McLeod, M.N. and Minson, D.J.
A note on Onozuka 3S cellulase as a replacement for Onozuka SS (P1500) cellulase when estimating forage digestibility in vitro 5(1980)347
McLeod, M.N. and Minson, D.J.
Accuracy of predicting digestibility by the cellulase technique; the effect of pretreatment of forage samples with neutral detergent or acid pepsin 7(1982)83
McLeod, M.N. and Wilson, J.R.
The effect of method of preservation of forages on their resistance to reducion in particle size with an artificial masticator 24 (1989)185
McMeniman, N.P. and Armstrong, D.G.
Nitrogen levels in low-roughage diets for efficient rumen microbial protein production 2(1977)255
McMeniman, N.P., Elliott, R. and Ash, A.J.
Supplementation of rice straw with crop by-products. I. Legume straw supplementation 19(1988)43
McNaughton, J.L., see Madasci, J.P. 20(1988) 69
McNew, R.W., see Zorrilla-Rios, J. 22(1988) 305
McReynolds, W.E., see Rule, D.C. 15(1986) 113
Melcion, J.P., see Lacassagne, L. 20(1988)59
Mennie, I., see Livingstone, R.M. 2(1977)31
Mensah-Asante, B., see Donkoh, A. 22(1988) 139
Mergounis, D.
Use of maize for ruminants and pig feeding in Greece 1(1976)125

Merritt, J.H.
Assessment of the production costs of fish protein hydrolysates 7(1982)147
Metwally, H., see Abouheif, M.A. 13(1985)215
Meyer, D.W., see Sanderson, M.A. 14(1986) 221
Michalet-Doreau, B., see Aufrere, J. 20(1988) 203
Michie, W., see Cowan, P.J. 4(1979)125
Michie, W., see Rose, S.P. 11(1984)221
Mika, V.
Some remarks on the determination of acid-detergent fibre in feeds with cetyltrimethyl-ammonium bromide versus determination of detergent fibre with alkylarylsulfonic acid 6(1981)91
Miller, C., see Caffrey, P.J. 19(1988)231
Miller, M.E., see Wohlt, J.E. 6(1981)115
Milliken, G., see Hinds, M.A. 12(1984/1985) 205
Milne, E., see Davidson, J. 7(1982)217
Milne, J., see Mbatya, P.B.A. 8(1983)229
Minson, D.J., see McLeod, M.N. 1(1976)61
Minson, D.J., see Goto, I. 2(1977)247
Minson, D.J., see McLeod, M.N. 3(1978)277
Minson, D.J., see McLeod, M.N. 5(1980)347
Minson, D.J.
A flexible, two-component feeding system derived from Blaxter's three-component, metabolisable-energy system 6(1981)223
Minson, D.J., see McLeod, M.N. 7(1982)83
Minson, D.J.
Feeding standards are necessary 7(1982)423
Minson, D.J., Butler, K.L., Grummitt, N. and Law, D.P.
Bias when predicting crude protein, dry matter digestibility and voluntary intake of tropical grasses by near-infrared reflectance 9(1983)221
Mirgani, T.
Effect of sodium hydroxide treatment of desert grass on carbohydrate digestibility by goats 17(1987)219
Miron, J., see Ben-Ghedalia, D. 10(1983/ 1984)269
Miron, J., see Ben-Ghedalia, D. 19(1988)219
Miron, J., see Ben-Ghedalia, D. 24(1989)289
Mitaru, B.N., see Baidoo, S.K. 18(1987)37
Mitaru, B.N., see Baidoo, S.K. 18(1987)45
Mitchell, K.G., see Barber, R.S. 2(1977)161
Mitchell, K.G., see Barber, R.S. 3(1978)163
Mitchell, K.G., see Barber, R.S. 4(1979)117
Mitchell, K.G., see Barber, R.S. 4(1979)233
Mitchell, K.G., see Barber, R.S. 5(1980)215
Mitchell, K.G., see Barber, R.S. 6(1981)35
Mitchell Jr., E.D., see Hibberd, C.A. 15(1986) 231
Mlodkowski, M., see Hanczakowski, P. 6(1981) 413
Mngulwi, J.G.J., see Kategile, J.A. 20(1988) 97
Modler, H.W., see Elliot, J.I. 17(1987)213
Moermans, R., see Eeckhout, W. 6(1981)367
Moermans, R.J., see Aerts, J.V. 3(1978)309
Moermans, R.J., see Cottyn, B.G. 11(1984) 137
Mohamed, K., Leclercq, B., Anwar, A., El-Alaily, H. and Soliman, H.
A comparative study of metabolisable energy in ducklings and domestic chicks 11 (1984)199
Mohammed, T.A., see Osman, A.G. 19(1988) 351
Mohammed, T.A. and Ali, O.M.
Effects of wood ash extract treatment on the feeding value and utilization of high-tannin sorghums by broiler chicks 22(1988)131
Moir, K.W. and Connor, J.K.
A comparison of three fibre methods for predicting the metabolizable energy content of sorghum grain for poultry 2(1977)197
Moisio, T. and Heikonen, M.
A titration method for silage assessment 22 (1988)341
Molina, B.T., see Djajanegara, A. 12(1984/ 1985)141
Molina, E., Boza, J. and Aguilera, J.F.
Nutritive value for ruminants of sugar cane bagasse ensiled after spray treatment with different levels of NaOH 9(1983)1
Molina, E., see Aguilera, J.F. 12(1984/1985) 171
Mollah, Y., see Leibholz, J. 15(1986)141
Mølle, KR. G.
Maize as a forage crop in Denmark 1(1976) 183
Molnar, S., see Greife, H.A. 5(1980)241
Montague Jr., W.C., see Walker, H.G. 9(1983) 283
Monteagudo, A., see Gonzalez, G. 1(1976)129
Monteagudo, A.
Cultivation of maize in irrigated areas for silage and grain production and their use by ruminants 1(1976)237
Montgomery, R.R., Nofsinger, G.W. and Bothast, R.J.
Preservation of high-moisture maize – a comparison of gaseous and liquid anhydrous ammonia with methylene-*bis*-propionate 5 (1980)337
Moo-Young, M. and Chahal, D.S.
Utilization of cattle manure for single-cell protein production with *Chaetomium cellulolyticum* 4(1979)199
Moo-Young, M., Chahal, D.S. and Vlach, D.
Utilization of cattle manure for Torula yeast production from straw hydrolysates 5(1980) 175
Moo-Young, M., see Chavez, E.R. 22(1988)1
Moo-Young, M., see Chavez, E.R. 22(1988)13
Moo-Young, M., see Chavez, E.R. 22(1988)23
Moore, K.J. and Lechtenberg, V.L.
Chemical composition and digestion in vitro of orchardgrass hay ammoniated by different techniques 17(1987)109
Mora, M.I., Fahey Jr., G.C., van der Aar, P.J. and Berger, L.L.
Improving utilization of crop residues. Eval-

uation of two chemicals used to improve digestibility of crop residues 9(1983)205
Moran, J.B., see Thompson, C. 16(1986/1987) 225
Moran, J.B., Lemerle, C. and Trigg, T.E.
The intake and digestion of maize silage-based diets by dairy cows and sheep 20 (1988)299
Morel, M.T., see Durand, M. 21(1988)197
Morgan, C.A. and Whittemore, C.T.
Energy evaluation of feeds and compounded diets for pigs – a review 7(1982)387
Morgan, C.A., Whittemore, C.T. and Cockburn, J.H.S.
The effect of level and source of protein, fibre and fat in the diet on the energy value of compounded pig feeds 11(1984)11
Morgan, C.A., Whittemore, C.T., Phillips, P. and Crooks, P.
The prediction of the energy value of compounded pig foods from chemical analysis 17(1987)81
Morgan, C.A. and Whittemore, C.T.
Dietary fibre and nitrogen excretion and retention by pigs 19(1988)185
Morgan, C.A., Whittemore, C.T. and Taylor, A. G.
Body composition and growth of young pigs as affected by protein source and a growth promoter 24(1989)219
Morita, Z., see Sekine, J. 25(1989)207
Morris, J.G., see Sidahmed, A.E. 8(1983)47
Morrison, I.M. and Brice, R.E.
The digestion of untreated and ammonia-treated barley straw in an artificial rumen 10(1983/1984)229
Morrison, I.M., see Robinson, P.H. 18(1987)75
Mosi, A., see Butterworth, M.H. 16(1986/1987)99
Mosi, A.K. and Butterworth, M.H.
The voluntary intake and digestibility of combinations of cereal crop residues and legume hay for sheep 12(1984/1985)241
Moss, B.R., El-Negoumy, A.M. and Newman, C.W.
Nutritional value of waxy barleys for broiler chickens 8(1983)25
Moughan, P.J., see Kies, A.K. 16(1986/1987) 169
Moughan, P.J., see Smith, W.C. 18(1987)143
Moughan, P.J., see Smith, W.C. 19(1988)105
Moughan, P.J., Wilson, M.N., Smits, C.H.M. and Smith, W.C.
An evaluation of skim milk powders subjected to different heating conditions during processing as dietary protein sources for the young pig 22(1988)203
Mould, F.L. and Ørskov, E.R.
Manipulation of rumen fluid pH and its influence on cellulolysis in sacco, dry matter degradation and the rumen microflora of sheep offered either hay or concentrate 10 (1983/1984)1
Mould, F.L., Ørskov, E.R. and Mann, S.O.
Associative effects of mixed feeds. I. Effects of type and level of supplementation and the influence of the rumen fluid pH on cellulolysis in vivo and dry matter digestion of various roughages 10(1983/1984)15
Mould, F.L., Ørskov, E.R. and Gauld, S.A.
Associative effects of mixed feeds. II. The effect of dietary addition of bicarbonate salts on the voluntary intake and digestibility of diets containing various proportions of hay and barley 10(1983/1984)31
Mowat, D.N., see Oji, U.I. 4(1979)177
Mowat, D.N., see Oji, U.I. 4(1979)187
Mshelia, T.A., see Kumaresan, A. 11(1984)45
Mugerwa, J.S., see Odwongo, W.O. 5(1980) 193
Muindi, P.J., Thomke, S. and Hanssen, J.F.
Protein quality studies on rats fed on cassava-root meal enriched with *Cephalosporium eichhorniae* 152 or with conventional plant-protein supplements 6(1981)197
Müller, H.L., see Kreuzer, M. 16(1986/1987) 233
Muñoz, F., see Alibés, X. 8(1983)63
Muñoz, F., see Alibés, X. 10(1983/1984)239
Muñoz, F., see Alibes, X. 11(1984)189
Murdock, F.R. and Kromann, R.P.
The digestibility and nutritive value for dairy heifers of rations containing cellulose fibre 2(1977)287
Murphy, G.E., see Hardin, A.C. 25(1989)99
Mussa, M.A., see Sarwatt, S.V. 22(1988)237
Myer, R.O. and Gorbet, D.W.
Waxy and normal grain sorghums with varying tannin contents in diets for young pigs 12(1984/1985)179
Myer, R.O., Barnett, R.D., Cornell, J.A. and Combs, G.E.
Nutritive value of diets containing triticale and varying mixtures of triticale and maize for growing-finishing swine 22(1988)217
Mzihirwa, Y.G., see Kategile, J.A. 6(1981)133

N

Nangole, F.N., Kayongo-Male, H. and Said, A.N.
Chemical composition, digestibility and feeding value of maize cobs 9(1983)121
Napoli, G.M. and Santini, F.J.
The effect of a protein energy supplement on pasture protein and fibre digestion in the rumen of grazing steers 25(1989)39
Narahari, D. and Kothandaraman, P.
The influence of processing and storage on hydrogen cyanide and tannin contents of para-rubber seed and its products 9(1983) 319
Narahari, D. and Kothandaraman, P.
Chemical composition and nutritional value of para-rubber seed and its products for chickens 10(1983/1984)257
Narasimhalu, P.

A comparison of laboratory methods involving single or two-stage cellulase and two-stage rumen inoculum–pepsin treatment for predicting digestibility and intake of hays in sheep 12(1984/1985)101
Narasimhalu, P., Graham, H. and Åman, P.
Effects of ozonation on the chemical composition and in vitro degradation of triticale internodes and red clover stems 24(1989)159
Natanam, R., Kadirvel, R. and Balagopal, R.
The effect of kernels of karanja (*Pongamia glabra* Vent) on growth and feed efficiency in broiler chicks to 4 weeks of age 25(1989) 201
Neale, R.J., see Oguntona, T. 12(1984/1985) 285
Nefzaoui, A. and Vanbelle, M.
Effects of feeding alkali-treated olive cake on intake, digestibility and rumen liquor parameters 14(1986)139
Negi, S.S.
Tannins in sal seed (*Shorea robusta*) and sal seed meal limit their utilization as livestock feeds 7(1982)161
Negi, S.S., see Lall, D. 9(1983)29
Negi, S.S., see Makkar, H.P.S. 20(1988)1
Negi, S.S., Singh, B. and Makkar, H.P.S.
Rumen degradability of nitrogen in typical cultivated grasses and leguminous fodders 22(1988)79
Neill, A.R., Bell, S.L. and Williams, K.C.
The dye absorption test as an indicator of the nutritive quality of heat-processed navy bean 9(1983)71
Nelson, M.L. and Finley, J.W.
Effect of soft white wheat addition to alfalfa-grass forage on heifer gain, diet digestibility and in vitro digestion kinetics 24 (1989)141
Nes, I.F., see Skrede, A. 20(1988)287
Neskar, M., see Rotter, B.A. 24(1989)233
Newman, C.W., see Moss, B.R. 8(1983)25
Nicholson, H.H., see Horton, G.M.J. 7(1982)1
Nicholson, H.H., see Mandell, I.B. 20(1988) 111
Nicholson, H.H., see Mann, M.E. 21(1988)57
Nikolić, J.A. and Jovanović, M.
Some properties of apple pomace ensiled with and without additives 15(1986)57
Njoku, P.C., see Okoh, P.N. 7(1982)359
Njoku, P.C.
Effect of dietary ascorbic acid (vitamin C) supplementation on the performance of broiler chickens in a tropical environment 16 (1986/1987)17
Nofsinger, G.W., see Montgomery, R.R. 5 (1980)337
Noland, P.R., Campbell, D.R., Sharp, R.N. and Johnson, Z.B.
Influence of pericarp and endosperm colour and type on digestibility of grain sorghum by pigs 2(1977)219
Noland, P.R., see Sharp, R.N. 2(1977)327
Noland, P.R., Campbell, D.R. and Johnson, Z.B.
Use of unextracted sunflower seeds as a protein source 5(1980)51
Nordkvist, E., Graham, H. and Åman, P.
Degradation in vitro and in sacco of red clover leaves and stems 17(1987)295
Nwokolo, E. and Oji, U.I.
Variation in metabolizable energy content of raw or autoclaved white and brown varieties of three tropical grain legumes 13 (1985)141
Nwokolo, E.
A comparison of metabolizable energy content of eight common feed ingredients determined with young guinea fowls (keets) and pullet chicks 15(1986)1

O

Obilana, A.B., see Alhassan, W.S. 17(1987) 285
Obilana, A.T., see Okoh, P.N. 7(1982)359
Ocio, E., Viñaras, R. and Rey, J.Ma.
House fly larvae meal grown on municipal organic waste as a source of protein in poultry diets 4(1979)227
Odebunmi, J.O., see Babatunde, G.M. 4 (1979)23
Oduye, O.O., see Babatunde, G.M. 4(1979)23
Odwongo, W.O. and Mugerwa, J.S.
Performance of calves on diets containing *Amaranthus* leaf meal 5(1980)193
Offer, N.W., see Husain, R.A.K. 16(1986/ 1987)297
Offer, N.W. and Husain, R.A.K.
Fish silage as a protein supplement for early-weaned calves 17(1987)165
Offer, N.W., see Hyslop, J.J. 25(1989)55
Ogle, R.B., see Göransson, L. 12(1984/1985) 159
Oguntona, T., Neale, R.J. and Lewis, D.
The effect of pelleting on the utilisation of yeast grown on n-alkanes by broiler chicks 12(1984/1985)285
Ohajuruka, O.A. and Palmquist, D.L.
Response of high-producing dairy cows to high levels of dried corn gluten feed 24 (1989)191
Ohlde, G. and Becker, K.
Suitability of cell-wall constituents as predictors of organic matter digestibility in some tropical and subtropical by-products 7(1982)191
Oji, U.I. and Mowat, D.N.
Nutritive value of thermoammoniated and steam-treated maize stover I. Intake, digestibility and nitrogen retention 4(1979) 177
Oji, U.I., Mowat, D.N. and Buchanan-Smith, J.G.
Nutritive value of thermoammoniated and

steam-treated maize stover II. Rumen metabolites and rate of passage 4(1979)187
Oji, U.I., see Nwokolo, E. 13(1985)141
Oke, O.L.
Problems in the use of cassava as animal feed 3(1978)345
Okoh, P.N., Obilana, A.T., Njoku, P.C. and Aduku, A.O.
Proximate analysis, amino acid composition and tannin content of improved Nigerian sorghum varieties and their potential in poultry feeds 7(1982)359
Oldham, J.D., see Phipps, R.H. 4(1979)163
Olsson, O., see Göransson, L. 7(1982)11
Omole, T.A. and Onwudike, O.C.
Investigations of the treatment of sawdust for rabbit feeding. 1. Effect of sodium hydroxide treatment 6(1981)43
Omole, T.A. and Onwudike, O.C.
Investigations of the treatment of sawdust for rabbit feeding. 2. Effect of boiling and incubation with poultry manure 6(1981)51
Omole, T.A., see Sonaiya, E.B. 8(1983)211
Onesirosan, P., see Smith, O.B. 11(1984)231
Onwudike, O.C., see Omole, T.A. 6(1981)43
Onwudike, O.C., see Omole, T.A. 6(1981)51
Onwudike, O.C.
Palm kernel meal as a feed for poultry. 1. Composition of palm kernel meal and availability of its amino acids to chicks 16(1986/1987)179
Onwudike, O.C.
Palm kernel meal as a feed for poultry. 2. Diets containing palm kernel meal for starter and grower pullets 16(1986/1987)187
Onwudike, O.C.
Palm kernel meal as a feed for poultry. 3. Replacement of groundnut cake by palm kernel meal in broiler diets 16(1986/1987) 195
Onwudike, O.C.
Palm kernel meal as a feed for poultry. 4. Use of palm kernel meal by laying birds 20 (1988)279
Opstvedt, J., Søbstad, G. and Hansen, P.
Fish protein concentrates for pre-ruminant calves: effects of processing temperatures and source of fish on protein digestibility and biological value 18(1987)181
Oram, R.N., see Green, A.G. 9(1983)271
Orpin, C.G.
The role of ciliate protozoa and fungi in the rumen digestion of plant cell walls 10(1983/1984)121
Orpin, C.G. and Letcher, A.J.
Effect of absence of ciliate protoza on rumen fluid volume, flow rate and bacterial populations in sheep 10(1983/1984)145
Orpin, C.G., Hazlewood, G.P. and Mann, S.P.
Possibilities of the use of recombinant-DNA techniques with rumen micro-organisms 21 (1988)161
Ørskov, E.R. and Grubb, D.A.
The use of whole barley diets fortified with solutions of urea, minerals and vitamins for lambs 2(1977)307
Ørskov, E.R., Macdearmid, A., Grubb, D.A. and Innes, G.M.
Utilization of alkali-treated grain. 1. Alkali-treated grain in complete diets for steers and lambs 6(1981)273
Ørskov, E.R., see Barnes, B.J. 6(1981)347
Ørskov, E.R., Barnes, B.J., Macdearmid, A., Williams, P.E.V. and Innes, G.M.
Utilization of alkali-treated grain. 3. Utilization by steers of NaOH-treated and rolled barley in silage-based diets 6(1981)355
Ørskov, E.R., Soliman, H.S. and Clark, C.F.S.
Use of fish protein hydrolysate in milk replacers 7(1982)135
Ørskov, E.R., Reid, G.W., Holland, S.M., Tait, C.A.G. and Lee, N.H.
The feeding value for ruminants of straw and whole-crop barley and oats treated with anhydrous or aqueous ammonia or urea 8 (1983)247
Ørskov, E.R., see Mould, F.L. 10(1983/1984) 1
Ørskov, E.R., see Mould, F.L. 10(1983/1984) 15
Ørskov, E.R., see Mould, F.L. 10(1983/1984) 31
Ørskov, E.R., see Istasse, L. 15(1986)167
Ørskov, E.R., see Istasse, L. 16(1986/1987)37
Ørskov, E.R., see Silva, A.T. 19(1988)277
Ørskov, E.R., see Silva, A.T. 19(1988)289
Ørskov, E.R., see Bhargava, P.K. 22(1988)295
Ortigues, I., Crickenberger, R.G., Fontenot, J.P., Harvey, R.W., McClure, W.H. and Linnerud, A.C.
Distribution of liquid supplement injected into large bales of roughage and comparison of nitrogen supplements for stocker cattle fed low-quality roughages 24(1989)13
Osafo, E.L.K., see Smith, O.B. 20(1988)189
Osbourn, D.F.
Principles governing the use of chemical methods for assessing the nutritive value of forages: a review 3(1978)265
Osbourn, D.F., Terry, R.A., Spooner, M.C. and Tetlow, R.M.
Use of processing to explore the factors affecting the digestion of forage cell walls 6 (1981)387
Osei, S.A. and Twumasi, I.K.
Effects of oven-dried cassava peel meal on the performance and carcass characteristics of broiler chickens 24(1989)247
O'Shea, J., see Lawlor, M.J. 4(1979)169
O'Shea, J. and Baldwin, A.
Evaluation of the increase in feeding value of barley straw following mild treatment with sulphur dioxide 15(1986)13
Oslage, H.J., see Schulz, E. 1(1976)9
Osman, A.G., El Tayeb, A.E., Sulieman, A.H. and Mohammed, T.A.
Effect of sorghum straw alone or in combination with molasses and nitrogen sources

on performance of Sudan desert lambs 19 (1988)351
Otwinowska, A., see Kowalczyk, J. 7(1982)225
Oura, R., see Sekine, J. 25(1989)207
Ovejero, F.M.A., see Fay, J.P. 16(1986/1987) 161
Overend, M.A., see Rooke, J.A. 13(1985)255
Owen, B.D., see Beames, R.M. 15(1986)285
Owen, E., see Alawa, J.P. 11(1984)149
Owen, E., see Alawa, J.P. 12(1984/1985)215
Owen, E., see Wrathall, J.H.M. 24(1989)57
Owens, F.N., see Goetsch, A.L. 18(1987)151

P

Pace, V., Barge, M.T., Settineri, D. and Malossini, F.
Comparison of forage digestibility in vitro with enzymic solubility 11(1984)125
Pahlow, G., see Weinberg, Z.G. 20(1988)335
Palmer, A.Z., see Potter, S.G. 12(1984/1985) 275
Palmquist, D.L., see Ohajuruka, O.A. 24 (1989)191
Pandit, N.N., see Singh, A.P. 3(1978)299
Pandita, N.N., see Singh, A.P. 9(1983)143
Panditharatne, S., see Jayasuriya, M.C.N. 3 (1978)219
Panigrahi, S., Bland, B. and Carlaw, P.M.
The nutritive value of tamarind seeds for broiler chicks 22(1988)285
Papadopoulos, G. and Ziras, E.
Nutrient composition of Greek cottonseed meal 18(1987)295
Papadopoulos, G., Fegeros, K. and Ziras, E.
Evaluation of Greek cottonseed meal. 2. Use in rations for fattening pigs 18(1987)303
Papadopoulos, M.C., El Boushy, A.R., Roodbeen, A.E. and Ketelaars, E.H.
Effects of processing time and moisture content on amino acid composition and nitrogen characteristics of feather meal 14(1986) 279
Papadopoulos, M.C.
The effect of enzymatic treatment on amino acid content and nitrogen characteristics of feather meal 16(1986/1987)151
Parigi-Bini, R. and Chiericato, G.M.
Influence of ensiling maize ears and grain on chemical composition, conservation losses and digestibility 1(1976)301
Parkins, J.J., see Bass, J.M. 7(1982)93
Parkins, J.J., see Hemingway, R.G. 15(1986) 123
Parr, W.H., see Machin, D.H. 15(1986)273
Parrish, F.W., see Madasci, J.P. 20(1988)69
Parthasarathy, M. and Pradhan, K.
Fermentation characteristics and feeding value of ensiled poultry litter containing wheat straw, bagasse or sawdust 7(1982)341
Partridge, G.G., Allan, S.J. and Findlay, M.
Studies on the nutritive value of roots, cabbage and grass silage for growing commercial rabbits 13(1985)299
Partridge, G.G., Findlay, M. and Fordyce, R.A.
Fat supplementation of diets for growing rabbits 16(1986/1987)109
Partridge, I.G., see Barber, R.S. 4(1979)233
Partridge, I.G., see Barber, R.S. 5(1980)215
Partridge, I.G., see Barber, R.S. 6(1981)35
Partridge, I.G.
The digestion of diets containing manioc by young growing pigs 12(1984/1985)119
Patchell, M.R., see Smith, W.C. 19(1988)105
Paterson, J.A., see Koeln, L.L. 11(1984)181
Paterson, J.A., see Goedeken, F.K. 13(1985) 47
Pathak, N.N., see Jain, V.K. 5(1980)315
Pathak, N.N., see Kamra, D.N. 25(1989)149
Patterson, D.C. and Walker, N.
The use of effluent from grass silage in the diet of finishing pigs. I. Variation in composition of effluents 4(1979)263
Patterson, D.C. and Walker, N.
The use of effluent from grass silage in the diet of finishing pigs. II. Assessment of nutritive value of fresh and stored effluent 4 (1979)275
Payne, E., see Wright, D.E. 2(1977)93
Paz, E.A., McDowell, L.R., Conrad, J.H. and Damron, B.L.
Biological availability of phosphorus from basic steel slag 11(1984)75
Pearce, G.R., see Hendrosoekarjo, S. 3(1978) 31
Pearce, G.R., see Stanogias, G. 3(1978)155
Pearce, G.R., see Jayasuriya, M.C.N. 8(1983) 271
Pearson, A.W., see Fenwick, G.R. 6(1981)421
Pearson, A.W., Greenwood, N.M., Butler, E.J., Curl, C.L. and Fenwick, G.R.
The involvement of trimethylamine oxide in fish meal in the production of egg taint 8 (1983)119
Pearson, G. and Carr, J.R.
Lupin-seed meal (*Lupinus angustifolius* cv. Uniwhite) as a protein supplement to barley-based diets for growing pigs 1(1976) 631
Pearson, G. and Carr, J.R.
A comparison between meals prepared from the seeds of different varieties of lupin as protein supplements to barley-based diets for growing pigs 2(1977)49
Pearson, G., see Smith, W.C. 18(1987)143
Pearson, G., see Smith, W.C. 19(1988)105
Peck, N.J., see Ashes, J.R. 3(1978)109
Peck, N.J., see Farrell, D.J. 9(1983)99
Pedroza, C., see Guardiola, C.M. 8(1983)129
Peer, D.J. and Leeson, S.
Feeding value of hydroponically sprouted barley for poultry and pigs 13(1985)183
Peer, D.J. and Leeson, S.
Nutrient content of hydroponically sprouted barley 13(1985)191
Peer, D.J. and Leeson, S.

Nutrient and trypsin inhibitor content of hydroponically sprouted soya beans 13 (1985)203
Peers, D.G., Taylor, A.G. and Whittemore, C.T.
The influence of feeding level and level of dietary inclusion on the digestibility of barley meal in the pig 2(1977)41
Peiniau, J., see Le Dividich, J. 3(1978)15
Peiniau, J., see Aumaitre, A. 24(1989)275
Pendlum, L.C., Boling, J.A. and Ely, D.G.
Abomasal and plasma nitrogen constituents of steers fed on different physical forms of shelled maize 2(1977)225
Penning, P.D., Corcuera, P. and Treacher, T.T.
Effect of dry-matter concentration of milk substitute and method of feeding on intake and performance by lambs 5(1980)321
Perdok, H.B., see Jayasuriya, M.C.N. 7(1982) 201
Perdok, H.B. and Leng, R.A.
Hyperexcitability in cattle fed ammoniated roughages 17(1987)121
Perera, H.G.D., see Jayasuriya, M.C.N. 7 (1982)221
Perez, J.M. and Aumaitre, A.
Waxy versus regular maize:energy value for growing pigs and utilization in piglet diets 4(1979)109
Petchey, A.M. and Mbatya, P.B.A.
The intake of steers and heifers fed on mixtures of alkali-treated or untreated straw with grass silage 2(1977)315
Petchey, A.M.
Performance of calves fed on milk replacers containing fish protein hydrolysate 7(1982) 141
Peterson, C.L., see Thomas, V.M. 11(1984)269
Peteva-Vantcheva, Z. and Platikanov, N.
Forage maize in the rations of dairy cows and fattening cattle in Bulgaria 1(1976)45
Peteva-Vantcheva, Z. and Jelev, K.
Maize silage and whole maize plant pellets as the basic forages in complete feeds for milk cows 3(1978)23
Pettersson, D., Hesselman, K. and Åman, P.
Nutritional value for chickens of dried distillers-spent-grain from barley and dehulled barley 17(1987)145
Pettersson, D. and Åman, P.
Effects of enzyme supplementation of diets based on wheat, rye or triticale on their productive value for broiler chickens 20(1988) 313
Petzel, M., see Hanczakowski, P. 4(1979)91
Pfeffer, E., see Abate, A. 14(1986)1
Pflugfelder, R.L., Rooney, L.W. and Schake, L.M.
The role of germination in sorghum reconstitution 14(1986)243
Phillips, P., see Morgan, C.A. 17(1987)81
Phipps, R.H. and Weller, R.F.
Effect of agronomic factors on the yield of maize and on its composition in relation to its use by ruminants 1(1976)251
Phipps, R.H. and Oldham, J.D.
Amino acid composition of maize silage produced in the United Kingdom 4(1979)163
Pike, D.J., see Wrathall, J.H.M. 24(1989)57
Pirie, R. and Greenhalgh, J.F.D.
Alkali treatment of straw for ruminants. I. Utilization of complete diets containing straw by beef cattle 3(1978)143
Pirie, R., see Greenhalgh, J.F.D. 3(1978)289
Pittman, R.J., see Barber, R.S. 3(1978)163
Pittman, R.J., see Barber, R.S. 4(1979)233
Pittman, R.J., see Barber, R.S. 5(1980)215
Platikanov, N., see Peteva-Vantcheva, Z. 1 (1976)45
Playne, M.J.
Differences between cattle and sheep in their digestion and relative intake of a mature tropical grass hay 3(1978)41
Playne, M.J.
Estimation of the digestibility of low-quality hays by cattle from measurements made with sheep 3(1978)51
Pollmann, D.S., Behnke, K.C. and Allee, G.L.
The effect of diet complexity and physical form on the performance of starter pigs and feed processing efficiency 8(1983)99
Pollmann, D.S. and Bandyk, C.A.
Stability of viable lactobacillus products 11 (1984)261
Pond, W.G., see Babatunde, G.M. 20(1988) 125
Potter, S.G., Ammerman, C.B., Henry, P.R., Becker, H.N. and Palmer, A.Z.
Effect of sugarcane condensed molasses solubles, sugarcane molasses and monensin on performance and volatile fatty acid production in finishing steers 12(1984/1985)275
Poutiainen, E., see Pulli, S. 1(1976)187
Poyyamozhi, V.S. and Kadirvel, R.
The value of banana stalk as a feed for goats 15(1986)95
Pradhan, K., see Parthasarathy, M. 7(1982) 341
Prasad, C.M., see Singh, K.S. 4(1979)143
Prasad, C.M., see Singh, K.S. 6(1981)63
Prasad, D.A., see Chandra, S. 12(1984/1985) 187
Prasad, J., see Singh, A.P. 9(1983)325
Prasad, T., see Lall, D. 23(1989)343
Preston, R.L., see Rule, D.C. 15(1986)113
Prieto, C., see Fonolla, J. 6(1981)405
Prieto, C., see Aguilera, J.F. 12(1984/1985) 171
Prieto, C., see Aguilera, J.F. 15(1986)33
Prieto, F., see Szylit, O. 3(1978)73
Pringle, D., Beames, R.M., Tait, R.M. and Litsky, J.
Effect of storage and processing of barley on its utilization by pigs and rats 9(1983)89
Pringle, D.B., see Beames, R.M. 15(1986)285
Proudfoot, F.G., see Zarkadas, C.G. 14(1986)

291
Proudfoot, F.G., see Hulan, H.W. 16(1986/1987)253
Proven, M.J., see McDonald, P. 8(1983)259
Pulli, S., Poutiainen, E. and Syrjala, L.
Prospects of growing maize in Finland 1 (1976)187
Pulungan, H., see Van Eys, J.E. 18(1987)197
Punj, M.L., see Seghal, J.P. 9(1983)155
Punj, M.L., see Jakhmola, R.C. 19(1988)1
Pusztai, A.J., see Williams, P.E.V. 12(1984/1985)1
Pyle, C., see Wright, D.E. 2(1977)93

R

Radosevich, S.R., see Sidahmed, A.E. 8(1983)47
Raharjo, Y. and Farrell, D.J.
A new biological method for determining amino acid digestibility in poultry feedstuffs using a simple cannula, and the influence of dietary fibre on endogenous amino acid output 12(1984/1985)29
Raj Reddy, M., see Krishna Reddy, G.V. 24 (1989)69
Rajaguru, A.S.B., see Ravindran, V. 17(1987)227
Rakes, J.M., see Daniels, L.B. 8(1983)19
Rana, V.K., see Langar, P.N. 13(1985)103
Rangkuti, M., see Van Eys, J.E. 18(1987)197
Rangnekar, D.V., Badve, V.C., Kharat, S.T., Sobale, B.N. and Joshi, A.L.
Effect of high-pressure steam treatment on chemical composition and digestibility in vitro of roughages 7(1982)61
Rao, N.M. and Fontenot, J.P.
Ensiling of rumen contents and blood with wheat straw 18(1987)67
Ravindran, V., Kornegay, E.T. and Rajaguru, A.S.B.
Influence of processing methods and storage time on the cyanide potential of cassava leaf meal 17(1987)227
Razzaque, M.A., Al-Nasser, A., Salman, A.J., Aderibigbe, A.O. and Church, D.C.
Corrugated cardboard and dried poultry manure as dietary ingredients for sheep 14 (1986)265
Razzaque, M.A., see Ilian, M.A. 19(1988)327
Razzaque, M.A., see Ilian, M.A. 19(1988)343
Reade, A.E., see Gregory, K.F. 2(1977)7
Rebolé, A. and Alvira, P.
Composition of vine-branches with leaves of *Vitis vinifera* L. fresh and ensiled using different additives (previous results) 16(1986/1987)89
Rebolé, A., Alvira, P. and González, G.
Digestibility in vivo of ensiled grapevines (branches and leaves): influence of the system of analysis in the detergent fibre scheme on the prediction of digestibility 22(1988)173
Rechkemmer, G., see Von Engelhardt, W. 23 (1989)43
Reddy, G.V.N. and Reddy, M.R.
Effect of ammonia treatment and processing of whole cotton plants as sole source of roughage in complete feeds for growing cross-bred calves 13(1985)93
Reddy, G.V.N. and Reddy, M.R.
Effects of ammonia treatment and processing of cotton straw as the sole source of roughage in complete feeds on rumen characteristics with Murrah buffaloes 14(1986)95
Reddy, M.R., see Reddy, G.V.N. 13(1985)93
Reddy, M.R., see Reddy, G.V.N. 14(1986)95
Redman, R.G., see Kellaway, R.C. 3(1978)201
Reid, G.W., see Ørskov, E.R. 8(1983)247
Reid, G.W., see Istasse, L. 15(1986)167
Reid, R.L. and Horvath, D.J.
Soil chemistry and mineral problems in farm livestock. A review 5(1980)95
Reid, S.J., see Hyslop, J.J. 25(1989)55
Rekib, A., see Singh, A.P. 9(1983)325
Rexen, B.
Enzyme solubility – a method for evaluating the digestibility of alkali-treated straw 2 (1977)205
Rexen, B., see Israelsen, M. 3(1978)227
Rexen, B.
Use of enzymes for improvement of feed 6 (1981)105
Rexen, F. and Vestergaard Thomsen, K.
The effect on digestibility of a new technique for alkali treatment of straw 1(1976)73
Rey, J.Ma., see Ocio, E. 4(1979)227
Rezende, S.A., see Brown, D.L. 18(1987)247
Richardson, C.R., see Bunting, L.D. 22(1988)61
Riggs, J.K., see Schake, L.M. 7(1982)257
Rigsby, L.L., see Akin, D.E. 19(1988)261
Rigsby, L.L., see Akin, D.E. 23(1989)305
Rioni, M., see Lanari, D. 1(1976)401
Rioni, M. and Lanari, D.
Bull-beef production from the maize crop in Italy in large and small units 1(1976)531
Riopérez, J., see Tortuero, F. 25(1989)79
Riou, Y., see Gouet, P. 2(1977)297
Ritchie, A.H. and Mackie, I.M.
Preparation of fish protein hydrolysates 7 (1982)125
Robert, P., Bertrand, D. and Demarquilly, C.
Prediction of forage digestibility by principal component analysis of near infrared reflectance spectra 16(1986/1987)215
Roberts, D.J.
The substitution of grass silage by a straw and concentrate mixture for dairy cows 20 (1988)135
Robertson, S., see Huhtanen, P. 21(1988)11
Robinson, J.J., see Kowalczyk, J. 7(1982)225
Robinson, P.H., Fadel, J.G. and Tamminga, S.
Evaluation of mathematical models to de-

scribe neutral detergent residue in terms of its susceptibility to degradation in the rumen 15(1986)249
Robinson, P.H., Brice, R.E. and Morrison, I.M.
Effect of straw treatment and laboratory of incubation on degradation of neutral detergent residue in the rumen 18(1987)75
Robinson, P.H., see Tamminga, S. 25(1989)89
Rodriguez, J., see Alibés, X. 8(1983)63
Rodriguez, J., see Alibes, X. 11(1984)189
Rodriguez, M.L., see Tortuero, F. 25(1989)79
Rollin, F., see Istasse, L. 25(1989)1
Romita, A.
Survey of the maize situation in Italy 1 (1976)109
Rönnau, K., see Von Engelhardt, W. 23(1989) 43
Roodbeen, A.E., see Papadopoulos, M.C. 14 (1986)279
Rooke, J.A., Brett, P.A., Overend, M.A. and Armstrong, D.G.
The energetic efficiency of rumen microbial protein synthesis in cattle given silage-based diets 13(1985)255
Rooke, J.A., Bell, S.L. and Armstrong, D.G.
The chemical composition of grass silages prepared with and without pre-treatment with inoculants containing *Lactobacillus plantarum* 13(1985)269
Rooke, J.A., see Lee, N.-H. 15(1986)301
Rooney, L.W., see Pflugfelder, R.L. 14(1986) 243
Rose, S.P. and Michie, W.
Meat and bone and fish meals in balancer feeds for choice-fed broilers 11(1984)221
Ross-Parker, H.M., see Jayasuriya, M.C.N. 7 (1982)201
Rossnagel, B.G., see Thacker, P.A. 19(1988) 191
Roth, F.X. and Kirchgessner, M.
Methanol-fermentation protein in veal calf nutrition 1(1976)33
Rotter, B.A., Neskar, M., Guenter, W. and Marquardt, R.R.
Effect of enzyme supplementation on the nutritive value of hulless barley in chicken diets 24(1989)233
Rubinstein, A., see Ben-Ghedalia, D. 15(1986) 47
Ruiz Jr., L.P., White, S.F. and Hove, E.L.
The alkaloid content of sweet lupin seed used in feeding trials on pigs and rats 2 (1977)59
Rule, D.C., Preston, R.L., Koes, R.M. and McReynolds, W.E.
Feeding value of sprouted wheat (*Triticum aestivum*) for beef cattle finishing diets 15 (1986)113
Rundgren, M.
Low-glucosinolate rapeseed products for pigs – a review 9(1983)239
Rundgren, M.
Evaluation of triticale given to pigs, poultry and rats 19(1988)359
Russell, J.R. and Buxton, D.R.
Storage of large round bales of hay harvested at different moisture concentrations and treated with sodium diacetate and/or covered with plastic 13(1985)69
Russell, J.R.
Influence of harvest date on the nutritive value and ensiling characteristics of maize stover 14(1986)11
Russell, J.R., see Adamu, A.M. 20(1988)241
Russell, J.R., see Adamu, A.M. 22(1988)227
Russo, V., Catalano, A., Mariani, P. and Del Monte, P.
Utilization of yeast grown on $_n$-paraffins in the feeding of heavy pigs, from weaning to slaughter 1(1976)25

S

Sadiq, M., see Slinger, S.J. 3(1978)251
Saiady, M.Y., see Fonnesbeck, P.V. 16(1986/1987)7
Said, A.N., see Nangole, F.N. 9(1983)121
Sakata, T., see Von Engelhardt, W. 23(1989) 43
Salcedo, T.S., see Gómez, G. 11(1984)57
Salman, A.J., see Razzaque, M.A. 14(1986) 265
Salman, A.J., see Ilian, M.A. 19(1988)327
Salman, A.J., see Ilian, M.A. 19(1988)343
Salmon, R.E.
Effects of pelleting, added sodium bentonite and fat in a wheat-based diet on performance and carcass characteristics of small white turkeys 12(1984/1985)223
Salvador, A., see González, G. 20(1988)177
Salyers, A.A. and McCarthy, R.E.
Assessing the importance of host-derived polysaccharides as carbon sources for bacteria growing in the human colon 23(1989) 109
Samuel, V., see Holzer, Z. 12(1984/1985)253
Samuel, V., see Levy, D. 16(1986/1987)63
Sanderson, M.A., Meyer, D.W. and Casper, H.H.
Dicoumarol concentrations in sweetclover hay treated with preservatives and in spoiled hay of high- and low-coumarin cultivars of sweetclover 14(1986)221
Santini, F.J., see Napoli, G.M. 25(1989)39
Santos, J., Gómez, G. and Alexander, J.C.
Production of fungal protein from rasped fresh cassava roots using 200- and 3000-litre fermentors 8(1983)313
Santos-Nunez, J., see Gregory, K.F. 2(1977)7
Sanz, R., see Fonolla, J. 6(1981)405
Sarwatt, S.V., Mussa, M.A. and Kategile, J.A.
The nutritive value of ensiled forages cut at three stages of growth 22(1988)237
Sauer, W.C. and Thacker, P.A.
Apparent ileal and faecal digestibility of

amino acids in barley-based diets supplemented with soya bean meal or canola meal for growing pigs 14(1986)183
Sauvant, D., Bertrand, D. and Giger, S.
Variations and prevision of the in sacco dry matter digestion of concentrates and by-products 13(1985)7
Sauveur, B. and Thapon, J.L.
Effects of dietary maize solubles and minerals on albumen quality of fresh and stored brown-shelled eggs 8(1983)177
Savić, R.
The use of techniques of maize cultivation to maximise feed for beef production 1(1976) 229
Savory, R., see Wood, J.F. 9(1983)307
Saxton, A.M., see Coelho, M. 20(1988)219
Schaefer, D.M., Brotz, P.G., Arp, S.C. and Cook, D.K.
Inoculation of corn silage and high-moisture corn with lactic acid bacteria and its effects on the subsequent fermentations and on feedlot performance of beef steers 25(1989) 23
Schake, L.M., Ellis, W.C., Suarez, W.A. and Riggs, J.K.
Preservation of sorghum plant portions harvested, processed and ensiled at ten stages of maturity 7(1982)257
Schake, L.M., see Pflugfelder, R.L. 14(1986) 243
Scheele, C.W., see Kan, C.A. 19(1988)97
Schemm, R.L., see Hibberd, C.A. 15(1986)231
Schenoni, P., see Giardini, A. 1(1976)313
Schneeberger, H.
Survey of the use of maize for livestock feeding in Switzerland 1(1976)149
Schöne, F., see Lüdke, H. 22(1988)33
Schöne, F., Lüdke, H., Hennig, A. and Jahreis, G.
Copper and iodine in pig diets with high glucosinolate rapeseed meal. II. Influence of iodine supplements for rations with rapeseed meal untreated or treated with copper ions on performance and thyroid hormone status of growing pigs 22(1988)45
Schultz, J.A. and Tolan, J.W.
Particle size distribution of a feed premix containing amprolium 3(1978)211
Schulz, E. and Oslage, H.J.
Composition and nutritive value of single-cell protein (SCP) 1(1976)9
Schwarz, F.J., see Kirchgessner, M. 6(1981) 337
Seghal, J.P. and Punj, M.L.
Utilization of alkali-treated and neutralized wheat straw-based rations for growing goat kids 9(1983)155
Sekine, J., Morita, Z., Oura, R. and Asahida, Y.
Water-drinking frequency and quantity of water drunk by steers fed hay or fresh forage 25(1989)207
Seligman, N.G., see Goldman, A. 18(1987)233
Setälä, J. and Syrjälä-Qvist, L.
Degradation of crude protein and quality of undegradable protein in untreated or formaldehyde-treated rapeseed meal 12 (1984/1985)19
Settineri, D., see Pace, V. 11(1984)125
Sharma, H.R., Ingalls, J.R. and McKirdy, J.A.
Feeding value of alkali-treated whole rye (*Secale cereale* L.) grain for lactating cows and its digestibility for sheep 10(1983/1984) 77
Sharma, H.R., White, B. and Ingalls, J.R.
Utilization of whole rape (Canola) seed and sunflower seeds as sources of energy and protein in calf starter diets 15(1986)101
Sharma, H.R., see Mbugi, P.K. 24(1989)267
Sharma, O.P., see Lall, D. 9(1983)29
Sharp, R.N., see Noland, P.R. 2(1977)219
Sharp, R.N., Noland, P.R. and Campbell, D.R.
Relationships between tristimulus colour values and digestibility of grain sorghum by swine 2(1977)327
Shelby, T., see Slabbert, N. 21(1988)31
Shen, H., Summers, J.D. and Leeson, S.
The influence of steam pelleting and grinding on the nutritive value of Canola rapeseed for poultry 8(1983)303
Shen, H., Summers, J.D. and Leeson, S.
Influence of supplemental fat and steam pelleting on the response of broilers to diets varying in energy density 13(1985)57
Shimada, A., see Alcántara, E. 23(1989)323
Shimojo, M. and Goto, I.
Effect of sodium silicate on forage digestion with rumen fluid of goats or cellulase using culture solutions adjusted for pH 24(1989) 173
Shin, H.T., see Greenhalgh, J.F.D. 3(1978)289
Shqueir, A.A., Brown, D.L. and Klasing, K.C.
Canavanine content and toxicity of sesbania leaf meal for growing chicks 25(1989)137
Sibbald, I.R., Elliot, J.I. and Cote, M.
Comparison of bioavailable energy values of diets measured with poultry and swine 8 (1983)191
Sidahmed, A.E., Morris, J.G., Radosevich, S.R. and Koong, L.J.
Seasonal changes in composition and intake of chaparral by Spanish goats 8(1983)47
Silanikove, N., Cohen, O., Levanon, D., Kipnis, T. and Kugenheim, Y.
Preservation and storage of green panic (*Panicum maximum*) as moist hay with urea 20(1988)87
Silva, A.T. and Ørskov, E.R.
Fibre degradation in the rumens of animals receiving hay, untreated or ammonia-treated straw 19(1988)277
Silva, A.T. and Ørskov, E.R.
The effect of five different supplements on the degradation of straw in sheep given untreated barley straw 19(1988)289
Silverside, D.E., see Machin, D.H. 15(1986)273

Singh, A.P. and Pandit, N.N.
Studies on fermentation of sorghum silage during storage - effect of urea and molasses 3(1978)299
Singh, A.P. and Pandita, N.N.
Studies on fermentation of sorghum silage during storage, and its effect on milch animals 9(1983)143
Singh, A.P., Prasad, J. and Rekib, A.
Note on high-nitrate para grass (*Brachiaria mutica*) silage given to rabbits 9(1983)325
Singh, B., see Negi, S.S. 22(1988)79
Singh, K., see Jakhmola, R.C. 19(1988)1
Singh, K.S., see Deolankar, R.P. 4(1979)133
Singh, K.S. and Prasad, C.M.
Feeding value of sunflower and groundnut cakes for broilers 4(1979)143
Singh, K.S., Prasad, C.M. and Brahmakshatriva, R.D.
Feeding value of sunflower- and groundnut-cakes for laying hens 6(1981)63
Singh, R., see Kamra, D.N. 9(1983)185
Singh, R., Kamra, D.N. and Jakhmola, R.C.
Ensiling of leguminous green forages in combination with different dry roughages and molasses 12(1984/1985)133
Singh, R., see Jakhmola, R.C. 19(1988)1
Siroka, P., see Baran, M. 15(1986)7
Siviter, J.W., see Smith, T. 10(1983/1984)189
Siviter, J.W., see Smith, T. 14(1986)29
Skoch, E.R., Behnke, K.C., Deyoe, C.W. and Binder, S.F.
The effect of steam-conditioning rate on the pelleting process 6(1981)83
Skraba, B., see Hanczakowski, P. 6(1981)189
Skraba, B., see Hanczakowski, P. 6(1981)413
Skraba, B., see Hanczakowski, P. 12(1984/1985)11
Skraba, B., see Hanczakowski, P. 24(1989)151
Skrede, A. and Nes, I.F.
Slaughterhouse by-products preserved by *Lactobacillus plantarum* fermentation as feed for mink and foxes 20(1988)287
Slabbert, N., Bolt, M.J., Shelby, T. and Campher, J.P.
The utilization by beef steers of a soft high lysine and a hard normal size maize cultivar in whole grain feedlot diets 21(1988)31
Slinger, S.J., Leeson, S., Summers, J.D. and Sadiq, M.
Influence of steam pelleting on the feeding value of tower and candle rapeseed products for broiler chickens 3(1978)251
Smart, M.E., see Mann, M.E. 21(1988)57
Smart, R.I., see Mbatya, P.B.A. 8(1983)221
Smart, R.I., see Mbatya, P.B.A. 8(1983)229
Smart, R.I., see Mbatya, P.B.A. 13(1985)281
Smart, R.I., see Mbatya, P.B.A. 13(1985)293
Smart, R.I., see Istasse, L. 16(1986/1987)37
Smith, M.A., Watson Jr., C.E., Watson, V.H. and Trevathan, L.E.
Effects of *Drechslera sorokiniana* infection on the yield and quality of tall fescue forage 15(1986)41
Smith, M.J., see Daniels, L.B. 8(1983)19
Smith, O.B. and Macleod, G.K.
Digestibility of wet cage layer excreta and an excreta-supplemented diet by cattle as determined by two methods 5(1980)77
Smith, O.B., Ilori, J.O. and Onesirosan, P.
The proximate composition and nutritive value of the winged bean *Psophocarpus tetragonolobus* (L.) DC for broilers 11 (1984)231
Smith, O.B. and Adegbola, A.A.
Studies on the feeding value of agro-industrial by-products. III. Digestibility of cocoa-pod and cocoa-pod-based diets by ruminants 13(1985)249
Smith, O.B., see Kabaija, E. 20(1988)171
Smith, O.B., Osafo, E.L.K. and Adegbola, A.A.
Studies on the feeding value of agro-industrial by-products: strategies for improving the utilisation of cocoa-pod-based diets by ruminants 20(1988)189
Smith, R.E., see Gregory, K.F. 2(1977)7
Smith, T., Grigera-Naón, J.J., Broster, W.H. and Siviter, J.W.
Ammonia versus sodium hydroxide treatment of straw for growing cattle 10(1983/1984)189
Smith, T., Grigera-Naón, J.J., Broster, W.H. and Siviter, J.W.
The use of barley straw as a feed 14(1986) 29
Smith, W.C., see Kies, A.K. 16(1986/1987)169
Smith, W.C., Moughan, P.J., Pearson, G. and James, K.A.C.
Comparative bioavailable energy values of five ground cereal grains measured with growing rats and pigs 18(1987)143
Smith, W.C., Moughan, P.J., Pearson, G. and Patchell, M.R.
A comparison of bioavailable energy values of ground cereal grains measured with adult cockerels and growing pigs 19(1988)105
Smith, W.C., see Moughan, P.J. 22(1988)203
Smits, C.H.M., see Moughan, P.J. 22(1988) 203
Sobale, B.N., see Rangnekar, D.V. 7(1982)61
Søbstad, G., see Opstvedt, J. 18(1987)181
Soliman, H., see Mohamed, K. 11(1984)199
Soliman, H.S., see Ørskov, E.R. 7(1982)135
Solomon, R., see Ben-Ghedalia, D. 18(1987)55
Solomon, R., see Ben-Ghedalia, D. 22(1988) 147
Sonaiya, E.B. and Omole, T.A.
Cassava meal and cassava peel meal in diets for growing pigs 8(1983)211
Sosulski, F.W., see Campbell, G.L. 16(1986/1987)243
Spears, J.W., see Guardiola, C.M. 8(1983)129
Spears, J.W., see Harvey, R.W. 24(1989)337
Spoelstra, S.F., see Courtin, M.G. 24(1989)97
Spooner, A.E., see Higgins, T.R. 16(1986/1987)1
Spooner, M.C., see Osbourn, D.F. 6(1981)387

Spragg, J.C., see Amaning-Kwarteng, K. 16 (1986/1987)75
Spurr, D.T., see Kernan, J.A. 6(1981)257
Spurr, D.T., see Kernan, J.A. 11(1984)301
Srećković, A.
Review of production and utilisation of maize for animal nutrition in Yugoslavia 1 (1976)115
Srivastava, R.V.N., see Kamra, D.N. 9(1983) 185
Stallcup, O.T., see Daniels, L.B. 8(1983)19
Stanogias, G. and Pearce, G.R.
Digestibility by cattle of diets containing dried pig faeces 3(1978)155
Staples, C.R. and Lough, D.S.
Efficacy of supplemental dietary neutralizing agents for lactating dairy cows. A review 23(1989)277
Stappers, H.P., see Grimbergen, A.H.M. 13 (1985)117
Steg, A., see Deinum, B. 10(1983/1984)301
Steg, A. and van der Meer, J.M.
Differences in chemical composition and digestibility of beet and cane molasses 13 (1985)83
Stehr, W. and Kirchgessner, M.
The relationship between the intake of herbage grazed by dairy cows and its digestibility 1(1976)53
Stephen, A.M.
The physiological effects of cellulose in the human large intestine 23(1989)241
Stern, M.D., see Hannah, S.M. 16(1986/1987) 51
Stern, M.D., see Waltz, D.M. 25(1989)111
Stewart, C.S., see Greenhalgh, J.F.D. 3(1978) 289
Stewart, C.S., see Theander, O. 10(1983/ 1984)89
Stewart, C.S., see Cheng, K.-J. 10(1983/1984) 93
Stewart, C.S., see Graham, H. 12(1984/1985) 195
Stewart, C.S., Fonty, G. and Gouet, Ph.
The establishment of rumen microbial communities 21(1988)69
Stobbs, the late T.H., see Aii, T. 5(1980)183
Streeter, C.L. and Horn, G.W.
Effect of treatment of wheat straw with ammonia and peracetic acid on digestibility in vitro and cell wall composition 7(1982) 325
Streeter, C.L., see Horn, G.W. 8(1983)35
Suarez, W.A., see Schake, L.M. 7(1982)257
Subramanian, P.R., Kadirvel, R., Viswanathan, K. and Chandrasekaran, D.
In vitro studies and short-term feeding trial in lambs to evaluate plantain sheath (*Musa sapientum*) as a feed for ruminants 20(1988) 343
Sudana, I.B. and Leng, R.A.
Effects of supplementing a wheat straw diet with urea or a urea–molasses block and/or cottonseed meal on intake and liveweight change of lambs 16(1986/1987)25
Sulieman, A.H., see Osman, A.G. 19(1988)351
Summers, J.D. and Leeson, S.
Utilization of processed pig hair in poultry diets 3(1978)243
Summers, J.D., see Slinger, S.J. 3(1978)251
Summers, J.D., Macleod, G.K. and Warner, W.C.
Chemical composition of culinary wastes and their potential as a feed for ruminants 5(1980)205
Summers, J.D., see Shen, H. 8(1983)303
Summers, J.D., see Leeson, S. 11(1984)211
Summers, J.D., see Shen, H. 13(1985)57
Summers, J.D. and Leeson, S.
Influence of nutrient density on feed consumption, weight gain and gut capacity of broilers, Leghorns and turkeys reared to 26 days of age 16(1986/1987)129
Summers, J.D., see Leeson, S. 19(1988)313
Sundstøl, F., see Kategile, J.A. 6(1981)133
Sundstøl, F., see Borhami, B.E.A. 7(1982)45
Sundstøl, F., see Borhami, B.E.A. 7(1982)53
Sundstøl, F.
Ammonia treatment of straw: methods for treatment and feeding experience in Norway 10(1983/1984)173
Sundstøl, F., see Wanapat, M. 12(1984/1985) 295
Sundstøl, F., see Dias-da-Silva, A.A. 14(1986) 67
Sundstøl, F., see Wanapat, M. 14(1986)215
Suttle, N.F.
Reducing the potential copper toxity of concentrates to sheep by the use of molybdenum and sulphur supplements 2(1977)235
Swan, H., see Karalazos, A. 2(1977)143
Swatson, H., see Atuahene, C.C. 17(1987)157
Syrjala, L., see Pulli, S. 1(1976)187
Syrjälä-Qvist, L., see Setälä, J. 12(1984/1985) 19
Szylit, O., Durand, M., Borgida, L.P., Atinkpahoun, H., Prieto, F. and Delort-Laval, J.
Raw and steam-pelleted cassava, sweet potato and yam *cayenensis* as starch sources for ruminant and chicken diets 3(1978)73

T

Tait, C.A.G., see Ørskov, E.R. 8(1983)247
Tait, C.A.G., see Istasse, L. 15(1986)167
Tait, R.M., see Pringle, D. 9(1983)89
Tait, R.M., see Beames, R.M. 15(1986)285
Tamminga, S., see Robinson, P.H. 15(1986) 249
Tamminga, S. and van Vuuren, A.M.
Formation and utilization of end products of ligno cellulose degradation in ruminants 21(1988)141
Tamminga, S., see Ibraham, M.N.M. 24(1989) 83
Tamminga, S., Robinson, P.H., Vogt, M. and

Boer, H.
Rumen ingesta kinetics of cell wall components in dairy cows 25(1989)89
Tangprasertchai, P., see McEllhiney, R.R. 8 (1983)139
Tangtaweewipat, S. and Elliott, R.
Nutritional value of pigeonpea (*Cajanus cajan*) meal in poultry diets 25(1989)123
Tatterson, I.N.
Fish silage – preparation, properties and uses 7(1982)153
Taylor, A. G., see Morgan, C.A. 24(1989)219
Taylor, A.G., see Peers, D.G. 2(1977)41
Teh, T.H., Hemken, R.W., Bremel, D.H. and Harmon, R.J.
Comparison of buffers on rumen functions, turnover rate and gastric secretions in Holstein steers 17(1987)257
Telfer, S.B., see Zervas, G. 21(1988)23
Teller, E. and Godeau, J.-M.
The effect of nitrogen supplements to maize silage on the amount and composition of the nitrogen fraction entering the intestine of cattle 13(1985)25
Terry, R.A., see Osbourn, D.F. 6(1981)387
Tetlow, R.M., see Osbourn, D.F. 6(1981)387
Tetlow, R.M., see Bolsen, K.K. 9(1983)37
Tetlow, R.M.
The effect of urea on the preservation and digestibility in vitro of perennial ryegrass 10 (1983/1984)49
Tetlow, R.M., see Woolford, M.K. 11(1984)159
Tetlow, R.M., Wilson, R.F. and Bolsen, K.K.
Effect of sodium acrylate and calcium and sodium hydroxides on the fermentation, digestibility, chemical composition and aerobic deterioration of whole-crop maize silage 12(1984/1985)91
Tetlow, R.M. and Mason, V.C.
Treatment of whole-crop cereals with alkali. 1. The influence of sodium hydroxide and ensiling on the chemical composition and in vitro digestibility of rye, barley and wheat crops harvested at increasing maturity and dry matter content 18(1987)257
Tetlow, R.M., Mason, V.C. and Deschard, G.
Treatment of whole-crop cereals with alkali. 2. Voluntary intake and digestibility by sheep of rye, barley and wheat crops ensiled with sodium hydroxide 18(1987)271
Tetlow, R.M., see Deschard, G. 18(1987)283
Tetlow, R.M., see Deschard, G. 19(1988)55
Tewe, O.O.
Effect of cassava-based diets varying in cyanide content on the performance and physiopathology of the African giant rat (*Cricetomys gambianus* Waterhouse) 11 (1984)1
Thacker, P.A., see Bowland, J.P. 12(1984/1985)75
Thacker, P.A., see Sauer, W.C. 14(1986)183
Thacker, P.A., Bell, J.M., Classen, H.L., Campbell, G.L. and Rossnagel, B.G.
The nutritive value of hulless barley for swine 19(1988)191
Tham, S.H., see Karunajeewa, H. 17(1987)271
Thanu, K., Kadirvel, R. and Ayyaluswami, P.
The effect of nutrient supplementation on the feeding value of kapok seed for poultry 9(1983)263
Thapon, J.L., see Sauveur, B. 8(1983)177
Theander, O., Hartley, R.D. and Stewart, C.S.
A note on the OECD collaborative experiment on ammonia-treated straw 10(1983/1984)89
Theander, O., see Graham, H. 12(1984/1985) 195
Theander, O., see Lindberg, J.E. 15(1986)197
Theander, O.
Possibilities for multipurpose fractionation and alternative uses of plant biomass 21 (1988)267
Theander, O., Westerlund, E., Åman, P. and Graham, H.
Plant cell walls and monogastric diets 23 (1989)205
Theodorou, M.K., see Akin, D.E. 19(1988)261
Theodorou, M.K., see Gascoyne, D.J. 21 (1988)183
Thiago, L.R.L., see Kellaway, R.C. 3(1978)201
Thiago, L.R.L. de S. and Kellaway, R.C.
Botanical composition and extent of lignification affecting digestibility of wheat and oat straw and paspalum hay 7(1982)71
Thiesen, J.
Detoxification of aflatoxins in groundnut meal 2(1977)67
Thivend, P., see Jouany, J.P. 15(1986)215
Thomas, H.R., see Harper, A.F. 8(1983)69
Thomas, V.M., Katz, R.J., Auld, D.L. and Peterson, C.L.
Value of mechanically extracted rape and safflower oilseed meals as protein supplements for growing lambs 11(1984)269
Thomke, S., see Muindi, P.J. 6(1981)197
Thomke, S., see Hesselman, K. 7(1982)351
Thompson, C. and Moran, J.B.
The rate of dry matter disappearance from nylon bags of maize silage/grain diets as influenced by basal ration and processing of test samples 16(1986/1987)225
Thompson, L.U., see McBurney, M.I. 23(1989) 261
Thomson, E., see Farrell, D.J. 9(1983)99
Timbers, G.E., see Elliot, J.I. 17(1987)213
Tjerneld, F., see Hahn-Hägerdal, B. 21(1988) 175
Tock, R.W., see Bunting, L.D. 22(1988)61
Tolan, J.W., see Schultz, J.A. 3(1978)211
Topps, J.H., see El-Sherbiny, A.E. 5(1980)33
Topps, J.H., see El-Sherbiny, A.E. 5(1980)43
Tortuero, F., Riopérez, J. and Rodriguez, M.L.
Nutritional value for rabbits of olive pulp and the effects on their visceral organs 25 (1989)79
Touchburn, S.P., see Chavez, E.R. 22(1988)1
Touchburn, S.P., see Chavez, E.R. 22(1988)13

Touchburn, S.P., see Chavez, E.R. 22(1988)23
Treacher, T.T., see Penning, P.D. 5(1980)321
Trenholm, H.L., see Hamilton, R.M.G. 11 (1984)293
Trenkle, A., see Adamu, A.M. 20(1988)241
Trenkle, A., see Adamu, A.M. 22(1988)227
Trevathan, L.E., see Smith, M.A. 15(1986)41
Treviño, J., Centeno, C. and Caballero, R.
The chemical composition of pea plant parts as related to harvesting time 16(1986/1987) 305
Trigg, T.E., see Moran, J.B. 20(1988)299
Tuite, J., see Cherney, J.H. 17(1987)45
Tullis, J.B. and Whittemore, C.T.
Digestibility of a fully hydrogenated tallow for growing pigs 5(1980)87
Twumasi, I.K., see Osei, S.A. 24(1989)247

U

Udedibie, A.B.I. and Igwe, F.O.
Dry matter yield and chemical composition of pigeon pea (*C. cajan*) leaf meal and the nutritive value of pigeon pea leaf meal and grain meal for laying hens 24(1989)111
Uden, P. and Van Soest, P.J.
The determination of digesta partical size in some herbivores 7(1982)35
Udén, P.
The effect of intake and hay: concentrate ratio upon digestibility and digesta passage 11(1984)167
Udén, P.
Digestibility and digesta retention in dairy cows receiving hay or silage at varying concentrate levels 11(1984)279
Udén, P.
The effect of grinding and pelleting hay on digestibility, fermentation rate, digesta passage and rumen and faecal particle size in cows 19(1988)145
Umunna, N.N., Klopfenstein, T.J., Hasimoglu, S. and Woods, W.R.
Evaluation of corn gluten meal with urea as a source of supplementary nitrogen for growing calves and lambs 7(1982)375
Umunna, N.N., Bartling, R.R. and Klopfenstein, T.J.
High temperature and pressure processing of maize cobs. I. Digestibility in vitro of processed cobs 12(1984/1985)151
Umunna, N.N., Bartling, R.R. and Klopfenstein, T.J.
High temperature and pressure processing of maize cobs. II. Animal response and effect of sodium metabisulphite 14(1986)161
Urio, N.A., see Kategile, J.A. 6(1981)133

V

Vadiveloo, J.
The intake and digestibility in goats of *Leucaena leucocephala* supplemented with dehydrated palm oil mill effluent 24(1989) 45
Vaid, J., see Lall, D. 9(1983)29
Valdivie, M., Compte, X. and Fundora, O.
The utilization of torula yeast in diets for White Leghorn birds during growth and laying periods 7(1982)185
Valdivieso, M., see Gómez, G. 11(1984)49
Valdivieso, M., see Gómez, G. 11(1984)57
Valentine, S.C. and Bartsch, B.D.
Digestibility of dry matter, nitrogen and energy by dairy cows fed whole or hammermilled lupin grain in oaten hay or oaten pasture based diets 16(1986/1987)143
Valentine, S.C. and Bartsch, B.D.
Fermentation of hammermilled barley, lupin, pea and faba bean grain in the rumen of dairy cows 16(1986/1987)261
van der Aar, P.J., see Mora, M.I. 9(1983)205
van der Honing, Y., see Van Es, A.J.S. 1(1976) 481
van der Meer, J.M., see Steg, A. 13(1985)83
Van Eenaeme, C., see Istasse, L. 25(1989)1
van Es, A.J.S., van der Honing, Y. and Boekholt, H.A.
The nutritive value (energy) of maize silage 1(1976)481
Van Eys, J.E., Pulungan, H., Rangkuti, M. and Johnson, W.L.
Cassava meal as supplement to napier grass diets for growing sheep and goats 18(1987) 197
van Houtert, M.F.J., see Jayasuriya, M.C.N. 7(1982)201
Van Nevel, C.J., see Kayouli, C. 10(1983/ 1984)165
Van Soest, P.J., see Uden, P. 7(1982)35
Van Soest, P.J., Mascarenhas Ferreira, A. and Hartley, R.D.
Chemical properties of fibre in relation to nutritive quality of ammonia-treated forages 10(1983/1984)155
van Soest, P.J., see Lau, M.M. 6(1981)123
van Vuuren, A.M., see Tamminga, S. 21(1988) 141
Van Weerden, E.J. and Huisman, J.
Digestibility of protein and amino acids of a fermentation single-cell protein for veal calves 2(1977)377
Vanacker, J.M., see Cottyn, B.G. 11(1984)137
Vanacker, J.M., see De Boever, J.L. 14(1986) 203
Vanacker, J.M., see De Boever, J.L. 19(1988) 247
Vanbelle, M., see Nefzaoui, A. 14(1986)139
Vandevoorde, L., Bruyneel, B. and Verstraete, W.
In-reactor cellulose conversion 21(1988)99
Vecchiettini, M., see Giardini, A. 1(1976)369
Vechiettini, M., see Giardini, A. 1(1976)313
Vedder, H.M., see Barton II, F.E. 15(1986)189
Veira, D.M., see Ivan, M. 9(1983)131
Vencl, B. and Flám, F.

Predicting the energetic value of forage from digestibility of organic matter in vitro or solubility in potassium hydroxide 6(1981) 235
Verstraete, W., see Vandevoorde, L. 21(1988) 99
Vervaeke, I.J., see Dierick, N.A. 23(1989)141
Vervaeke, I.J., Dierick, N.A., Demeyer, D.I. and Decuypere, J.A.
Approach to the energetic importance of fibre digestion in pigs. II. An experimental approach to hindgut digestion 23(1989)169
Vestergaard Thomsen, K., see Rexen, F. 1 (1976)73
Vestergaard Thomsen, K., see Waagepetersen, J. 2(1977)131
Vestergaard Thomsen, K., see El Khidir, O.A. 7(1982)277
Vestergaard Thomsen, K., see El Khidir, O.A. 9(1983)197
Vignon, B., see Laurent, F. 13(1985)131
Viñaras, R., see Ocio, E. 4(1979)227
Viswanathan, K., see Subramanian, P.R. 20 (1988)343
Viswanathan, K., see Chandrasekaran, D. 22 (1988)321
Viswanathan, K., Kadirvel, R. and Chandrasekaran, D.
Nutritive value of banana stalk (*Musa cavendishi*) as a feed for sheep 22(1988)327
Vlach, D., see Moo-Young, M. 5(1980)175
Vogt, M., see Tamminga, S. 25(1989)89
Volenec, J.J., see Cherney, J.H. 17(1987)45
Von Engelhardt, W., Rönnau, K., Rechkemmer, G. and Sakata, T.
Absorption of short-chain fatty acids and their role in the hindgut of monogastric animals 23(1989)43

W

Waagepetersen, J. and Vestergaard Thomsen, K.
Effect on digestibility and nitrogen content of barley straw of different ammonia treatments 2(1977)131
Wagner, D.G., see Hibberd, C.A. 15(1986)231
Wainman, F.W., see De Boever, J.L. 14(1986) 203
Waldroup, P.W., Bussell, W.D., Burke, A.B. and Johnson, Z.B.
Comparison of the relative biological availability and stability of commercial sources of vitamin K activity 3(1978)329
Waldroup, P.W., Whelchel, D.L. and Johnson, Z.B.
Variation in nutrient content of samples of dried bakery product 7(1982)419
Walker, H.G., Mackay, B.E. and Montague Jr., W.C.
Composition and enzymatic digestibility of oregon grass straws 9(1983)283
Walker, N.
The composition of processed abattoir blood and its use in finishing-pig diets 2(1977)21
Walker, N.
Treatment of the haem fraction of abattoir blood with formic acid to produce a solid state, low dry matter protein supplement for use in finishing-pig diets 2(1977)367
Walker, N., see Patterson, D.C. 4(1979)263
Walker, N., see Patterson, D.C. 4(1979)275
Walker, N.
A comparison of wheat- or barley-based diets given ad libitum as meal or pellets to finishing pigs 22(1988)263
Walker, N.
Observations on the feeding of non-lactating sows on graded levels of liquid whey 24 (1989)327
Walli, T.K., see Bhargava, P.K. 22(1988)295
Walter, K.
Economic aspects of high-plane feeding of bulls with maize silage 1(1976)513
Waltz, D.M. and Stern, M.D.
Evaluation of various methods for protecting soya bean protein from degradation by rumen bacteria 25(1989)111
Wanapat, M., Sundstøl, F. and Garmo, T.H.
A comparison of alkali treatment methods to improve the nutritive value of straw. I. Digestibility and metabolizability 12(1984/ 1985)295
Wanapat, M., Sundstøl, F. and Hall, J.M.R.
A comparison of alkali treatment methods used to improve the nutritive value of straw. II. In sacco and in vitro degradation relative to in vivo digestibility 14(1986)215
Ward, J.K., see Faulkner, D.B. 12(1984/1985) 125
Warner, W.C., see Summers, J.D. 5(1980)205
Waterworth, D.G. and Heath, M.E.
Pruteen in the diet of breeding pigs: reproductive performance 6(1981)297
Watson, C., see Batterham, E.S. 13(1985)171
Watson, V.H., see Smith, M.A. 15(1986)41
Watson Jr., C.E., see Smith, M.A. 15(1986)41
Weibel, D.E., see Hibberd, C.A. 15(1986)231
Weigand, E. and Kirchgessner, M.
Protein and energy value of vinasse for pigs 5(1980)221
Weinberg, Z.G., Pahlow, G., Dinter, B. and Ashbell, G.
The effect of treatment with urea, sorbic acid, or dehydration on orange peel silage 20(1988)335
Weir, J., see Hyslop, J.J. 25(1989)55
Weller, R.F., see Phipps, R.H. 1(1976)251
Werner, D., see Arieli, A. 23(1989)333
Westerlund, E., see Theander, O. 23(1989)205
Westlake, K., Mackie, R.I. and Dutton, M.F.
In vitro metabolism of mycotoxins by bacterial, protozoal and ovine ruminal fluid preparations 25(1989)169
Whelchel, D.L., see Waldroup, P.W. 7(1982) 419
White, B., see Sharma, H.R. 15(1986)101

White, S.F., see Ruiz Jr., L.P. 2(1977)59
Whitehead, D.C., Goulden, K.M. and Hartley, R.D.
Fractions of nitrogen, sulphur, phosphorus, calcium and magnesium in the herbage of perennial ryegrass as influenced by fertilizer nitrogen 14(1986)231
Whittemore, C.T., see Peers, D.G. 2(1977)41
Whittemore, C.T.
The potato (*Solanum tuberosum*) as a source of nutrients for pigs, calves and fowl – a review 2(1977)171
Whittemore, C.T., see Tullis, J.B. 5(1980)87
Whittemore, C.T., see Morgan, C.A. 7(1982) 387
Whittemore, C.T., see Morgan, C.A. 11(1984) 11
Whittemore, C.T., see Morgan, C.A. 17(1987) 81
Whittemore, C.T., see Morgan, C.A. 19(1988) 185
Whittemore, C.T., see Morgan, C.A. 24(1989) 219
Wignjosoesastro, N. and Young, A.W.
Digestibility of diets containing increasing levels of NaOH-treated or untreated wheat straw 7(1982)331
Wijeyatunge, C., see Jayasuriya, M.C.N. 7 (1982)221
Wilcock, J.M., see Hyslop, J.J. 25(1989)55
Wilkins, R.J.
Discussion on session 2 1(1976)273
Wilkinson, J.M.
Discussion on session I 1(1976)215
Wilkinson, J.M.
Voluntary intake and efficiency of utilisation of whole-crop maize silage 1(1976)441
Wilkinson, J.M. and Gonzalez Santillana, R.
Ensiled alkali-treated straw. I. Effect of level and type of alkali on the composition and digestibility in vitro of ensiled barley straw 3(1978)117
Wilkinson, J.M. and Gonzalez Santillana, R.
Ensiled alkali-treated straw. II. The nutritive value for young beef cattle of mixtures of ensiled or frozen alkali-treated straw and ryegrass silage 3(1978)133
Wilkinson, J.M.
Laboratory evaluation of ensiled mixtures of alkali-treated cattle excreta and whole-crop maize 3(1978)335
Wilkinson, J.M., see Gonzalez Santillana, R. 3(1978)381
Wilkinson, J.M., see Woolford, M.K. 8(1983) 107
Williams, A.G.
Factors affecting the formation of polysaccharide-degrading enzymes by rumen micro-organisms 21(1988)191
Williams, J.E., see Goedeken, F.K. 13(1985) 47
Williams, K.C., see Neill, A.R. 9(1983)71
Williams, P.E.V., see Ørskov, E.R. 6(1981)355
Williams, P.E.V.
Digestibility studies on ammonia-treated straw 10(1983/1984)213
Williams, P.E.V., Macdearmid, A., Innes, G.M. and Gauld, S.A.
Ammonia-treated barley straw and rolled barley offered either together, in a mixed ration, or successively to beef steers 10(1983/1984)247
Williams, P.E.V., Innes, G.M. and Brewer, A.
Ammonia treatment of straw via the hydrolysis of urea. I. Effects of dry matter and urea concentrations on the rate of hydrolysis of urea 11(1984)103
Williams, P.E.V., Innes, G.M. and Brewer, A.
Ammonia treatment of straw via the hydrolysis of urea. II. Additions of soya bean (urease), sodium hydroxide and molasses; effects on the digestibility of urea-treated straw 11(1984)115
Williams, P.E.V., Pusztai, A.J., Macdearmid, A. and Innes, G.M.
The use of kidney beans (*Phaseolus vulgaris*) as protein supplements in diets for young rapidly growing beef steers 12(1984/1985)1
Williams, P.E.V., see Fallon, R.J. 14(1986)103
Wilson, J.R., see McLeod, M.N. 24(1989)185
Wilson, M.N., see Moughan, P.J. 22(1988)203
Wilson, P.N., Brigstocke, T.D.A. and Cuthbert, N.H.
Some factors affecting the future composition of UK compound animal feeds 6(1981) 1
Wilson, R.F., see Bolsen, K.K. 9(1983)37
Wilson, R.F., see Tetlow, R.M. 12(1984/1985) 91
Wiseman, J.
A note on the nutritive value of dried instant coffee residue for broiler chickens and turkey poults 10(1983/1984)285
Wiseman, J., see Green, S. 21(1988)43
Wohlt, J.E., Fiallo, J.F. and Miller, M.E.
Composition of by-products of the essential-oil industry and their potential as feeds for ruminants 6(1981)115
Wolf, I., see Flachowsky, G. 17(1987)235
Wolfram, D., see Flachowsky, G. 17(1987)235
Wolsink, J.H., see Barton II, F.E. 15(1986)189
Wood, J.F., Carter, P.M. and Savory, R.
Investigations into the effects of processing on the retention of the carotenoid fractions of *Leucaena leucocephala* during storage, and the effects of processing on mimosine concentration 9(1983)307
Wood, J.F.
The functional properties of feed raw materials and their effect on the production and quality of feed pellets 18(1987)1
Woods, W.R., see Umunna, N.N. 7(1982)375
Woolford, M.K. and Cook, J.E.
A note on the effect on the aerobic deterioration of maize silage of the manipulation of the microflora by means of antibiotics 3 (1978)89

Woolford, M.K., Wilkinson, J.M. and Cook, J.E.
Investigations on the effect of sodium bentonite and other moisture absorbents on the production of effluent from grass silage 8 (1983)107
Woolford, M.K. and Tetlow, R.M.
The effect of anhydrous ammonia and moisture content on the preservation and chemical composition of perennial ryegrass hay 11(1984)159
Wrathall, J.H.M., Owen, E. and Pike, D.J.
Upgrading barley straw for goats: the effectiveness of a sodium hydroxide and urea dip method 24(1989)57
Wright, D.E., Payne, E., Pyle, C., Aitken, W.M. and Kirton, A.H.
Poly-unsaturated fat in young ruminants 2 (1977)93
Wyers, M., see Cherbut, C. 20(1988)327
Wyss, U. and Bickel, H.
Ripe beans of *Canavalia ensiformis* (jackbean) as feed ingredient for monogastric animals 20(1988)325

Y

Yosef, E., see Ben-Ghedalia, D. 19(1988)219
Yosef, E., see Ben-Ghedalia, D. 22(1988)247
Yosef, E., see Ben-Ghedalia, D. 22(1988)255
Yosef, E., see Ben-Ghedalia, D. 24(1989)289
Young, A.W., see Wignjosoesastro, N. 7(1982) 331
Yulzari, A., see Goldman, A. 18(1987)233

Z

Zacchi, G., see Hahn-Hägerdal, B. 21(1988) 175
Zarkadas, C.G., Hulan, H.W. and Proudfoot, F.G.
The amino acid and mineral composition of white fish meal containing enzyme-digested or untreated stickwater solids 14(1986)291
Zarkadas, C.G., see Hulan, H.W. 16(1986/1987)253
Zemmelink, G., see Ibraham, M.N.M. 24 (1989)83
Zervas, G.
Treatment of dairy sheep with soluble glass boluses containing copper, cobalt and selenium 19(1988)79
Zervas, G., Telfer, S.B., Carlos, G. and Anderson, P.
The effect of soluble-glass boluses containing copper, cobalt and selenium on the blood composition of ewes 21(1988)23
Zimmer, E. and Zscheischler, J.
Use of maize for livestock feeding in the Federal Republic of Germany 1(1976)157
Zimmer, E.
Efficiency of harvesting and conservation 1 (1976)289
Zimmer, E.
Discussion on session 3 1(1976)327
Ziras, E., see Papadopoulos, G. 18(1987)295
Ziras, E., see Papadopoulos, G. 18(1987)303
Zorilla-Rios, J., see Horn, G.W. 24(1989)201
Zorrilla-Rios, J., Horn, G.W. and McNew, R.W.
Effect of ammoniation and energy supplementation on the utilization of wheat straw by sheep 22(1988)305
Zscheischler, J., see Zimmer, E. 1(1976)157

SUBJECT INDEX

Animal Feed Science and Technology, volumes 1–25

A

Acid treatment
Studies on ammonia-treated straw. II. Fixation of ammonia in treated straw by spraying with acids 7(1982)53

Accuracy of predicting digestibility by the cellulase technique; the effect of pretreatment of forage samples with neutral detergent or acid pepsin 7(1982)83

Aflatoxins
Detoxification of aflatoxins in groundnut meal 2(1977)67

In vitro metabolism of mycotoxins by bacterial, protozoal and ovine ruminal fluid preparations 25(1989)169

African giant rat
Effect of cassava-based diets varying in cyanide content on the performance and physiopathology of the African giant rat (*Cricetomys gambianus* Waterhouse) 11 (1984)1

African yam
Variation in metabolizable energy content of raw or autoclaved white and brown varieties of three tropical grain legumes 13 (1985)141

Agronomic factors
Cultivation of maize in irrigated areas for silage and grain production and their use by ruminants 1(1976)237

Effect of agronomic factors on the yield of maize and on its composition in relation to its use by ruminants 1(1976)251

Agronomic factors affecting the growth and composition of the maize plant 1(1976)263

Soil chemistry and mineral problems in farm livestock. A review 5(1980)95

The quality of the green fraction of leaf protein concentrate from *Chenopodium quinoa* Willd. grown at different levels of fertilizer nitrogen 11(1984)239

The nutritional value of crop residue components from several wheat cultivars grown at different fertilizer levels 11(1984)301

Estimating yield and nutrient losses due to rainfall on field-drying alfalfa hay 16(1986/1987)7

Agropyron elongatum
Effect of lactate on the in vitro digestion of *Agropyron elongatum* by rumen microorganisms 16(1986/1987)161

Albumen quality
Effects of dietary maize solubles and minerals on albumen quality of fresh and stored brown-shelled eggs 8(1983)177

Alcaligenes eutrophus
Nitrogen metabolism in broiler chickens consuming the bacterial strain *Alcaligenes eutrophus* 5(1980)241

Alfalfa
Microwave drying of alfalfa compared to field- and oven-drying: effects on forage quality 16(1986/1987)1

Microfloral and compositional changes in alfalfa hay treated with sodium diacetate and stored at different moisture contents 17(1987)45

Composition and fibre digestion in morphological components of an alfalfa–timothy sward 19(1988)135

Effect of soft white wheat addition to alfalfa-grass forage on heifer gain, diet digestibility and in vitro digestion kinetics 24(1989)141

Alkali treatment
The effect on digestibility of a new technique for alkali treatment of straw 1(1976)73

Review article: the alkali treatment of straws 2(1977)105

Enzyme solubility – a method for evaluating the digestibility of alkali-treated straw 2 (1977)205

The intake of steers and heifers fed on mixtures of alkali-treated or untreated straw with grass silage 2(1977)315

Ensiled alkali-treated straw. I. Effect of level and type of alkali on the composition and digestibility in vitro of ensiled barley straw 3(1978)117

Ensiled alkali-treated straw. II. The nutritive value for young beef cattle of mixtures of ensiled or frozen alkali-treated straw and ryegrass silage 3(1978)133

Alkali treatment of straw for ruminants. I. Utilization of complete diets containing straw by beef cattle 3(1978)143

Cellulase insoluble fibre as a measure of unavailable organic matter in cattle compounds containing alkali-treated straw 3 (1978)227

Akali treatment of straw for ruminants. II. Nutritive value of straw ensiled after alkali treatment. 3(1978)289

Laboratory evaluation of ensiled mixtures of alkali-treated cattle excreta and whole-crop maize 3(1978)335

Effect of time of storage and method of drying on the digestibility in vitro of alkali-treated barley straw 3(1978)381

Effect of level of sodium hydroxide treatment and volume of solution on the nutritive value of maize cobs 4(1979)1

Performance of heifers fed on diets based on NaOH-treated maize cobs and the effect of supplementary urea and source of carbohydrates 4(1979)97

Investigations of the treatment of sawdust for rabbit feeding. 1. Effect of sodium hydroxide treatment 6(1981)43

Titratable groups and soluble phenolic compounds as indicators of the digestibility of chemically treated roughages 6(1981)123

Simplified method for alkali treatment of low-quality roughages for use by small-holders in developing countries 6(1981)133

Utilization of alkali-treated grain. 1. Alkali-treated grain in complete diets for steers and lambs 6(1981)273

Utilization of alkali-treated grain. 2. Utilization by steers of diets based on hay or straw and mixed with either NaOH-treated or rolled barley 6(1981)347

Utilization of alkali-treated grain. 3. Utilization by steers of NaOH-treated and rolled barley in silage-based diets 6(1981)355

Ammonia and sodium hydroxide treatment of wheat straw in diets for fattening steers 7 (1982)1

Effect of alkali-treated rice straw supplemented with spent tea leaf and thyroprotein on milk yield, milk composition and certain physiological parameters of dairy cows 7(1982)201

Note on the lysinoalanine content of alkali-treated barley 7(1982)217

Intake and digestibility in sheep fed on alkali-treated straw and concentrates, as complete diets or separate feeds 7(1982)271

Digestibility of diets containing increasing levels of NaOH-treated or untreated wheat straw 7(1982)331

Methods of improving the utilization of cereal straw by ruminants. I. Supplements of urea, molasses and dried grass and treatment with sodium hydroxide 8(1983)221

Nutritive value for ruminants of sugar cane bagasse ensiled after spray treatment with different levels of NaOH 9(1983)1

Utilization of alkali-treated and neutralized wheat straw-based rations for growing goat kids 9(1983)155

Feeding value of alkali-treated whole rye (*Secale cereale* L.) grain for lactating cows and its digestibility for sheep 10(1983/1984)77

Effect of sodium acrylate and calcium and sodium hydroxides on the fermentation, digestibility, chemical composition and aerobic deterioration of whole-crop maize silage 12(1984/1985)91

Effect of sodium hydroxide treatment and/or extrusion cooking on the nutritive value of peanut hulls 12(1984/1985)187

A comparison of alkali treatment methods to improve the nutritive value of straw. I. Digestibility and metabolizability 12(1984/1985)295

Organic waste ash as possible source of alkali for animal feed treatment 13(1985)237

Effects of feeding alkali-treated olive cake on intake, digestibility and rumen liquor parameters 14(1986)139

A comparison of alkali treatment methods used to improve the nutritive value of straw. II. In sacco and in vitro degradation relative to in vivo digestibility 14(1986)215

Corrugated cardboard and dried poultry manure as dietary ingredients for sheep 14 (1986)265

The response of screened manure fibre to ozone and sodium hydroxide treatments 15 (1986)47

The effect of grinding groundnut hulls either with or without alkali treatment on digestibility of diet and on ruminal and blood components 15(1986)69

The nutritive value of barley for monogastric animals and for ruminants as affected by artificial drying and alkali and anaerobic storage 15(1986)285

Effect of sodium hydroxide treatment of desert grass on carbohydrate digestibility by goats 17(1987)219

Digestibility and beef performance after feeding NaOH- or KOH-treated pig wastes 17 (1987)235

Treatment of whole-crop cereals with alkali. 1. The influence of sodium hydroxide and ensiling on the chemical composition and in vitro digestibility of rye, barley and wheat crops harvested at increasing maturity and dry matter content 18(1987)257

Treatment of whole-crop cereals with alkali. 2. Voluntary intake and digestibility by sheep of rye, barley and wheat crops ensiled with sodium hydroxide 18(1987)271

Treatment of whole-crop cereals with alkali. 3. Voluntary intake and digestibility studies in sheep given immature wheat ensiled with sodium hydroxide, urea or ammonia 18(1987)283

Treatment of whole-crop cereals with alkali. 4. Voluntary intake and growth in steers given

wheat ensiled with sodium hydroxide, urea or ammonia 19(1988)55

Fermentation and utilization by lambs of sugarcane harvested fresh and ensiled with and without NaOH. 4. Ruminal kinetics 23 (1989)323

Alkali-treated soya flour for piglets 24(1989)1

Upgrading barley straw for goats: the effectiveness of a sodium hydroxide and urea dip method 24(1989)57

Alkaloid content

The alkaloid content of sweet lupin seed used in feeding trials on pigs and rats 2(1977)59

Almond hulls

Nutritive value of almond hulls for sheep 8 (1983)63

Amaranthus hybridus

Performance of calves on diets containing *Amaranthus* leaf meal 5(1980)193

Amino acids, essential

Utilization of processed pig hair in poultry diets 3(1978)243

Nitrogen metabolism in broiler chickens consuming the bacterial strain *Alcaligenes eutrophus* 5(1980)241

Estimation of optimum supplements of essential amino acids to a wheat and barley-based diet in rats using a nitrogen balance technique 15(1986)21

Palm kernel meal as a feed for poultry. 1. Composition of palm kernel meal and availability of its amino acids to chicks 16(1986/1987)179

Amino acids, general

Amino acid supplementation of methanol-grown dried microbial cells in diets for young chicks 4(1979)63

Ammonia treatment

Detoxification of aflatoxins in groundnut meal 2(1977)67

Effect on digestibility and nitrogen content of barley straw of different ammonia treatments 2(1977)131

The effect of ammoniation on the intake and nutritive value of straw 4(1979)169

Nutritive value of thermoammoniated and steam-treated maize stover I. Intake, digestibility and nitrogen retention 4(1979) 177

Nutritive value of thermoammoniated and steam-treated maize stover II. Rumen metabolites and rate of passage 4(1979)187

Preservation of high-moisture maize – a comparison of gaseous and liquid anhydrous ammonia with methylene-*bis*-propionate 5 (1980)337

Effects of treatment of barley straw with anhydrous ammonia and supplementation with heat-treated protein meals on feed intake and liveweight performance of growing lambs 6(1981)145

New crop residues and forages for Western Canada: assessment of feeding value in vitro and response to ammonia treatment 6(1981)257

Different sources of ammonia for improving the nutritive value of low quality roughages 6(1981)377

Ammonia and sodium hydroxide treatment of wheat straw in diets for fattening steers 7 (1982)1

Studies on ammonia-treated straw. I. The effects of type and level of ammonia, moisture content and treatment time on the digestibility in vitro and enzyme soluble organic matter of oat straw 7(1982)45

Studies on ammonia-treated straw. II. Fixation of ammonia in treated straw by spraying with acids 7(1982)53

Effect of treatment of wheat straw with ammonia and peracetic acid on digestibility in vitro and cell wall composition 7(1982)325

Ammoniation of wheat straw and native grass hay during baling of large round bales 8 (1983)35

The effect of prolonged storage on the digestibility and nitrogen content of ammonia-treated barley straw 8(1983)147

The feeding value for ruminants of straw and whole-crop barley and oats treated with anhydrous or aqueous ammonia or urea 8 (1983)247

The effect of urease enzyme on treatment time and the nutritive value of straw treated with ammonia as urea 8(1983)271

Methods of improving the utilization of cereal straw by ruminants. III. A note on the effect of ensiling straw treated with urea 9(1983)181

Effect of wilting and the additives, straw, molasses and urea on the fermentation pattern of maize silage 9(1983)185

Composition and enzymatic digestibility of oregon grass straws 9(1983)283

The effects of ammonia on the fermentation of chopped sugarcane 9(1983)291

The effect of urea on the preservation and digestibility in vitro of perennial ryegrass 10(1983/1984)49

Digestion of organic matter and nitrogen in ammoniated barley straw 10(1983/1984)65

A note on the OECD collaborative experiment on ammonia-treated straw 10(1983/1984)89

Chemical properties of fibre in relation to nutritive quality of ammonia-treated forages 10(1983/1984)155

Ammonia treatment of straw: methods for treatment and feeding experience in Norway 10(1983/1984)173

Ammonia versus sodium hydroxide treatment of straw for growing cattle 10(1983/1984) 189

A comparison between degradation in vitro and in sacco of constituents of untreated and ammonia-treated barley straw 10(1983/1984)199

Digestibility studies on ammonia-treated straw 10(1983/1984)213

Commercial experience of treating straw with ammonia 10(1983/1984)223

The digestion of untreated and ammonia-treated barley straw in an artificial rumen 10(1983/1984)229

Anhydrous ammonia-treated cereal straw for animal feeding. Some results from the Mediterranean area 10(1983/1984)239

Ammonia-treated barley straw and rolled barley offered either together, in a mixed ration, or successively to beef steers 10 (1983/1984)247

Ammonia treatment of straw via the hydrolysis of urea. I. Effects of dry matter and urea concentrations on the rate of hydrolysis of urea 11(1984)103

Ammonia treatment of straw via the hydrolysis of urea. II. Additions of soya bean (urease), sodium hydroxide and molasses; effects on the digestibility of urea-treated straw 11(1984)115

The effect of anhydrous ammonia and moisture content on the preservation and chemical composition of perennial ryegrass hay 11(1984)159

Influence of heat sterilization and ammoniation on straw composition and degradation by pure cultures of cellulolytic rumen bacteria 12(1984/1985)195

Effect of ammonia treatment and processing of whole cotton plants as sole source of roughage in complete feeds for growing cross-bred calves 13(1985)93

Effects of ethanolamine and ethylenediamine treatments on the composition and degradability of wheat straw 13(1985)131

Methods of improving the utilization of cereal straw by ruminants. IV. The effect on intake of level of supplementation with urea and dried grass 13(1985)281

Methods of improving the utilization of cereal straw by ruminants. V. The effect of molasses and dried grass 13(1985)293

The use of barley straw as a feed 14(1986)29

Treatment of barley straw with ammonia and sulphur dioxide gases under laboratory conditions 14(1986)41

Effect of ammonia level and time of exposure to ammonia on nutritional and preservatory characteristics of dry and high-moisture coastal bermuda grass hay 14 (1986)55

Urea as a source of ammonia for improving the nutritive value of wheat straw 14(1986) 67

Aqueous ammonia treatment of wheat straw: voluntary intake and digestibility in cattle 14(1986)81

Effects of ammonia treatment and processing of cotton straw as the sole source of roughage in complete feeds on rumen characteristics with Murrah buffaloes 14(1986) 95

Near infrared reflectance spectroscopy of untreated and ammoniated barley straw 15 (1986)189

Chemical composition and digestion in vitro of orchardgrass hay ammoniated by different techniques 17(1987)109

Hyperexcitability in cattle fed ammoniated roughages 17(1987)121

Alkaline hydrogen peroxide treatments of crop residues to increase ruminal dry matter disappearance in sacco 17(1987)179

Treatment of whole-crop cereals with alkali. 3. Voluntary intake and digestibility studies in sheep given immature wheat ensiled with sodium hydroxide, urea or ammonia 18(1987)283

Treatment of whole-crop cereals with alkali. 4. Voluntary intake and growth in steers given wheat ensiled with sodium hydroxide, urea or ammonia 19(1988)55

Effects of ammonia and sulphur dioxide gases on the composition and digestion of barley straw 19(1988)121

The effect of ammoniation on the nutritive value of wheat, barley and oat straws. I. Changes in chemical composition in relation to digestibility in vitro and cell wall degradability 19(1988)159

The effect of ammoniation on the nutritive value of wheat, barley and oat straws.

II. Digestibility and energy value measurements in vivo and their prediction from laboratory measurements 19(1988)173

Fibre degradation in the rumens of animals receiving hay, untreated or ammonia-treated straw 19(1988)277

The effect of variation in the water content of wheat straw before ammoniation on its nutritive value for beef cattle 20(1988)111

The effect of treatment with urea, sorbic acid, or dehydration on orange peel silage 20 (1988)335

The feeding value of ammoniated flax straw, wheat straw and wheat chaff for beef cattle 21(1988)57

Near infra-red spectra of ammonia-treated straw and of isolated cell walls 21(1988)209

Upgrading of straw by ammoniation 21(1988) 287

Effect of ammoniation and energy supplementation on the utilization of wheat straw by sheep 22(1988)305

Effect of urea treatment on rumen degradation characteristics of rice straws 24(1989) 83

Influence of stage of forage maturity and ammoniation of wheat straw on ruminal degradation of wheat forage tissues 24 (1989)201

Oven and stack ammoniation of grass hays. 1. Changes in chemical composition in relation to digestibility in vitro and cell-wall degradability 24(1989)299

Oven and stack ammoniation of grass hays. 2. Voluntary intake, digestibility and faecal nitrogen excretion in sheep in relation to hay composition 24(1989)313

Effects of urea treatment on chemical composition and in vitro digestibility on meadow hays of Northern Portugal 25(1989)157

Amprolium

Particle size distribution of a feed premix containing amprolium 3(1978)211

Amylase

Prediction of the digestibility of ruminant feeds from their solubility in enzyme solutions 2(1977)153

Antibiotics

A note on the effect on the aerobic deterioration of maize silage of the manipulation of the microflora by means of antibiotics 3 (1978)89

Efficacy of virginiamycin and a commercially-available lactobacillus probiotic in swine diets 8(1983)69

Apple pomace

Feeding value of apple pomace silage for sheep 11(1984)189

Some properties of apple pomace ensiled with and without additives 15(1986)57

Nutritive value of apple pomace for ruminants 16(1986/1987)311

Arginine

Limiting amino acids in dried microbial cells given to young turkeys 4(1979)209

Arhar

Nutritional evaluation of gram (*Cicer arietinum*) and arhar (*Cajanus cajan*) chunies for cattle 5(1980)315

Artichoke

The chemical composition and nutritive value for ruminants of four possible energy crops and their residues of anaerobic fermentation 15(1986)197

Aspergillus fumigatus

Production of fungal protein from rasped fresh cassava roots using 200- and 3000-litre fermentors 8(1983)313

Avoparcin

Effect of avoparcin on performance, body composition and meat quality of finishing bulls 8(1983)237

In vitro effects of avoparcin on protein degradability and rumen fermentation 15 (1986)215

B

Bacteria

A note on the effect on the aerobic deterioration of maize silage of the manipulation of the microflora by means of antibiotics 3 (1978)89

Nitrogen metabolism in broiler chickens consuming the bacterial strain *Alcaligenes eutrophus* 5(1980)241

Protein quality studies on rats fed on cassava-root meal enriched with *Cephalosporium eichhorniae* 152 or with conventional plant-protein supplements 6(1981)197

The effect of some pre-ensiling treatments on silage composition and nitrogen disappearance in the rumen 8(1983)259

Electron microscopy of bacteria involved in the digestion of plant cell walls 10(1983/ 1984)93

Effect of absence of ciliate protoza on rumen fluid volume, flow rate and bacterial populations in sheep 10(1983/1984)145

Stability of viable lactobacillus products 11 (1984)261

Influence of heat sterilization and ammoniation on straw composition and degradation by pure cultures of cellulolytic rumen bacteria 12(1984/1985)195

Effects of molasses/urea and bacterial inoculant additives on silage quality, dry matter recovery, and feeding value for cattle 12 (1984/1985)205

The chemical composition of grass silages prepared with and without pre-treatment with inoculants containing *Lactobacillus plantarum* 13(1985)269

Evaluation of a dual flow continuous culture system for estimating bacterial fermentation in vivo of mixed diets containing various soya bean products 16(1986/1987)51

Population changes of fibrolytic rumen bacteria in the presence of phenolic acids and plant extracts 19(1988)261

The establishment of rumen microbial communities 21(1988)69

Possibilities of the use of recombinant-DNA techniques with rumen micro-organisms 21 (1988)161

Characterization of the flora of the large bowel of pigs: a status report 23(1989)79

Fibre-degrading microorganisms in the monogastric digestive tract 23(1989)91

Assessing the importance of host-derived polysaccharides as carbon sources for bacteria growing in the human colon 23(1989) 109

In vitro fermentation of dietary fiber by human fecal organisms 23(1989)121

Inoculation of corn silage and high-moisture corn with lactic acid bacteria and its effects on the subsequent fermentations and on feedlot performance of beef steers 25(1989) 23

Evaluation of various methods for protecting soya bean protein from degradation by rumen bacteria 25(1989)111

In vitro metabolism of mycotoxins by bacterial, protozoal and ovine ruminal fluid preparations 25(1989)169

Bagaruwa seeds

Biochemical evaluation of bagaruwa seeds (*Acacia nilotica*) for use as livestock feed 11(1984)45

Balanites kernel cake

The use of Balanites kernel cake in a diet for fattening sheep 9(1983)301

Banana stalk

The value of banana stalk as a feed for goats 15(1986)95

Nutritive value of banana stalk (*Musa cavendishi*) as a feed for sheep 22(1988)327

Barley

Lupin-seed meal (*Lupinus angustifolius* cv. Uniwhite) as a protein supplement to barley-based diets for growing pigs 1(1976) 631

Swedes (*Brassica napus*) for growing pigs: chemical composition and use as a replacement for barley in the diet 2(1977)31

A comparison between meals prepared from the seeds of different varieties of lupin as protein supplements to barley-based diets for growing pigs 2(1977)49

The effect of supplementary protein on the performance of pigs given swedes (*Brassica napus*) as a replacement for barley in the diet 2(1977)277

The use of whole barley diets fortified with solutions of urea, minerals and vitamins for lambs 2(1977)307

Nutritive value of leaf protein coagulum and cereals mixtures 4(1979)91

The determination of available lysine in barley using *Tetrahymena pyriformis* W. 5 (1980)33

A comparison of three methods for the determination of available lysine in barley 5 (1980)43

Utilization of alkali-treated grain. 1. Alkali-treated grain in complete diets for steers and lambs 6(1981)273

Nutritional evaluation, using rats, of diets based on cereals and ethanol-grown yeast 6(1981)309

Utilization of alkali-treated grain. 2. Utilization by steers of diets based on hay or straw and mixed with either NaOH-treated or rolled barley 6(1981)347

Utilization of alkali-treated grain. 3. Utilization by steers of NaOH-treated and rolled barley in silage-based diets 6(1981)355

Influence of increasing levels of β-glucanase on the productive value of barley diets for broiler chickens 7(1982)351

Nutritional value of waxy barleys for broiler chickens 8(1983)25

Evaluation of optimum lysine and threonine supplements to a wheat and barley-based diet in rats 8(1983)163

The effect of calcium and sodium hydroxides and of sodium acrylate on the fermentation

and digestibility in vitro of ensiled whole-crop wheat and barley harvested at different stages of maturity 9(1983)37

Effect of storage and processing of barley on its utilization by pigs and rats 9(1983)89

Effects of milling and pelleting of maize, barley and wheat on their metabolizable energy value for cockerels and chicks 9(1983) 99

Associative effects of mixed feeds. II. The effect of dietary addition of bicarbonate salts on the voluntary intake and digestibility of diets containing various proportions of hay and barley 10(1983/1984)31

Anaerobically stored high moisture cereals for growing pigs 12(1984/1985)159

Feeding value of hydroponically sprouted barley for poultry and pigs 13(1985)183

Nutrient content of hydroponically sprouted barley 13(1985)191

Apparent ileal and faecal digestibility of amino acids in barley-based diets supplemented with soya bean meal or canola meal for growing pigs 14(1986)183

The effect of monensin on rumen fermentation in sheep fed on all-roughage and barley/roughage diets 15(1986)7

Estimation of optimum supplements of essential amino acids to a wheat and barley-based diet in rats using a nitrogen balance technique 15(1986)21

The effect of β-glucanase on the utilization of starch and nitrogen by broiler chickens fed on barley of low- or high-viscosity 15(1986) 83

The nutritive value of barley for monogastric animals and for ruminants as affected by artificial drying and alkali and anaerobic storage 15(1986)285

Fermentation of hammermilled barley, lupin, pea and faba bean grain in the rumen of dairy cows 16(1986/1987)261

Nutritional value for chickens of dried distillers-spent-grain from barley and dehulled barley 17(1987)145

The effect of dietary barley on carbohydrate digestibility of sulphur dioxide-treated wheat straw by sheep 18(1987)55

Treatment of whole-crop cereals with alkali. 1. The influence of sodium hydroxide and ensiling on the chemical composition and in vitro digestibility of rye, barley and wheat crops harvested at increasing maturity and dry matter content 18(1987)257

Treatment of whole-crop cereals with alkali. 2. Voluntary intake and digestibility by sheep of rye, barley and wheat crops ensiled with sodium hydroxide 18(1987)271

The nutritive value of hulless barley for swine 19(1988)191

The effects of barley, unmolassed sugar-beet pulp and molasses supplements on organic matter, nitrogen and fibre digestion in the rumen of cattle given a silage diet 20(1988) 259

Amino acids flow to and absorption from the small intestine of sheep fed barley and sulfur dioxide-treated straw at different ratios 22(1988)147

A comparison of wheat- or barley-based diets given ad libitum as meal or pellets to finishing pigs 22(1988)263

The effect of mechanical treatment on the nutritive value of high-moisture barley and the performance of growing-finishing pigs 24(1989)121

Effect of enzyme supplementation on the nutritive value of hulless barley in chicken diets 24(1989)233

Factors effecting the digestibility by pigs of energy and protein in wheat, barley and sorghum diets supplemented with canola meal 24(1989)253

Effect of formaldehyde treatment on the rumen digestion of barley and wheat grain and its influence on hay intake and digestion by sheep 25(1989)55

The effect of β-glucanase supplementation on nutrient digestibility and growth in broilers given diets containing barley, oat groats or wheat 25(1989)193

Barley grain

Note on the lysinoalanine content of alkali-treated barley 7(1982)217

Milling barley and lupin grain in diets for cattle 12(1984/1985)57

Microbial dynamics in barley grain stored under controlled atmosphere 19(1988)299

Barley meal

The influence of feeding level and level of dietary inclusion on the digestibility of barley meal in the pig 2(1977)41

Barley straw

Effect on digestibility and nitrogen content of barley straw of different ammonia treatments 2(1977)131

Ensiled alkali-treated straw. I. Effect of level and type of alkali on the composition and digestibility in vitro of ensiled barley straw 3(1978)117

Alkali treatment of straw for ruminants. I. Utilization of complete diets containing straw by beef cattle 3(1978)143

Akali treatment of straw for ruminants. II. Nutritive value of straw ensiled after alkali treatment. 3(1978)289

Effect of time of storage and method of drying on the digestibility in vitro of alkali-treated barley straw 3(1978)381

Effects of treatment of barley straw with anhydrous ammonia and supplementation with heat-treated protein meals on feed intake and liveweight performance of growing lambs 6(1981)145

Studies on ammonia-treated straw. II. Fixation of ammonia in treated straw by spraying with acids 7(1982)53

The feeding value for ruminants of straw and whole-crop barley and oats treated with anhydrous or aqueous ammonia or urea 8 (1983)247

The effect of urease enzyme on treatment time and the nutritive value of straw treated with ammonia as urea 8(1983)271

Digestion of organic matter and nitrogen in ammoniated barley straw 10(1983/1984)65

A comparison between degradation in vitro and in sacco of constituents of untreated and ammonia-treated barley straw 10(1983/1984)199

Digestibility studies on ammonia-treated straw 10(1983/1984)213

The digestion of untreated and ammonia-treated barley straw in an artificial rumen 10(1983/1984)229

Ammonia-treated barley straw and rolled barley offered either together, in a mixed ration, or successively to beef steers 10 (1983/1984)247

A comparison of alkali treatment methods to improve the nutritive value of straw. I. Digestibility and metabolizability 12(1984/1985)295

Treatment of barley straw with ammonia and sulphur dioxide gases under laboratory conditions 14(1986)41

A comparison of alkali treatment methods used to improve the nutritive value of straw. II. In sacco and in vitro degradation relative to in vivo digestibility 14(1986)215

Evaluation of the increase in feeding value of barley straw following mild treatment with sulphur dioxide 15(1986)13

Near infrared reflectance spectroscopy of untreated and ammoniated barley straw 15 (1986)189

Effect of straw treatment and laboratory of incubation on degradation of neutral detergent residue in the rumen 18(1987)75

Fresh and dried brewer's grains as protein supplements to barley straw diets given to pregnant beef cows 19(1988)33

Effects of ammonia and sulphur dioxide gases on the composition and digestion of barley straw 19(1988)121

The effect of ammoniation on the nutritive value of wheat, barley and oat straws. I. Changes in chemical composition in relation to digestibility in vitro and cell wall degradability 19(1988)159

The effect of ammoniation on the nutritive value of wheat, barley and oat straws. II. Digestibility and energy value measurements in vivo and their prediction from laboratory measurements 19(1988)173

Fibre degradation in the rumens of animals receiving hay, untreated or ammonia-treated straw 19(1988)277

The effect of five different supplements on the degradation of straw in sheep given untreated barley straw 19(1988)289

Upgrading of straw by ammoniation 21(1988) 287

Effect of soaking, ensilage and hydrogen peroxide treatment of barley straw on rumen degradability 22(1988)295

Upgrading barley straw for goats: the effectiveness of a sodium hydroxide and urea dip method 24(1989)57

Basic steel slag

Biological availability of phosphorus from basic steel slag 11(1984)75

Bean, winged

The proximate composition and nutritive value of the winged bean *Psophocarpus tetragonolobus* (L.) DC for broilers 11 (1984)231

Bermuda grass

Effect of ammonia level and time of exposure to ammonia on nutritional and preservatory characteristics of dry and high-moisture coastal bermuda grass hay 14 (1986)55

Bermudagrass

Intake, digestion and daily gain by cattle consuming bermudagrass (*Cynodon dactylon*) and supplemented with different combinations of ground corn, vegetable oil, uera and corn gluten and blood meals 25(1989)99

β-glucanase
Influence of increasing levels of β-glucanase on the productive value of barley diets for broiler chickens 7(1982)351

The effect of β-glucanase on the utilization of starch and nitrogen by broiler chickens fed on barley of low- or high-viscosity 15(1986) 83

The effect of β-glucanase supplementation on nutrient digestibility and growth in broilers given diets containing barley, oat groats or wheat 25(1989)193

Bicarbonate salts
Associative effects of mixed feeds. II. The effect of dietary addition of bicarbonate salts on the voluntary intake and digestibility of diets containing various proportions of hay and barley 10(1983/1984)31

Blood
The composition of processed abattoir blood and its use in finishing-pig diets 2(1977)21

Treatment of the haem fraction of abattoir blood with formic acid to produce a solid state, low dry matter protein supplement for use in finishing-pig diets 2(1977)367

Blood meal as a source of protein for grower/finisher pigs 3(1978)191

Ensiling of rumen contents and blood with wheat straw 18(1987)67

Comparison of cottonseed cake and dried blood as nitrogen sources for growing Sudan Nubian goats 22(1988)333

Blood meal
Protein quality of mixtures of animal protein meals 9(1983)79

Effect of blood meal processing method on performance and nitrogen utilization by growing cattle 24(1989)337

Bone meal
Protein quality of mixtures of animal protein meals 9(1983)79

Meat and bone and fish meals in balancer feeds for choice-fed broilers 11(1984)221

Botanical composition
Botanical composition and extent of lignification affecting digestibility of wheat and oat straw and paspalum hay 7(1982)71

Brewer's dried grains
Influence of the drying process on the composition of brewer's dried grains 6(1981)163

Evaluation of brewer's dried grains in the diets of laying hens 6(1981)169

Evaluation of brewer's dried grains in the diets of broiler chickens 6(1981)179

Fresh and dried brewer's grains as protein supplements to barley straw diets given to pregnant beef cows 19(1988)33

Buckwheat
A nutritional evaluation of buckwheat (*Fagopyrum esculentum*) 3(1978)95

Buffalo
Nutritive evaluation of uromol–wheat straw as a basal ration for buffaloes 13(1985)103

Effects of ammonia treatment and processing of cotton straw as the sole source of roughage in complete feeds on rumen characteristics with Murrah buffaloes 14(1986) 95

Colonization of guinea grass by anaerobic rumen fungi in swamp buffalo and cattle 22 (1988)161

Buffers
Comparison between two methods of feeding concentrate to sheep given a diet high or low in concentrate with or without buffering substances 16(1986/1987)37

Comparison of buffers on rumen functions, turnover rate and gastric secretions in Holstein steers 17(1987)257

Efficacy of supplemental dietary neutralizing agents for lactating dairy cows. A review 23(1989)277

By-products–animal waste
The composition of processed abattoir blood and its use in finishing-pig diets 2(1977)21

Nutritive value of rumen contents for monogastric animals 2(1977)351

Utilization by sheep of dried pig faeces containing a high concentration of copper 3 (1978)31

Digestibility by cattle of diets containing dried pig faeces 3(1978)155

Utilization of processed pig hair in poultry diets 3(1978)243

Protein digestibility of hydrolyzed hog hair meal for pigs 3(1978)323

The effects of air dried poultry manure on performance, organ weights and haematological measurements of weaner pigs reared in the tropics 4(1979)23

Utilization of cattle manure for single-cell protein production with *Chaetomium cellulolyticum* 4(1979)199

Digestibility of wet cage layer excreta and an excreta-supplemented diet by cattle as determined by two methods 5(1980)77

Utilization of cattle manure for Torula yeast production from straw hydrolysates 5(1980) 175

Performance of fattening cattle and lactating beef cows on diets high in chemically

treated wheat straw and poultry litter 5 (1980)299

Investigations of the treatment of sawdust for rabbit feeding. 2. Effect of boiling and incubation with poultry manure 6(1981)51

Fermentation characteristics and feeding value of ensiled poultry litter containing wheat straw, bagasse or sawdust 7(1982)341

Nutritive value of ensiled broiler litter for cattle 8(1983)19

The utilization by broiler chickens of poultry offal hydrolysed in formic acid 11(1984)247

Chemical preparation of NaOH–keratin hydrolysate for improving the nutritive value of wheat straw 13(1985)215

Thermofractographic determination of monocarbonyl compounds in animal waste fats used as feed fat 13(1985)227

Organic waste ash as possible source of alkali for animal feed treatment 13(1985)237

Effects of processing time and moisture content on amino acid composition and nitrogen characteristics of feather meal 14(1986) 279

The utilisation by growing pigs of ruminant offal hydrolysed in formic acid 15(1986)273

The effect of enzymatic treatment on amino acid content and nitrogen characteristics of feather meal 16(1986/1987)151

Digestibility and beef performance after feeding NaOH- or KOH-treated pig wastes 17 (1987)235

Ensiling of rumen contents and blood with wheat straw 18(1987)67

Evaluation of cardboard and dried poultry waste as feed ingredients for ruminants 18 (1987)209

Animal excreta as ruminant feed – scope and limitations under Indian conditions 19 (1988)1

Slaughterhouse by-products preserved by *Lactobacillus plantarum* fermentation as feed for mink and foxes 20(1988)287

Nutritive value of rice straw (*Oryza sativa*) ensiled with animal excreta and rumen digesta 24(1989)69

Ensiling characteristics of cattle waste fermented with wheat straw and green berseem (*Trifolium alexandrium*) 25(1989)149

By-products–apple pomace

Some properties of apple pomace ensiled with and without additives 15(1986)57

By-products–banana stalk

The value of banana stalk as a feed for goats 15(1986)95

By-products–brewer's dried grains

Influence of the drying process on the composition of brewer's dried grains 6(1981)163

Evaluation of brewer's dried grains in the diets of laying hens 6(1981)169

Evaluation of brewer's dried grains in the diets of broiler chickens 6(1981)179

By-products–canning industry residues

By-products from the canning industry as feedstuff for ruminants: digestibility and its prediction from chemical composition and laboratory bioassays 25(1989)67

By-products–cardboard

Evaluation of cardboard and dried poultry waste as feed ingredients for ruminants 18 (1987)209

By-products–cocoa

Studies on the feeding value of agro-industrial by-products. III. Digestibility of cocoa-pod and cocoa-pod-based diets by ruminants 13 (1985)249

Studies on the feeding value of agro-industrial by-products: strategies for improving the utilisation of cocoa-pod-based diets by ruminants 20(1988)189

By-products–coffee residue

A note on the nutritive value of dried instant coffee residue for broiler chickens and turkey poults 10(1983/1984)285

The nutritional value of dried coffee pulp (DCP) in broiler chickens's diets 22(1988) 139

By-products–cottonseed

A review of the nutritive value and utilization of whole cottonseed, cottonseed meal and associated by-products by dairy cattle 18 (1987)89

By-products–crop residues

New crop residues and forages for Western Canada: assessment of feeding value in vitro and response to ammonia treatment 6(1981)257

Improving utilization of crop residues. Evaluation of two chemicals used to improve digestibility of crop residues 9(1983)205

The nutritional value of crop residue components from several wheat cultivars grown at different fertilizer levels 11(1984)301

Alkaline hydrogen peroxide treatments of crop residues to increase ruminal dry matter disappearance in sacco 17(1987)179

Supplementation of rice straw with crop by-products. I. Legume straw supplementation 19(1988)43

Nutritive value of banana stalk (*Musa cavendishi*) as a feed for sheep 22(1988)327

By-products–culinary waste
Chemical composition of culinary wastes and their potential as a feed for ruminants 5 (1980)205

By-products–distillery waste
Distillery effluents as animal feed: the use of condensed beet molasses stillage (CBMS) in broiler feeding 5(1980)233

Nitrogen balance and abomasal protein-nitrogen flow in growing ruminants fed methane digester effluent in combination with distillers dried grains 13(1985)47

Nutritional value for chickens of dried distillers-spent-grain from barley and dehulled barley 17(1987)145

By-products–dried bakery product
Variation in nutrient content of samples of dried bakery product 7(1982)419

By-products–fish silage
Fish silage as a protein supplement for early-weaned calves 17(1987)165

By-products–general
Suitability of cell-wall constituents as predictors of organic matter digestibility in some tropical and subtropical by-products 7(1982)191

Variations and prevision of the in sacco dry matter digestion of concentrates and by-products 13(1985)7

Use of the rumen simulation technique (RUSITEC) to compare microbial digestion of various by-products 21(1988)197

By-products–grapevine
Digestibility in vivo of ensiled grapevines (branches and leaves): influence of the system of analysis in the detergent fibre scheme on the prediction of digestibility 22 (1988)173

By-products–maize whole stillage
Maize whole stillage as a supplemental protein source for growing lambs and finishing steers 11(1984)181

By-products–oil industry
Composition of by-products of the essential-oil industry and their potential as feeds for ruminants 6(1981)115

By-products–olive cake
Effects of feeding alkali-treated olive cake on intake, digestibility and rumen liquor parameters 14(1986)139

By-products–orange peel
The effect of treatment with urea, sorbic acid, or dehydration on orange peel silage 20 (1988)335

By-products–palm oil mill effluent
The intake and digestibility in goats of *Leucaena leucocephala* supplemented with dehydrated palm oil mill effluent 24(1989)45

By-products–sal seed meal
Tannins in sal seed (*Shorea robusta*) and sal seed meal limit their utilization as livestock feeds 7(1982)161

By-products–sawdust
Investigations of the treatment of sawdust for rabbit feeding. 1. Effect of sodium hydroxide treatment 6(1981)43

Investigations of the treatment of sawdust for rabbit feeding. 2. Effect of boiling and incubation with poultry manure 6(1981)51

By-products–silage effluent
The use of effluent from grass silage in the diet of finishing pigs. I. Variation in composition of effluents 4(1979)263

The use of effluent from grass silage in the diet of finishing pigs. II. Assessment of nutritive value of fresh and stored effluent 4(1979) 275

Investigations on the effect of sodium bentonite and other moisture absorbents on the production of effluent from grass silage 8(1983)107

By-products–sisal waste
Feed value of Tanzanian sisal wastes: chemical composition and digestibility in vitro 14 (1986)255

By-products–vine
Composition of vine-branches with leaves of *Vitis vinifera* L. fresh and ensiled using different additives (previous results) 16(1986/1987)89

By-products–whey
The nutritive value of liquid whey, either sour or sweet, when given in restricted amounts to the growing pig 3(1978)163

The evaluation of fresh sweet liquid whey as a protein supplement to maize for growing pigs 14(1986)171

By-products–wood residue
Composition of kermes oak browse as affected by shade and stage of maturity 21(1988)1

Pilot plant production of sulphur dioxide-treated mesquite wood. II. Digestibility and feedlot performance by lambs 22(1988) 61

C

Cabbage
Cabbage (*Brassica oleracea*) in the diet of growing–finishing pigs 5(1980)69

Studies on the nutritive value of roots, cabbage and grass silage for growing commercial rabbits 13(1985)299

Calcium
Fractions of nitrogen, sulphur, phosphorus, calcium and magnesium in the herbage of perennial ryegrass as influenced by fertilizer nitrogen 14(1986)231

Compositional quality of certain uncoventional calcium and phosphorus sources in India for use as mineral supplements for livestock 23(1989)343

Calcium, magnesium
Solubility and flow of calcium, magnesium and phosphorus in the digestive tract of sheep given maize or alfalfa silages 9(1983) 131

Calcium hydroxide treatment
The effect of calcium hydroxide treatment on the digestibility of chopped oat straw supplemented with a solution containing urea, calcium, phosphorus, sodium, trace elements and vitamins 7(1982)93

The effect of calcium and sodium hydroxides and of sodium acrylate on the fermentation and digestibility in vitro of ensiled whole-crop wheat and barley harvested at different stages of maturity 9(1983)37

Effect of sodium acrylate and calcium and sodium hydroxides on the fermentation, digestibility, chemical composition and aerobic deterioration of whole-crop maize silage 12(1984/1985)91

The utilization of untreated and calcium hydroxide treated wheat straw by sheep 12 (1984/1985)141

Digestion rates in and outflow rates from the rumen of sheep fed untreated or calcium hydroxide-treated wheat straw 25(1989) 179

Calcium soap
The effects on feed intake, growth and digestibility of nutrients of including calcium soaps of fat in diets for young calves 14 (1986)103

Calcium-soaps
Feed intake, milk yield and milk composition by replacing unprotected fat by Ca-soaps for dairy cows 22(1988)193

Calorimetry
A calorimetric investigation of the nutritive value of formic acid- and formaldehyde–formic acid-treated grass silages 6(1981)245

Canavanine
Canavanine content and toxicity of sesbania leaf meal for growing chicks 25(1989)137

Cane
Proximate chemical analysis of final cane molasses and effect of feeding 30% molasses on intestinal sucrase and maltase activities in the rat 3(1978)15

Canola meal
Apparent ileal and faecal digestibility of amino acids in barley-based diets supplemented with soya bean meal or canola meal for growing pigs 14(1986)183

Canola meal as a protein supplement for growing–finishing pigs 18(1987)37

The nutritive value of canola meal for early-weaned pigs 18(1987)45

Factors effecting the digestibility by pigs of energy and protein in wheat, barley and sorghum diets supplemented with canola meal 24(1989)253

Canola rapeseed
The influence of steam pelleting and grinding on the nutritive value of Canola rapeseed for poultry 8(1983)303

Carbon tetrachloride
Long term feeding of concentrate fumigated with carbon tetrachloride to sheep 2(1977) 77

Cardboard
Corrugated cardboard and dried poultry manure as dietary ingredients for sheep 14 (1986)265

Evaluation of cardboard and dried poultry waste as feed ingredients for ruminants 18 (1987)209

Carotenes
Losses of beta-carotene in red clover in an acid medium during ensiling 4(1979)81

The enzymic nature of the degradation of beta-carotene in red clover and in other forage crops during silage-making with acid additives 5(1980)59

Losses of beta-carotene in unwilted forage crops during silage-making and feeding 9 (1983)63

Investigations into the effects of processing on the retention of the carotenoid fractions of *Leucaena leucocephala* during storage, and the effects of processing on mimosine concentration 9(1983)307

Casein
The apparent and true ileal digestibility of nitrogen and amino acids in lactic casein for the growing pig 16(1986/1987)169

Nutritive value of Nigerian rubber seed (*Hevea brasiliensis*) meal and oil. III. Performance characteristics, relative organ weights, hematocrit and plasma metabolites of growing female rats fed corn diets containing rubber seed meal, soya bean meal or casein 20(1988)125

Cassava
Further thermotolerant fungi for the conversion of cassava starch to protein 2(1977)7

Raw and steam-pelleted cassava, sweet potato and yam *cayenensis* as starch sources for ruminant and chicken diets 3(1978)73

Problems in the use of cassava as animal feed 3(1978)345

Cassava meal and cassava peel meal in diets for growing pigs 8(1983)211

Effect of cassava-based diets varying in cyanide content on the performance and physiopathology of the African giant rat (*Cricetomys gambianus* Waterhouse) 11 (1984)1

Cassava for animal feeding: effect of variety and plant age on production of leaves and roots 11(1984)49

Cassava chips
Effect of variety and plant age on the cyanide content of whole-root cassava chips and its reduction by sun-drying 11(1984)57

Cassava flour
The maize replacement value of cassava flour in rations for growing ewe lambs 20(1988) 233

Cassava meal
Protein quality studies on rats fed on cassava-root meal enriched with *Cephalosporium eichhorniae* 152 or with conventional plant-protein supplements 6(1981)197

Influence of processing methods and storage time on the cyanide potential of cassava leaf meal 17(1987)227

Cassava meal as supplement to napier grass diets for growing sheep and goats 18(1987) 197

Effects of oven-dried cassava peel meal on the performance and carcass characteristics of broiler chickens 24(1989)247

Cassava roots
Production of fungal protein from rasped fresh cassava roots using 200- and 3000-litre fermentors 8(1983)313

Cassava tops
Net energy for gain (NE_{gain}) of cassava tops 20(1988)97

Cattle–beef
The use of techniques of maize cultivation to maximise feed for beef production 1(1976) 229

Maize silage with or without NPN, dehydrated whole-crop maize pellets or high moisture maize grain for finishing bulls 1 (1976)347

Maize grain or ears in concentrate diets for young fattening bulls 1(1976)393

Rate of growth, feed utilisation and digestibility observed with diets containing ensiled maize grain or maize ears fed alone or in mixtures 1(1976)401

Utilising the maize crop in various forms for beef production 1(1976)417

Levels, sources and methods of nitrogen supplementation of maize silage for beef production 1(1976)465

Economic aspects of high-plane feeding of bulls with maize silage 1(1976)513

For beef cattle, all forms of maize can be used 1(1976)521

Bull-beef production from the maize crop in Italy in large and small units 1(1976)531

Dried suger beet pulp as a high energy feed for beef cattle 1(1976)643

Abomasal and plasma nitrogen constituents of steers fed on different physical forms of shelled maize 2(1977)225

Alkali treatment of straw for ruminants. I. Utilization of complete diets containing straw by beef cattle 3(1978)143

Performance of fattening cattle and lactating beef cows on diets high in chemically treated wheat straw and poultry litter 5 (1980)299

Utilization of alkali-treated grain. 1. Alkali-treated grain in complete diets for steers and lambs 6(1981)273

Utilization of alkali-treated grain. 2. Utilization by steers of diets based on hay or straw and mixed with either NaOH-treated or rolled barley 6(1981)347

Utilization of alkali-treated grain. 3. Utilization by steers of NaOH-treated and rolled barley in silage-based diets 6(1981)355

Ammonia and sodium hydroxide treatment of wheat straw in diets for fattening steers 7 (1982)1

The digestion in the small intestine of young bulls of the protein of rapeseed meal

treated or untreated with formaldehyde 7 (1982)225

Monensin-sodium as a performance-promoting additive for fattening bulls and its impact on carcass and meat quality characteristics 7(1982)401

Effect of avoparcin on performance, body composition and meat quality of finishing bulls 8(1983)237

Ammonia versus sodium hydroxide treatment of straw for growing cattle 10(1983/1984) 189

Ammonia-treated barley straw and rolled barley offered either together, in a mixed ration, or successively to beef steers 10 (1983/1984)247

Maize whole stillage as a supplemental protein source for growing lambs and finishing steers 11(1984)181

The use of kidney beans (*Phaseolus vulgaris*) as protein supplements in diets for young rapidly growing beef steers 12(1984/1985) 1

Milling barley and lupin grain in diets for cattle 12(1984/1985)57

Improving the intake and nutritive value of wheat straw for beef cows 12(1984/1985) 125

Wheat silage for fattening young male cattle 12(1984/1985)253

Effect of sugarcane condensed molasses solubles, sugarcane molasses and monensin on performance and volatile fatty acid production in finishing steers 12(1984/1985)275

Methods of improving the utilization of cereal straw by ruminants. IV. The effect on intake of level of supplementation with urea and dried grass 13(1985)281

Methods of improving the utilization of cereal straw by ruminants. V. The effect of molasses and dried grass 13(1985)293

Effect of dietary nitrogen source on microbial protein synthesis, dietary protein degradation and nutrient digestion in steers 14 (1986)117

Effects of dietary concentrates on rumen digestion of fibrous feedstuffs 14(1986)193

Feeding value of sprouted wheat (*Triticum aestivum*) for beef cattle finishing diets 15 (1986)113

High grain content maize silage in fattening diets of young male cattle 16(1986/1987)63

Relative intakes, digestibility and bacterial protein synthesis by sheep and cattle fed high-roughage diets 16(1986/1987)75

The effects of different sources of nitrogen supplementation on the digestion of fibre components in the rumen of steers 17(1987) 65

Comparison of buffers on rumen functions, turnover rate and gastric secretions in Holstein steers 17(1987)257

Effects of whole or ground corn with different forms of hay in 85% concentrate diets on digestion and passage rate in beef heifers 18(1987)151

Fresh and dried brewer's grains as protein supplements to barley straw diets given to pregnant beef cows 19(1988)33

Treatment of whole-crop cereals with alkali. 4. Voluntary intake and growth in steers given wheat ensiled with sodium hydroxide, urea or ammonia 19(1988)55

The effect of variation in the water content of wheat straw before ammoniation on its nutritive value for beef cattle 20(1988)111

Effects of ruminal degradability of the protein supplement on the utilization of maize-stover silage by growing beef cattle 20 (1988)241

The utilization by beef steers of a soft high lysine and a hard normal size maize cultivar in whole grain feedlot diets 21(1988)31

The feeding value of ammoniated flax straw, wheat straw and wheat chaff for beef cattle 21(1988)57

Rumen degradability of nitrogen in typical cultivated grasses and leguminous fodders 22(1988)79

Effects of added dietary urea on the utilization of maize stover silage by growing beef cattle 22(1988)227

Effect of level and frequency of maize supplementation on the utilization of low-quality grass hay by beef cows 24(1989)129

Inoculation of corn silage and high-moisture corn with lactic acid bacteria and its effects on the subsequent fermentations and on feedlot performance of beef steers 25(1989) 23

The effect of a protein energy supplement on pasture protein and fibre digestion in the rumen of grazing steers 25(1989)39

Water-drinking frequency and quantity of water drunk by steers fed hay or fresh forage 25(1989)207

Cattle–calf

Methanol-fermentation protein in veal calf nutrition 1(1976)33

Effect of weight at slaughter on the efficiency of fattening young cattle 1(1976)505

The replacement of skim milk by dried microbial cells and whey in milk diets for calves 2(1977)85

Digestibility of protein and amino acids of a fermentation single-cell protein for veal calves 2(1977)377

Ensiled alkali-treated straw. II. The nutritive value for young beef cattle of mixtures of ensiled or frozen alkali-treated straw and ryegrass silage 3(1978)133

Spent tea leaf as a ruminant feed 3(1978)219

Performance of calves on diets containing *Amaranthus* leaf meal 5(1980)193

Effects of sodium chloride and sodium bicarbonate in the diet on the performance of calves 5(1980)309

Performance of calves fed on milk replacers containing fish protein hydrolysate 7(1982) 141

Evaluation of corn gluten meal with urea as a source of supplementary nitrogen for growing calves and lambs 7(1982)375

Effect of ammonia treatment and processing of whole cotton plants as sole source of roughage in complete feeds for growing cross-bred calves 13(1985)93

The effects on feed intake, growth and digestibility of nutrients of including calcium soaps of fat in diets for young calves 14 (1986)103

Utilization of whole rape (Canola) seed and sunflower seeds as sources of energy and protein in calf starter diets 15(1986)101

Effect of formaldehyde-treated soya bean meal and urea in starters on nitrogen quality, degradibility in sacco, sheep digestibility and calf performance 16(1986/1987)287

Fish silage as a protein supplement for early-weaned calves 17(1987)165

Fish protein concentrates for pre-ruminant calves: effects of processing temperatures and source of fish on protein digestibility and biological value 18(1987)181

The effects of method of processing of starters, tallow inclusion and roughage supplementation on the performance of early-weaned calves 19(1988)231

Evaluation of pea protein concentrate as a source of protein in milk replacers for Holstein calves 24(1989)267

Effect of blood meal processing method on performance and nitrogen utilization by growing cattle 24(1989)337

Cattle–dairy

Forage maize in the rations of dairy cows and fattening cattle in Bulgaria 1(1976)45

The relationship between the intake of herbage grazed by dairy cows and its digestibility 1 (1976)53

The digestibility and nutritive value for dairy heifers of rations containing cellulose fibre 2(1977)287

Maize silage and whole maize plant pellets as the basic forages in complete feeds for milk cows 3(1978)23

Performance of heifers fed on diets based on NaOH-treated maize cobs and the effect of supplementary urea and source of carbohydrates 4(1979)97

Nutritional evaluation of gram (*Cicer arietinum*) and arhar (*Cajanus cajan*) chunies for cattle 5(1980)315

Dry matter, protein, energy and fibre intake by dairy heifers grazing a Rhodes grass (*Chloris gayana*) pasture 6(1981)15

Feed consumption and performance of dairy cows with alternate feedings of grass silage and maize silage 6(1981)337

Effect of alkali-treated rice straw supplemented with spent tea leaf and thyroprotein on milk yield, milk composition and certain physiological parameters of dairy cows 7(1982)201

Studies on fermentation of sorghum silage during storage, and its effect on milch animals 9(1983)143

Effect of intake on digestibility of plant cell walls and cell contents of complete diets for ruminants 9(1983)197

Feeding value of alkali-treated whole rye (*Secale cereale* L.) grain for lactating cows and its digestibility for sheep 10(1983/1984)77

The prediction of the net energy value of compound feedstuffs for dairy cattle 11(1984) 137

The effect of intake and hay: concentrate ratio upon digestibility and digesta passage 11 (1984)167

Digestibility and digesta retention in dairy cows receiving hay or silage at varying concentrate levels 11(1984)279

The use of barley straw as a feed 14(1986)29

Aqueous ammonia treatment of wheat straw: voluntary intake and digestibility in cattle 14(1986)81

Sugar beet pulp products for dairy cows 15 (1986)123

Concentrates for dairy cows: effects of feeding method, proportion in diet and type 15 (1986)167

Digestibility of dry matter, nitrogen and energy by dairy cows fed whole or hammer-milled lupin grain in oaten hay or oaten pasture based diets 16(1986/1987)143

Fermentation of hammermilled barley, lupin, pea and faba bean grain in the rumen of dairy cows 16(1986/1987)261

A review of the nutritive value and utilization of whole cottonseed, cottonseed meal and associated by-products by dairy cattle 18 (1987)89

A note on the use of sugar-beet pulp silage and molasses in the diet of lactating dairy cows 20(1988)13

The substitution of grass silage by a straw and concentrate mixture for dairy cows 20 (1988)135

The intake and digestion of maize silage-based diets by dairy cows and sheep 20 (1988)299

Feed intake and milk yield at increasing supplement of a palmitic and stearic acid-rich type of fat in comparison with animal fat 22(1988)179

Feed intake, milk yield and milk composition by replacing unprotected fat by Ca-soaps for dairy cows 22(1988)193

Efficacy of supplemental dietary neutralizing agents for lactating dairy cows. A review 23(1989)277

Response of high-producing dairy cows to high levels of dried corn gluten feed 24 (1989)191

Discriminating power of the nylon bag technique and the pepsin-cellulase method to detect less commonly used components included in concentrates for dairy cows 25 (1989)1

Rumen ingesta kinetics of cell wall components in dairy cows 25(1989)89

Cattle–general

The potato (*Solanum tuberosum*) as a source of nutrients for pigs, calves and fowl – a review 2(1977)171

The intake of steers and heifers fed on mixtures of alkali-treated or untreated straw with grass silage 2(1977)315

Differences between cattle and sheep in their digestion and relative intake of a mature tropical grass hay 3(1978)41

Estimation of the digestibility of low-quality hays by cattle from measurements made with sheep 3(1978)51

Raw and steam-pelleted cassava, sweet potato and yam *cayenensis* as starch sources for ruminant and chicken diets 3(1978)73

Digestibility by cattle of diets containing dried pig faeces 3(1978)155

Cellulase insoluble fibre as a measure of unavailable organic matter in cattle compounds containing alkali-treated straw 3 (1978)227

Digestibility of wet cage layer excreta and an excreta-supplemented diet by cattle as determined by two methods 5(1980)77

Nutritive value of ensiled broiler litter for cattle 8(1983)19

Comparative digestibility of feedstuffs by sheep and cows 12(1984/1985)47

Effects of molasses/urea and bacterial inoculant additives on silage quality, dry matter recovery, and feeding value for cattle 12 (1984/1985)205

The effect of nitrogen supplements to maize silage on the amount and composition of the nitrogen fraction entering the intestine of cattle 13(1985)25

Nitrogen balance and abomasal protein-nitrogen flow in growing ruminants fed methane digester effluent in combination with distillers dried grains 13(1985)47

Studies on the feeding value of agro-industrial by-products. III. Digestibility of cocoa-pod and cocoa-pod-based diets by ruminants 13 (1985)249

The energetic efficiency of rumen microbial protein synthesis in cattle given silage-based diets 13(1985)255

Hyperexcitability in cattle fed ammoniated roughages 17(1987)121

Nutritive value of three cultivars of triticale for sheep, pigs and poultry 18(1987)19

The effect of grinding and pelleting hay on digestibility, fermentation rate, digesta passage and rumen and faecal particle size in cows 19(1988)145

Rumen fermentation, urease activity and performance of cattle given palm kernel cake-based diet 20(1988)79

The effects of barley, unmolassed sugar-beet pulp and molasses supplements on organic matter, nitrogen and fibre digestion in the rumen of cattle given a silage diet 20(1988) 259

Colonization of guinea grass by anaerobic rumen fungi in swamp buffalo and cattle 22 (1988)161

Comparative fiber degradation by mixed rumen fungi from Australian and U.S.A. cattle 23(1989)305

Distribution of liquid supplement injected into large bales of roughage and comparison of nitrogen supplements for stocker cattle fed low-quality roughages 24(1989)13

Effect of urea treatment on rumen degradation characteristics of rice straws 24(1989) 83

Effect of soft white wheat addition to alfalfa-grass forage on heifer gain, diet digestibility and in vitro digestion kinetics 24(1989)141

Utilization of sugar beet molasses and monensin for finishing dairy bullocks 25(1989) 11

Intake, digestion and daily gain by cattle consuming bermudagrass (*Cynodon dactylon*) and supplemented with different combinations of ground corn, vegetable oil, urea and corn gluten and blood meals 25(1989)99

Cell content

Apparent and true digestibility of the Weende components, cell content and cell wall of ryegrass 1(1976)607

Cell wall

Apparent and true digestibility of the Weende components, cell content and cell wall of ryegrass 1(1976)607

Use of processing to explore the factors affecting the digestion of forage cell walls 6 (1981)387

Suitability of cell-wall constituents as predictors of organic matter digestibility in some tropical and subtropical by-products 7(1982)191

Effect of treatment of wheat straw with ammonia and peracetic acid on digestibility in vitro and cell wall composition 7(1982)325

Effect of intake on digestibility of plant cell walls and cell contents of complete diets for ruminants 9(1983)197

Electron microscopy of bacteria involved in the digestion of plant cell walls 10(1983/1984)93

The role of ciliate protozoa and fungi in the rumen digestion of plant cell walls 10(1983/1984)121

Lignin-polysaccharide complexes of the plant cell wall and their effect on microbial degradation in the rumen 21(1988)219

SO_2-treated straw as a silage additive: the participation of soluble and cell wall monosaccharide residues in lucerne silage fermentation 22(1988)255

Oven and stack ammoniation of grass hays. 1. Changes in chemical composition in relation to digestibility in vitro and cell-wall degradability 24(1989)299

Rumen ingesta kinetics of cell wall components in dairy cows 25(1989)89

Cellulase

Prediction of the digestibility of ruminant feeds from their solubility in enzyme solutions 2(1977)153

Enzyme solubility – a method for evaluating the digestibility of alkali-treated straw 2 (1977)205

Prediction of the dry matter digestibility of tropical grasses using a pepsin-cellulase assay 2(1977)247

Cellulase insoluble fibre as a measure of unavailable organic matter in cattle compounds containing alkali-treated straw 3 (1978)227

A note on Onozuka 3S cellulase as a replacement for Onozuka SS (P1500) cellulase when estimating forage digestibility in vitro 5(1980)347

Accuracy of predicting digestibility by the cellulase technique; the effect of pretreatment of forage samples with neutral detergent or acid pepsin 7(1982)83

Comparison of forage digestibility in vitro with enzymic solubility 11(1984)125

A comparison of laboratory methods involving single or two-stage cellulase and two-stage rumen inoculum–pepsin treatment for predicting digestibility and intake of hays in sheep 12(1984/1985)101

The use of an enzymatic technique to predict digestibility, metabolizable and net energy of compound feedstuffs for ruminants 14 (1986)203

The use of a cellulase technique to predict digestibility, metabolizable and net energy of forages 19(1988)247

Effect of sodium silicate on forage digestion with rumen fluid of goats or cellulase using culture solutions adjusted for pH 24(1989) 173

Cellulolysis

Manipulation of rumen fluid pH and its influence on cellulolysis in sacco, dry matter degradation and the rumen microflora of sheep offered either hay or concentrate 10 (1983/1984)1

Associative effects of mixed feeds. I. Effects of type and level of supplementation and the influence of the rumen fluid pH on cellulolysis in vivo and dry matter digestion of various roughages 10(1983/1984)15

Cellulose

The digestibility and nutritive value for dairy heifers of rations containing cellulose fibre 2(1977)287

Some remarks on the determination of acid-detergent fibre in feeds with cetyltrimethylammonium bromide versus determination of detergent fibre with alkylarylsulfonic acid 6(1981)91

The effects of sulphur supplementation on cellulose digestion in vitro and on nutrient digestion, nitrogen metabolism and rumen characteristics of lambs fed on good quality

fescue and tropical star grass hays 8(1983) 129

Effect of defaunation on the loss of energy in wethers fed different quantities of cellulose and normal or steamflaked maize starch 16 (1986/1987)233

In-reactor cellulose conversion 21(1988)99

The physiological effects of cellulose in the human large intestine 23(1989)241

Cephalosporium eichhorniae

Further thermotolerant fungi for the conversion of cassava starch to protein 2(1977)7

Protein quality studies on rats fed on cassava-root meal enriched with *Cephalosporium eichhorniae* 152 or with conventional plant-protein supplements 6(1981)197

Cereals, general

A comparison of bioavailable energy values of ground cereal grains measured with adult cockerels and growing pigs 19(1988)105

Chaetomium cellulolyticum

Utilization of cattle manure for single-cell protein production with *Chaetomium cellulolyticum* 4(1979)199

Nutritive value of single-cell protein produced by *Chaetomium cellulolyticum* grown on maize stover and pulp-mill sludge 11(1984) 211

Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. I. Composition and nutritive evaluation 22(1988)1

Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. II. Effect of method of drying and response to amino acid supplementation 22(1988)13

Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. III. Nutritive value for chicks and piglets 22(1988)23

Chaparral

Seasonal changes in composition and intake of chaparral by Spanish goats 8(1983)47

Chenopodium quinoa

The quality of the green fraction of leaf protein concentrate from *Chenopodium quinoa* Willd. grown at different levels of fertilizer nitrogen 11(1984)239

Chicken–broiler

Nutritive value of spruce muka (foliage) for the growing chick 3(1978)63

Utilization of processed pig hair in poultry diets 3(1978)243

Influence of steam pelleting on the feeding value of tower and candle rapeseed products for broiler chickens 3(1978)251

Trypsin inhibitor, mineral availability, and performance of broiler chickens fed on diets based on rice bran 4(1979)133

Feeding value of sunflower and groundnut cakes for broilers 4(1979)143

House fly larvae meal grown on municipal organic waste as a source of protein in poultry diets 4(1979)227

Distillery effluents as animal feed: the use of condensed beet molasses stillage (CBMS) in broiler feeding 5(1980)233

Nitrogen metabolism in broiler chickens consuming the bacterial strain *Alcaligenes eutrophus* 5(1980)241

Evaluation of brewer's dried grains in the diets of broiler chickens 6(1981)179

Influence of age on the nutrient utilization of diets for broilers 6(1981)405

Influence of increasing levels of β-glucanase on the productive value of barley diets for broiler chickens 7(1982)351

Nutritional value of waxy barleys for broiler chickens 8(1983)25

The influence of steam pelleting and grinding on the nutritive value of Canola rapeseed for poultry 8(1983)303

A note on the nutritive value of dried instant coffee residue for broiler chickens and turkey poults 10(1983/1984)285

Substitution of finger millet (*Eleusine coracana*) and bulrush millet (*Pennisetum typhoides*) for maize in broiler feeds 10(1983/1984)291

Meat and bone and fish meals in balancer feeds for choice-fed broilers 11(1984)221

The proximate composition and nutritive value of the winged bean *Psophocarpus tetragonolobus* (L.) DC for broilers 11 (1984)231

The utilization by broiler chickens of poultry offal hydrolysed in formic acid 11(1984)247

The effect of pelleting on the utilisation of yeast grown on n-alkanes by broiler chicks 12(1984/1985)285

Influence of supplemental fat and steam pelleting on the response of broilers to diets varying in energy density 13(1985)57

The effect of β-glucanase on the utilization of starch and nitrogen by broiler chickens fed on barley of low- or high-viscosity 15(1986) 83

Effect of dietary ascorbic acid (vitamin C) supplementation on the performance of broiler chickens in a tropical environment 16(1986/1987)17

Influence of nutrient density on feed consumption, weight gain and gut capacity of broilers, Leghorns and turkeys reared to 26 days of age 16(1986/1987)129

Palm kernel meal as a feed for poultry. 3. Replacement of groundnut cake by palm kernel meal in broiler diets 16(1986/1987) 195

Nutritive value of irradiated and β-glucanase-treated wild oat groats (*Avena fatua* L.) for broiler chickens 16(1986/1987)243

The effect of adding white fish meal containing enzyme digested or untreated stickwater solids to diets for broiler chickens 16 (1986/1987)253

Utilisation of L- and DL-amino acids by broiler chicks fed diets containing Pruteen 17(1987)1

Oil palm slurry (OPS) as a partial replacement for maize in the diets of broiler chickens 17(1987)157

The energy content of full-fat soya beans in meal and pelleted feeds for adult cocks and broilers 19(1988)97

Feeding and nutritive value of hominy and corn grits for poultry 19(1988)313

The metabolisable energy (AME_n) of fats for broilers in relation to their chemical composition 20(1988)45

Treatment of low-tannin sorghum grain for broiler feed 20(1988)69

Utilization of aquatic plants *Elodea canadensis* and *Hydrilla verticillata* in broiler chick diets 20(1988)155

Effects of enzyme supplementation of diets based on wheat, rye or triticale on their productive value for broiler chickens 20 (1988)313

The nutritional value for broilers of fats characterized by short-chain fatty acids as affected by level of inclusion and age 22(1988) 105

Nutritional value of triticale (Carman) for broiler diets 22(1988)119

Effects of wood ash extract treatment on the feeding value and utilization of high-tannin sorghums by broiler chicks 22(1988)131

The nutritional value of dried coffee pulp (DCP) in broiler chickens's diets 22(1988) 139

The effect of fat level and type on the utilization of triticale (cultivar Carman) by broiler chicks 22(1988)273

The nutritive value of tamarind seeds for broiler chicks 22(1988)285

Effects of oven-dried cassava peel meal on the performance and carcass characteristics of broiler chickens 24(1989)247

The effect of β-glucanase supplementation on nutrient digestibility and growth in broilers given diets containing barley, oat groats or wheat 25(1989)193

Chicken–chick

Amino acid supplementation of methanol-grown dried microbial cells in diets for young chicks 4(1979)63

Nutritive value of leaf-protein concentrate from potato haulm for rats and chicks 6 (1981)413

Growth performance of, and mimosine excretion by, young chicks fed on *Leucaena leucocephala* 7(1982)247

Metabolisable energy value of fats in chicks and adult cockerels 7(1982)365

Effects of milling and pelleting of maize, barley and wheat on their metabolizable energy value for cockerels and chicks 9(1983) 99

The effect of nutrient supplementation on the feeding value of kapok seed for poultry 9 (1983)263

A comparative study of metabolisable energy in ducklings and domestic chicks 11(1984) 199

A comparison of metabolizable energy content of eight common feed ingredients determined with young guinea fowls (keets) and pullet chicks 15(1986)1

Palm kernel meal as a feed for poultry. 1. Composition of palm kernel meal and availability of its amino acids to chicks 16(1986/ 1987)179

Palm kernel meal as a feed for poultry. 2. Diets containing palm kernel meal for starter and grower pullets 16(1986/1987)187

Utilization of tannin-containing and tannin-free faba beans (*Vicia faba*) by young chicks: effects of pelleting feeds on energy, protein and starch digestibility 20(1988)59

Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. III. Nutritive value for chicks and piglets 22(1988)23

Canavanine content and toxicity of sesbania leaf meal for growing chicks 25(1989)137

The effect of kernels of karanja (*Pongamia glabra* Vent) on growth and feed efficiency in broiler chicks to 4 weeks of age 25(1989) 201

Chicken–cockerel

Metabolisable energy value of fats in chicks and adult cockerels 7(1982)365

Comparison of bioavailable energy values of diets measured with poultry and swine 8 (1983)191

Effects of milling and pelleting of maize, barley and wheat on their metabolizable energy value for cockerels and chicks 9(1983) 99

Observations on the chemical and nutritive content of white winter and spring wheats contaminated with deoxynivalenol (vomitoxin) 11(1984)293

Feeding value of hydroponically sprouted barley for poultry and pigs 13(1985)183

A comparison of bioavailable energy values of ground cereal grains measured with adult cockerels and growing pigs 19(1988)105

Effect of enzyme supplementation on the nutritive value of hulless barley in chicken diets 24(1989)233

Chicken–general

The potato (*Solanum tuberosum*) as a source of nutrients for pigs, calves and fowl – a review 2(1977)171

A comparison of three fibre methods for predicting the metabolizable energy content of sorghum grain for poultry 2(1977)197

Estimation of lysine in compounded poultry diets after enzymic digestion 3(1978)57

Raw and steam-pelleted cassava, sweet potato and yam *cayenensis* as starch sources for ruminant and chicken diets 3(1978)73

Rapeseed meal and its use in poultry diets. A review 5(1980)255

Proximate analysis, amino acid composition and tannin content of improved Nigerian sorghum varieties and their potential in poultry feeds 7(1982)359

Chemical composition and nutritional value of para-rubber seed and its products for chickens 10(1983/1984)257

A new biological method for determining amino acid digestibility in poultry feedstuffs using a simple cannula, and the influence of dietary fibre on endogenous amino acid output 12(1984/1985)29

Nutritional value for chickens of dried distillers-spent-grain from barley and dehulled barley 17(1987)145

Nutritional value of pigeonpea (*Cajanus cajan*) meal in poultry diets 25(1989)123

Chicken–laying hen

Feeding value of sunflower- and groundnut-cakes for laying hens 6(1981)63

Effects of the addition of silica to the diet of the laying hen, with particular reference to egg-shell quality 6(1981)157

Evaluation of brewer's dried grains in the diets of laying hens 6(1981)169

Rapeseed meal tannins and egg taint 6(1981) 421

The utilization of torula yeast in diets for White Leghorn birds during growth and laying periods 7(1982)185

The involvement of trimethylamine oxide in fish meal in the production of egg taint 8 (1983)119

Influence of nutrient density on feed consumption, weight gain and gut capacity of broilers, Leghorns and turkeys reared to 26 days of age 16(1986/1987)129

The influence of oat groats and dietary level of lysine on the laying performance of cross-bred hens 17(1987)271

Evaluation of triticale given to pigs, poultry and rats 19(1988)359

Palm kernel meal as a feed for poultry. 4. Use of palm kernel meal by laying birds 20 (1988)279

Dry matter yield and chemical composition of pigeon pea (*C. cajan*) leaf meal and the nutritive value of pigeon pea leaf meal and grain meal for laying hens 24(1989)111

Chopping

Influence of processing methods and storage time on the cyanide potential of cassava leaf meal 17(1987)227

Chunies

Nutritional evaluation of gram (*Cicer arietinum*) and arhar (*Cajanus cajan*) chunies for cattle 5(1980)315

Citrus pulp

Composition of citrus pulp 5(1980)1

Citrus pulp in diets for fattening lambs 5 (1980)11

Citrus pulp in diets for fattening rabbits 5 (1980)23

Clover

Losses of beta-carotene in red clover in an acid medium during ensiling 4(1979)81

The enzymic nature of the degradation of beta-carotene in red clover and in other forage crops during silage-making with acid additives 5(1980)59

Degradation in vitro and in sacco of red clover leaves and stems 17(1987)295

Cobalt
Treatment of dairy sheep with soluble glass boluses containing copper, cobalt and selenium 19(1988)79

The effect of soluble-glass boluses containing copper, cobalt and selenium on the blood composition of ewes 21(1988)23

Cocoa-pod
Studies on the feeding value of agro-industrial by-products: strategies for improving the utilisation of cocoa-pod-based diets by ruminants 20(1988)189

Coffee pulp
Changes in nutrient intake and performance by goats fed coffee pulp-based diets followed by a commercial concentrate 14 (1986)1

The nutritional value of dried coffee pulp (DCP) in broiler chickens's diets 22(1988) 139

Coffee residue
A note on the nutritive value of dried instant coffee residue for broiler chickens and turkey poults 10(1983/1984)285

Colour value
Influence of pericarp and endosperm colour and type on digestibility of grain sorghum by pigs 2(1977)219

Relationships between tristimulus colour values and digestibility of grain sorghum by swine 2(1977)327

Compound feeds
Cellulase insoluble fibre as a measure of unavailable organic matter in cattle compounds containing alkali-treated straw 3 (1978)227

Some factors affecting the future composition of UK compound animal feeds 6(1981)1

Energy evaluation of feeds and compounded diets for pigs – a review 7(1982)387

The prediction of the net energy value of compound feedstuffs for dairy cattle 11(1984) 137

The use of an enzymatic technique to predict digestibility, metabolizable and net energy of compound feedstuffs for ruminants 14 (1986)203

The prediction of the energy value of compounded pig foods from chemical analysis 17(1987)81

Cooking
Effect of sodium hydroxide treatment and/or extrusion cooking on the nutritive value of peanut hulls 12(1984/1985)187

Copper
Reducing the potential copper toxity of concentrates to sheep by the use of molybdenum and sulphur supplements 2(1977)235

Utilization by sheep of dried pig faeces containing a high concentration of copper 3 (1978)31

Olaquindox as performance-promoting feed additive for growing pigs 4(1979)117

Treatment of dairy sheep with soluble glass boluses containing copper, cobalt and selenium 19(1988)79

The effect of age of regrowth on content and release of manganese, iron, zinc and copper from four tropical forages incubated in sacco in rumen of sheep 20(1988)171

The effect of soluble-glass boluses containing copper, cobalt and selenium on the blood composition of ewes 21(1988)23

Copper and iodine in pig diets with high glucosinolate rapeseed meal. I. Performance and thyroid hormone status of growing pigs fed on a diet with rapeseed meal treated with copper sulphate solution or untreated and supplements of iodine, copper or a quinoxaline derivative 22(1988)33

Copper and iodine in pig diets with high glucosinolate rapeseed meal. II. Influence of iodine supplements for rations with rapeseed meal untreated or treated with copper ions on performance and thyroid hormone status of growing pigs 22(1988)45

Corn gluten feed
Evaluation of corn gluten meal with urea as a source of supplementary nitrogen for growing calves and lambs 7(1982)375

Response of high-producing dairy cows to high levels of dried corn gluten feed 24 (1989)191

Cotton
Effect of ammonia treatment and processing of whole cotton plants as sole source of roughage in complete feeds for growing cross-bred calves 13(1985)93

Cotton straw
Effects of ammonia treatment and processing of cotton straw as the sole source of roughage in complete feeds on rumen characteristics with Murrah buffaloes 14(1986) 95

Cottonseed
A review of the nutritive value and utilization of whole cottonseed, cottonseed meal and associated by-products by dairy cattle 18 (1987)89

Cottonseed cake
Comparison of cottonseed cake and dried blood as nitrogen sources for growing Sudan Nubian goats 22(1988)333

Cottonseed meal
Effects of supplementing a wheat straw diet with urea or a urea–molasses block and/or cottonseed meal on intake and liveweight change of lambs 16(1986/1987)25

Nutrient composition of Greek cottonseed meal 18(1987)295

Evaluation of Greek cottonseed meal. 2. Use in rations for fattening pigs 18(1987)303

Chemical and biological evaluation of Nigerian cottonseed meal protein quality 20 (1988)251

Cowpea
Variation in metabolizable energy content of raw or autoclaved white and brown varieties of three tropical grain legumes 13 (1985)141

Crop residues
Alkaline hydrogen peroxide treatments of crop residues to increase ruminal dry matter disappearance in sacco 17(1987)179

Culinary wastes
Chemical composition of culinary wastes and their potential as a feed for ruminants 5 (1980)205

Cyanide
The influence of processing and storage on hydrogen cyanide and tannin contents of para-rubber seed and its products 9(1983) 319

Effect of cassava-based diets varying in cyanide content on the performance and physiopathology of the African giant rat (*Cricetomys gambianus* Waterhouse) 11 (1984)1

Effect of variety and plant age on the cyanide content of whole-root cassava chips and its reduction by sun-drying 11(1984)57

Influence of processing methods and storage time on the cyanide potential of cassava leaf meal 17(1987)227

Cystine
Meat meal in the diet of the early-weaned pig. II. Amino acid supplementation 4(1979)43

D

Defaunation
Effect of defaunation on straw digestion in sacco and on particle retention in the rumen 10(1983/1984)165

Effect of defaunation on the loss of energy in wethers fed different quantities of cellulose and normal or steamflaked maize starch 16 (1986/1987)233

Effect of defaunating the rumen 21(1988)229

Deoxynivalenol
Observations on the chemical and nutritive content of white winter and spring wheats contaminated with deoxynivalenol (vomitoxin) 11(1984)293

Deterioration
A note on the effect on the aerobic deterioration of maize silage of the manipulation of the microflora by means of antibiotics 3 (1978)89

Losses of beta-carotene in red clover in an acid medium during ensiling 4(1979)81

Chemicals for preventing moulding in damp hay 6(1981)323

Losses of beta-carotene in unwilted forage crops during silage-making and feeding 9 (1983)63

A comparison between degradation in vitro and in sacco of constituents of untreated and ammonia-treated barley straw 10(1983/ 1984)199

The effect of anhydrous ammonia and moisture content on the preservation and chemical composition of perennial ryegrass hay 11(1984)159

Effect of dietary nitrogen source on microbial protein synthesis, dietary protein degradation and nutrient digestion in steers 14 (1986)117

Nutrient losses during deterioration of hay in relation to changes in biochemical composition and microbial growth 15(1986)149

Estimating yield and nutrient losses due to rainfall on field-drying alfalfa hay 16(1986/ 1987)7

Effect of formaldehyde treatment on the degradation of acid-preserved fish silage protein in vitro 16(1986/1987)297

Whole-crop peas. I. Changes in botanical and chemical composition and rumen in vitro degradability during maturation 17(1987) 15

Degradation in vitro and in sacco of red clover leaves and stems 17(1987)295

Microbial dynamics in barley grain stored under controlled atmosphere 19(1988)299

Fungal contamination of hydroponic forage 20 (1988)163

Deteroriation
The enzymic nature of the degradation of beta-carotene in red clover and in other forage crops during silage-making with acid additives 5(1980)59

Dicoumarol
Dicoumarol concentrations in sweetclover hay treated with preservatives and in spoiled hay of high- and low-coumarin cultivars of sweetclover 14(1986)221

Digesta particle size
The determination of digesta partical size in some herbivores 7(1982)35

Digesta retention
Digestibility and digesta retention in dairy cows receiving hay or silage at varying concentrate levels 11(1984)279

Digestibility–*Agropyron elongatum*
Effect of lactate on the in vitro digestion of *Agropyron elongatum* by rumen microorganisms 16(1986/1987)161

Digestibility–alfalfa
Composition and fibre digestion in morphological components of an alfalfa–timothy sward 19(1988)135

Digestibility–amino acids and proteins
Apparent ileal and faecal digestibility of amino acids in barley-based diets supplemented with soya bean meal or canola meal for growing pigs 14(1986)183

Digestibility–animal waste
Digestibility by cattle of diets containing dried pig faeces 3(1978)155

Digestibility of wet cage layer excreta and an excreta-supplemented diet by cattle as determined by two methods 5(1980)77

Digestibility and beef performance after feeding NaOH- or KOH-treated pig wastes 17 (1987)235

Digestibility–barley
The effect of calcium and sodium hydroxides and of sodium acrylate on the fermentation and digestibility in vitro of ensiled whole-crop wheat and barley harvested at different stages of maturity 9(1983)37

Effect of storage and processing of barley on its utilization by pigs and rats 9(1983)89

The digestion by sheep of barley and maize-based diets containing either meat and bone meal or soya bean meal 15(1986)301

Digestibility–barley meal
The influence of feeding level and level of dietary inclusion on the digestibility of barley meal in the pig 2(1977)41

Digestibility–bermudagrass
Intake, digestion and daily gain by cattle consuming bermudagrass (*Cynodon dactylon*) and supplemented with different combinations of ground corn, vegetable oil, urea and corn gluten and blood meals 25(1989)99

Digestibility–canola meal
Factors effecting the digestibility by pigs of energy and protein in wheat, barley and sorghum diets supplemented with canola meal 24(1989)253

Digestibility–cassava
Net energy for gain (NE_{gain}) of cassava tops 20(1988)97

Digestibility–cell contents
Effect of intake on digestibility of plant cell walls and cell contents of complete diets for ruminants 9(1983)197

Digestibility–cell wall
Use of processing to explore the factors affecting the digestion of forage cell walls 6 (1981)387

Effect of intake on digestibility of plant cell walls and cell contents of complete diets for ruminants 9(1983)197

Digestibility–compound feeds
The use of an enzymatic technique to predict digestibility, metabolizable and net energy of compound feedstuffs for ruminants 14 (1986)203

Digestibility–crop residues
Improving utilization of crop residues. Evaluation of two chemicals used to improve digestibility of crop residues 9(1983)205

Digestibility–desert grass
Effect of sodium hydroxide treatment of desert grass on carbohydrate digestibility by goats 17(1987)219

Digestibility–dry matter
The accuracy of the pepsin–cellulase technique for estimating the dry matter digestibility in vivo of grasses and legumes 3(1978)277

Bias when predicting crude protein, dry matter digestibility and voluntary intake of tropical grasses by near-infrared reflectance 9(1983)221

Variations and prevision of the in sacco dry matter digestion of concentrates and by-products 13(1985)7

Digestibility–faba bean
Utilization of tannin-containing and tannin-free faba beans (*Vicia faba*) by young chicks: effects of pelleting feeds on energy, protein and starch digestibility 20(1988)59

Digestibility–fibre
The effects of different sources of nitrogen supplementation on the digestion of fibre components in the rumen of steers 17(1987) 65

Approach to the energetic importance of fibre digestion in pigs. I. Importance of fermentation in the overall energy supply 23(1989) 141

Approach to the energetic importance of fibre digestion in pigs. II. An experimental approach to hindgut digestion 23(1989)169

Digestion and fermentation pattern of various dietary fiber sources in the rat 23(1989)195

Digestibility–fish meal
Fish protein concentrates for pre-ruminant calves: effects of processing temperatures and source of fish on protein digestibility and biological value 18(1987)181

Digestibility–fodder beet
The chemical composition, digestibility and energy value of fodder-beet roots 18(1987) 225

Digestibility–forage, general
Some remarks on the analytical procedure of Van Soest for the prediction of forage digestibility 3(1978)309

Variability in structural carbohydrates and in vitro digestibility of forages. 4. Fenugreek (*Trigonella foenum-graecum*) 4(1979)17

A note on Onozuka 3S cellulase as a replacement for Onozuka SS (P1500) cellulase when estimating forage digestibility in vitro 5(1980)347

Effect of high-pressure steam treatment on chemical composition and digestibility in vitro of roughages 7(1982)61

Accuracy of predicting digestibility by the cellulase technique; the effect of pretreatment of forage samples with neutral detergent or acid pepsin 7(1982)83

Relative intakes, digestibility and bacterial protein synthesis by sheep and cattle fed high-roughage diets 16(1986/1987)75

Prediction of forage digestibility by principal component analysis of near infrared reflectance spectra 16(1986/1987)215

Digestibility–forages, general
Comparison of laboratory methods for predicting the organic matter digestibility of forages 2(1977)337

Digestibility–general
Prediction of the digestibility of ruminant feeds from their solubility in enzyme solutions 2(1977)153

Comparative digestibility of feedstuffs by sheep and cows 12(1984/1985)47

Digestibility–grapevine
Digestibility in vivo of ensiled grapevines (branches and leaves): influence of the system of analysis in the detergent fibre scheme on the prediction of digestibility 22 (1988)173

Digestibility–grass
Apparent and true digestibility of the Weende components, cell content and cell wall of ryegrass 1(1976)607

Prediction of the dry matter digestibility of tropical grasses using a pepsin-cellulase assay 2(1977)247

Comparison of forage digestibility in vitro with enzymic solubility 11(1984)125

Effect of sodium silicate on forage digestion with rumen fluid of goats or cellulase using culture solutions adjusted for pH 24(1989) 173

Digestibility–groundnut
The effect of grinding groundnut hulls either with or without alkali treatment on digestibility of diet and on ruminal and blood components 15(1986)69

Digestibility–hay
Apparent and true digestibility of the Weende components, cell content and cell wall of ryegrass 1(1976)607

Differences between cattle and sheep in their digestion and relative intake of a mature tropical grass hay 3(1978)41

Estimation of the digestibility of low-quality hays by cattle from measurements made with sheep 3(1978)51

The effect of intake and hay: concentrate ratio upon digestibility and digesta passage 11 (1984)167

Digestibility and digesta retention in dairy cows receiving hay or silage at varying concentrate levels 11(1984)279

The voluntary intake and digestibility of combinations of cereal crop residues and legume hay for sheep 12(1984/1985)241

Chemical composition and digestion in vitro of orchardgrass hay ammoniated by different techniques 17(1987)109

The effect of grinding and pelleting hay on digestibility, fermentation rate, digesta passage and rumen and faecal particle size in cows 19(1988)145

Oven and stack ammoniation of grass hays. 1. Changes in chemical composition in relation to digestibility in vitro and cell-wall degradability 24(1989)299

Oven and stack ammoniation of grass hays. 2. Voluntary intake, digestibility and faecal nitrogen excretion in sheep in relation to hay composition 24(1989)313

Effects of urea treatment on chemical composition and in vitro digestibility on meadow hays of Northern Portugal 25(1989)157

Digestibility–jackbean
Ripe beans of *Canavalia ensiformis* (jackbean) as feed ingredient for monogastric animals 20(1988)325

Digestibility–legumes
Comparison of forage digestibility in vitro with enzymic solubility 11(1984)125

Digestibility–*Leucaena leucocephala*
The intake and digestibility in goats of *Leucaena leucocephala* supplemented with dehydrated palm oil mill effluent 24(1989)45

Digestibility–lupin
Digestibility and energy value of sweet lupin seed (*Lupinus albus* var. multolupa) in pigs 12(1984/1985)171

Digestibility of dry matter, nitrogen and energy by dairy cows fed whole or hammer-milled lupin grain in oaten hay or oaten pasture based diets 16(1986/1987)143

Digestibility–maize
Rate of growth, feed utilisation and digestibility observed with diets containing ensiled maize grain or maize ears fed alone or in mixtures 1(1976)401

The digestion and metabolism of the energy and protein of maize 1(1976)455

Measurement and prediction of digestibility of forage maize in The Netherlands 10 (1983/1984)301

Effect of sodium acrylate and calcium and sodium hydroxides on the fermentation, digestibility, chemical composition and aerobic deterioration of whole-crop maize silage 12(1984/1985)91

High temperature and pressure processing of maize cobs. I. Digestibility in vitro of processed cobs 12(1984/1985)151

The intake and digestibility by sheep of oat straw and maize stover offered with different levels of noug (*Guizotia abyssinica*) meal 16(1986/1987)99

Digestibility–manioc
The digestion of diets containing manioc by young growing pigs 12(1984/1985)119

Digestibility–mixed feeds
Associative effects of mixed feeds. I. Effects of type and level of supplementation and the influence of the rumen fluid pH on cellulolysis in vivo and dry matter digestion of various roughages 10(1983/1984)15

Associative effects of mixed feeds. II. The effect of dietary addition of bicarbonate salts on the voluntary intake and digestibility of diets containing various proportions of hay and barley 10(1983/1984)31

Digestibility–molasse
Differences in chemical composition and digestibility of beet and cane molasses 13 (1985)83

Digestibility–nitrogen
The apparent and true ileal digestibility of nitrogen and amino acids in lactic casein for the growing pig 16(1986/1987)169

Digestibility–olive cake
Effects of feeding alkali-treated olive cake on intake, digestibility and rumen liquor parameters 14(1986)139

Digestibility–organic matter
Suitability of cell-wall constituents as predictors of organic matter digestibility in some tropical and subtropical by-products 7(1982)191

The effect of high levels of molasses in combinations with hay on digestibility of organic matter, microbial protein synthesis and volatile fatty acid production in vitro 7(1982)277

A comparison between fermentation heat of forages and organic matter digestibility determined by in vitro incubation with rumen fluid 23(1989)333

Digestibility–paper diets
Intake, digestibility and energy utilization of corrugated paper diets 17(1987)57

Digestibility–pasture plants
The relationship between the intake of herbage grazed by dairy cows and its digestibility 1 (1976)53

The analytical and biological accuracy of estimating the dry matter digestibility of different legume species 1(1976)61

Solubility of the protein of tropical pasture species and the rate of its digestion in the rumen 5(1980)183

Digestibility–pea
Whole-crop peas. II. Digestion of early- and late-harvested crops in the gastrointestinal tract of pigs 17(1987)33

Digestibility–plantain sheath
In vitro studies and short-term feeding trial in lambs to evaluate plantain sheath (*Musa sapientum*) as a feed for ruminants 20 (1988)343

Digestibility–potato
The effect of different patterns of thermal processing of potatoes on their digestibility by growing pigs 4(1979)295

Digestibility–prediction
Comparison of methods for predicting digestibility of feeds 20(1988)203

Digestibility–protected fat
The effects on feed intake, growth and digestibility of nutrients of including calcium soaps of fat in diets for young calves 14 (1986)103

Digestibility–proteins and amino acids
Digestibility of protein and amino acids of a fermentation single-cell protein for veal calves 2(1977)377

The apparent and true ileal digestibility of nitrogen and amino acids in lactic casein for the growing pig 16(1986/1987)169

Digestibility–rapeseed
Degradation of crude protein and quality of undegradable protein in untreated or formaldehyde-treated rapeseed meal 12 (1984/1985)19

Effect of graded levels of raw and processed rapeseed on feed digestibility and nutrient utilization in young pigs 24(1989)275

Digestibility–rapeseed meal
The digestion in the small intestine of young bulls of the protein of rapeseed meal treated or untreated with formaldehyde 7 (1982)225

Digestibility–ryegrass
The effect of urea on the preservation and digestibility in vitro of perennial ryegrass 10(1983/1984)49

Digestibility–silage
Digestibility and digesta retention in dairy cows receiving hay or silage at varying concentrate levels 11(1984)279

The effect of supplementing grass silage with sodium hydroxide-treated chopped or milled straw on intake and digestibility by wether sheep 12(1984/1985)215

The intake and digestion of maize silage-based diets by dairy cows and sheep 20 (1988)299

The digestibility and metabolisable energy content of grass silage and their prediction from laboratory measurements 24(1989)27

Digestibility–sisal waste
Feed value of Tanzanian sisal wastes: chemical composition and digestibility in vitro 14 (1986)255

Digestibility–sorghum grain
Influence of pericarp and endosperm colour and type on digestibility of grain sorghum by pigs 2(1977)219

Relationships between tristimulus colour values and digestibility of grain sorghum by swine 2(1977)327

Digestibility–soya bean
Effects of micronization on the digestibility of whole soya beans and rapeseeds for the growing pig 3(1978)179

Effect of formaldehyde-treated soya bean meal and urea in starters on nitrogen quality, degradibility in sacco, sheep digestibility and calf performance 16(1986/1987)287

Digestibility–straw
The effect on digestibility of a new technique for alkali treatment of straw 1(1976)73

Effect on digestibility and nitrogen content of barley straw of different ammonia treatments 2(1977)131

Enzyme solubility – a method for evaluating the digestibility of alkali-treated straw 2 (1977)205

Ensiled alkali-treated straw. I. Effect of level and type of alkali on the composition and digestibility in vitro of ensiled barley straw 3(1978)117

Effect of time of storage and method of drying on the digestibility in vitro of alkali-treated barley straw 3(1978)381

Titratable groups and soluble phenolic compounds as indicators of the digestibility of chemically treated roughages 6(1981)123

Studies on ammonia-treated straw. I. The effects of type and level of ammonia, moisture content and treatment time on the digestibility in vitro and enzyme soluble organic matter of oat straw 7(1982)45

Botanical composition and extent of lignification affecting digestibility of wheat and oat straw and paspalum hay 7(1982)71

The effect of calcium hydroxide treatment on the digestibility of chopped oat straw supplemented with a solution containing urea, calcium, phosphorus, sodium, trace elements and vitamins 7(1982)93

Intake and digestibility in sheep fed on alkali-treated straw and concentrates, as complete diets or separate feeds 7(1982)271

Effect of treatment of wheat straw with ammonia and peracetic acid on digestibility in vitro and cell wall composition 7(1982)325

Digestibility of diets containing increasing levels of NaOH-treated or untreated wheat straw 7(1982)331

The effect of prolonged storage on the digestibility and nitrogen content of ammonia-treated barley straw 8(1983)147

Composition and enzymatic digestibility of oregon grass straws 9(1983)283

Digestion of organic matter and nitrogen in ammoniated barley straw 10(1983/1984)65

Effect of defaunation on straw digestion in sacco and on particle retention in the rumen 10(1983/1984)165

Digestibility studies on ammonia-treated straw 10(1983/1984)213

The digestion of untreated and ammonia-treated barley straw in an artificial rumen 10(1983/1984)229

Ammonia treatment of straw via the hydrolysis of urea. II. Additions of soya bean (urease), sodium hydroxide and molasses; effects on the digestibility of urea-treated straw 11(1984)115

The effect of milling and sodium hydroxide treatment on intake and digestibility of wheat straw by sheep 11(1984)149

The utilization of untreated and calcium hydroxide treated wheat straw by sheep 12 (1984/1985)141

Aqueous ammonia treatment of wheat straw: voluntary intake and digestibility in cattle 14(1986)81

The intake and digestibility by sheep of oat straw and maize stover offered with different levels of noug (*Guizotia abyssinica*) meal 16(1986/1987)99

The effect of dietary barley on carbohydrate digestibility of sulphur dioxide-treated wheat straw by sheep 18(1987)55

Digestibility–sugar beet

Dried suger beet pulp as a high energy feed for beef cattle 1(1976)643

Digestion of diets containing molassed or plain sugar-beet pulp by growing pigs 23 (1989)67

Digestibility–tallow

Digestibility of a fully hydrogenated tallow for growing pigs 5(1980)87

Digestibility–Tilley and Terry test

Improving the reliability of the two-stage in vitro assay for ruminant feed digestibility by calibration against in vivo data from a wide range of sources 18(1987)233

Digestibility–timothy

Composition and fibre digestion in morphological components of an alfalfa–timothy sward 19(1988)135

Digestibility–wheat

The effect of calcium and sodium hydroxides and of sodium acrylate on the fermentation and digestibility in vitro of ensiled whole-crop wheat and barley harvested at different stages of maturity 9(1983)37

Effect of soft white wheat addition to alfalfa-grass forage on heifer gain, diet digestibility and in vitro digestion kinetics 24(1989)141

Drechslera sorokiniana

Effects of *Drechslera sorokiniana* infection on the yield and quality of tall fescue forage 15(1986)41

Dried bakery product

Variation in nutrient content of samples of dried bakery product 7(1982)419

Dried microbial cells

The replacement of skim milk by dried microbial cells and whey in milk diets for calves 2(1977)85

Limiting amino acids in dried microbial cells given to young turkeys 4(1979)209

Drying methods

Effect of time of storage and method of drying on the digestibility in vitro of alkali-treated barley straw 3(1978)381

Influence of the drying process on the composition of brewer's dried grains 6(1981)163

Effect of variety and plant age on the cyanide content of whole-root cassava chips and its reduction by sun-drying 11(1984)57

Microwave drying of alfalfa compared to field- and oven-drying: effects on forage quality 16(1986/1987)1

Estimating yield and nutrient losses due to rainfall on field-drying alfalfa hay 16(1986/1987)7

Influence of processing methods and storage time on the cyanide potential of cassava leaf meal 17(1987)227

The effect of treatment with urea, sorbic acid, or dehydration on orange peel silage 20 (1988)335

Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. II. Effect of method of drying and response to amino acid supplementation 22(1988)13

Drying treatment drastically affects feed evaluation and feed quality with certain tropical forage species 22(1988)69

Duck
A comparative study of metabolisable energy in ducklings and domestic chicks 11(1984) 199

Dye absorption test
The dye absorption test as an indicator of the nutritive quality of heat-processed navy bean 9(1983)71

E

Economics
Economic aspects of high-plane feeding of bulls with maize silage 1(1976)513

Energetics of stockfeed ~~oduction 3(1978)1

Assessment of the production costs of fish protein hydrolysates 7(1982)147

Editorial–general
The dilemma of animal feeds and nutrition 1 (1976)1

Egg taint
Rapeseed meal tannins and egg taint 6(1981) 421

The involvement of trimethylamine oxide in fish meal in the production of egg taint 8 (1983)119

Elodea
Utilization of aquatic plants *Elodea canadensis* and *Hydrilla verticillata* in broiler chick diets 20(1988)155

Energy–metabolizable
A comparative study of metabolisable energy in ducklings and domestic chicks 11(1984) 199

Effect of prolonged wilting in poor conditions on the fermentation quality, metabolisability and net energy value of silage given to sheep 12(1984/1985)109

Variation in metabolizable energy content of raw or autoclaved white and brown varieties of three tropical grain legumes 13 (1985)141

The use of an enzymatic technique to predict digestibility, metabolizable and net energy of compound feedstuffs for ruminants 14 (1986)203

A comparison of metabolizable energy content of eight common feed ingredients determined with young guinea fowls (keets) and pullet chicks 15(1986)1

The metabolisable energy (AME_n) of fats for broilers in relation to their chemical composition 20(1988)45

The digestibility and metabolisable energy content of grass silage and their prediction from laboratory measurements 24(1989)27

Energy–net
The use of an enzymatic technique to predict digestibility, metabolizable and net energy of compound feedstuffs for ruminants 14 (1986)203

The use of a cellulase technique to predict digestibility, metabolizable and net energy of forages 19(1988)247

Net energy for gain (NE_{gain}) of cassava tops 20(1988)97

Energy balance
Energy and nitrogen balance, intake and growth rate of lambs fed on unwilted grass silages treated with a formaldehyde–acid mixture or untreated 9(1983)107

Dietary fiber and energy balance: integration of the human ileostomy and in vitro fermentation models 23(1989)261

Energy density
Influence of supplemental fat and steam pelleting on the response of broilers to diets varying in energy density 13(1985)57

Energy intake
Dry matter, protein, energy and fibre intake by dairy heifers grazing a Rhodes grass (*Chloris gayana*) pasture 6(1981)15

Energy metabolism
Energy metabolism in lambs fed silage diets with sucrose or a growth promoter 16 (1986/1987)203

Effect of defaunation on the loss of energy in wethers fed different quantities of cellulose and normal or steamflaked maize starch 16 (1986/1987)233

Energy systems
The energy systems used in the EEC countries: their application to 24 commercial pig-feeds and relationships between them 6(1981)211

A flexible, two-component feeding system derived from Blaxter's three-component, metabolisable-energy system 6(1981)223

Prediction of the energy value of commercial pig feeds, according to the various energy systems used in the European Community, that are based on simple analytical parameters of the feed 6(1981)367

Energy utilisation
Protein and energy utilisation in rats of diets based on lupin seed (*Lupinus albus* var. multolupa) 15(1986)33

Intake, digestibility and energy utilization of corrugated paper diets 17(1987)57

Energy value
The digestion and metabolism of the energy and protein of maize 1(1976)455

Dried suger beet pulp as a high energy feed for beef cattle 1(1976)643

A comparison of three fibre methods for predicting the metabolizable energy content of sorghum grain for poultry 2(1977)197

Waxy versus regular maize:energy value for growing pigs and utilization in piglet diets 4(1979)109

Protein and energy value of vinasse for pigs 5 (1980)221

Predicting the energetic value of forage from digestibility of organic matter in vitro or solubility in potassium hydroxide 6(1981) 235

Metabolisable energy value of fats in chicks and adult cockerels 7(1982)365

Energy evaluation of feeds and compounded diets for pigs – a review 7(1982)387

Comparison of bioavailable energy values of diets measured with poultry and swine 8 (1983)191

Effects of milling and pelleting of maize, barley and wheat on their metabolizable energy value for cockerels and chicks 9(1983) 99

The prediction of the net energy value of compound feedstuffs for dairy cattle 11(1984) 137

Digestibility and energy value of sweet lupin seed (*Lupinus albus* var. multolupa) in pigs 12(1984/1985)171

Utilization of whole rape (Canola) seed and sunflower seeds as sources of energy and protein in calf starter diets 15(1986)101

The prediction of the energy value of compounded pig foods from chemical analysis 17(1987)81

Comparative bioavailable energy values of five ground cereal grains measured with growing rats and pigs 18(1987)143

The chemical composition, digestibility and energy value of fodder-beet roots 18(1987) 225

The energy content of full-fat soya beans in meal and pelleted feeds for adult cocks and broilers 19(1988)97

A comparison of bioavailable energy values of ground cereal grains measured with adult cockerels and growing pigs 19(1988)105

The effect of ammoniation on the nutritive value of wheat, barley and oat straws. II. Digestibility and energy value measurements in vivo and their prediction from laboratory measurements 19(1988)173

Approach to the energetic importance of fibre digestion in pigs. I. Importance of fermentation in the overall energy supply 23(1989) 141

Approach to the energetic importance of fibre digestion in pigs. II. An experimental approach to hindgut digestion 23(1989)169

Enzymes, general
The enzymic nature of the degradation of beta-carotene in red clover and in other forage crops during silage-making with acid additives 5(1980)59

Use of enzymes for improvement of feed 6 (1981)105

Composition and enzymatic digestibility of oregon grass straws 9(1983)283

The effect of enzymatic treatment on amino acid content and nitrogen characteristics of feather meal 16(1986/1987)151

Factors affecting the formation of polysaccharide-degrading enzymes by rumen microorganisms 21(1988)191

Effect of enzyme supplementation on the nutritive value of hulless barley in chicken diets 24(1989)233

Ethanol
Production of ethanol from lignocellulosic materials 21(1988)175

Ethanolamine
Effects of ethanolamine and ethylenediamine treatments on the composition and degradability of wheat straw 13(1985)131

Ethylenediamine
Effects of ethanolamine and ethylenediamine treatments on the composition and degradability of wheat straw 13(1985)131

F

Faba bean
Fermentation of hammermilled barley, lupin, pea and faba bean grain in the rumen of dairy cows 16(1986/1987)261

Utilization of tannin-containing and tannin-free faba beans (*Vicia faba*) by young chicks: effects of pelleting feeds on energy, protein and starch digestibility 20(1988)59

Faecal particle size
The effect of grinding and pelleting hay on digestibility, fermentation rate, digesta passage and rumen and faecal particle size in cows 19(1988)145

Fats
Poly-unsaturated fat in young ruminants 2 (1977)93

Digestibility of a fully hydrogenated tallow for growing pigs 5(1980)87

Metabolisable energy value of fats in chicks and adult cockerels 7(1982)365

Variability for protein and oil quality in *Lupinus albus* 9(1983)271

The effect of level and source of protein, fibre and fat in the diet on the energy value of compounded pig feeds 11(1984)11

Effects of pelleting, added sodium bentonite and fat in a wheat-based diet on performance and carcass characteristics of small white turkeys 12(1984/1985)223

Effect of processing on the utilization by young pigs of the fat in soya beans and sunflower seeds 12(1984/1985)267

Influence of supplemental fat and steam pelleting on the response of broilers to diets varying in energy density 13(1985)57

Performance and carcass characteristics of Sudan desert sheep and goats on high roughage diets with added fat 13(1985)147

Effect of dietary protein and fat levels on the utilization of the fat in sunflower seeds by the young pig 13(1985)159

Thermofractographic determination of monocarbonyl compounds in animal waste fats used as feed fat 13(1985)227

Fat supplementation of diets for growing rabbits 16(1986/1987)109

Influence of nutrient density on feed consumption, weight gain and gut capacity of broilers, Leghorns and turkeys reared to 26 days of age 16(1986/1987)129

High-fat maize in diets for pigs and sows 17 (1987)201

Isoacids in ruminant nutrition: their role in ruminal and intermediary metabolism and possible influences on performances – a review 18(1987)169

Investigations on the nutritional value for ruminants of various fats 19(1988)111

Use of fat in diets of sheep in hot environments. I. Effects on performance, carcass characteristics and lipid composition of plasma 19(1988)327

Use of fat in diets of sheep in hot environments. II. Effect on rumen metabolism 19 (1988)343

The metabolisable energy (AME_n) of fats for broilers in relation to their chemical composition 20(1988)45

Feed intake and milk yield at increasing supplement of a palmitic and stearic acid-rich type of fat in comparison with animal fat 22(1988)179

Feed intake, milk yield and milk composition by replacing unprotected fat by Ca-soaps for dairy cows 22(1988)193

The effect of fat level and type on the utilization of triticale (cultivar Carman) by broiler chicks 22(1988)273

Fatty acids
Absorption of short-chain fatty acids and their role in the hindgut of monogastric animals 23(1989)43

Feather meal
Protein quality of mixtures of animal protein meals 9(1983)79

Effects of processing time and moisture content on amino acid composition and nitrogen characteristics of feather meal 14(1986) 279

The effect of enzymatic treatment on amino acid content and nitrogen characteristics of feather meal 16(1986/1987)151

Feeding standards
Who needs feeding standards? 6(1981)101

Feeding standards are necessary 7(1982)423

Fenugreek
Variability in structural carbohydrates and in vitro digestibility of forages. 4. Fenugreek (*Trigonella foenum-graecum*) 4(1979)17

Fermentation–animal waste
Fermentation characteristics and feeding value of ensiled poultry litter containing wheat straw, bagasse or sawdust 7(1982)341

Slaughterhouse by-products preserved by *Lactobacillus plantarum* fermentation as feed for mink and foxes 20(1988)287

Fermentation–colon
In vitro fermentation of dietary fiber by human fecal organisms 23(1989)121

Approach to the energetic importance of fibre digestion in pigs. I. Importance of fermentation in the overall energy supply 23(1989) 141

Digestion and fermentation pattern of various dietary fiber sources in the rat 23(1989)195

The rumen as a model of the microbiology of fibre digestion in the human intestine 23 (1989)227

Fermentation–forage
A comparison between fermentation heat of forages and organic matter digestibility determined by in vitro incubation with rumen fluid 23(1989)333

Fermentation–fresh crops
The chemical composition and nutritive value for ruminants of four possible energy crops and their residues of anaerobic fermentation 15(1986)197

Fermentation–ilium
Dietary fiber and energy balance: integration of the human ileostomy and in vitro fermentation models 23(1989)261

Fermentation–in vitro
The effect of high levels of molasses in combinations with hay on digestibility of organic matter, microbial protein synthesis and volatile fatty acid production in vitro 7(1982)277

Fermentation–rumen
Nitrogen levels in low-roughage diets for efficient rumen microbial protein production 2(1977)255

Inhibition of ruminal deamination in vitro by formaldehyde treatment of sunflower-seed, soya bean and fish meals: response curves to protective treatment 2(1977)267

Rumen and post-rumen fermentation of spent tea leaf protein and other protein sources studied by the nylon bag method 7(1982) 221

The energetic efficiency of rumen microbial protein synthesis in cattle given silage-based diets 13(1985)255

The effect of monensin on rumen fermentation in sheep fed on all-roughage and barley/roughage diets 15(1986)7

In vitro effects of avoparcin on protein degradability and rumen fermentation 15 (1986)215

Evaluation of a dual flow continuous culture system for estimating bacterial fermentation in vivo of mixed diets containing various soya bean products 16(1986/1987)51

Fermentation of hammermilled barley, lupin, pea and faba bean grain in the rumen of dairy cows 16(1986/1987)261

Effect of straw treatment and laboratory of incubation on degradation of neutral detergent residue in the rumen 18(1987)75

The effect of grinding and pelleting hay on digestibility, fermentation rate, digesta passage and rumen and faecal particle size in cows 19(1988)145

Fibre degradation in the rumens of animals receiving hay, untreated or ammonia-treated straw 19(1988)277

The effect of five different supplements on the degradation of straw in sheep given untreated barley straw 19(1988)289

Rumen fermentation, urease activity and performance of cattle given palm kernel cake-based diet 20(1988)79

Formation and utilization of end products of ligno cellulose degradation in ruminants 21 (1988)141

The rumen as a model of the microbiology of fibre digestion in the human intestine 23 (1989)227

Fermentation and utilization by lambs of sugarcane harvested fresh and ensiled with and without NaOH. 4. Ruminal kinetics 23 (1989)323

Effect of urea treatment on rumen degradation characteristics of rice straws 24(1989) 83

Fermentation–silage
Effect of maize silage harvest stage on yield, plant composition and fermentation losses 1(1976)313

Studies on fermentation of sorghum silage during storage - effect of urea and molasses 3(1978)299

The effect of silage additives containing formaldehyde on the fermentation of ryegrass ensiled at different dry matter levels and on the nutritive value of direct-cut silage 7(1982)303

Studies on the fermentation of chopped sugarcane 7(1982)411

The effect of calcium and sodium hydroxides and of sodium acrylate on the fermentation and digestibility in vitro of ensiled whole-crop wheat and barley harvested at different stages of maturity 9(1983)37

Studies on fermentation of sorghum silage during storage, and its effect on milch animals 9(1983)143

Effect of wilting and the additives, straw, molasses and urea on the fermentation pattern of maize silage 9(1983)185

The effects of ammonia on the fermentation of chopped sugarcane 9(1983)291

Effect of sodium acrylate and calcium and sodium hydroxides on the fermentation, digestibility, chemical composition and aerobic deterioration of whole-crop maize silage 12(1984/1985)91

The effects of inoculation, addition of glucose and mincing on fermentation and proteolysis in ryegrass ensiled in laboratory silos 19 (1988)85

SO_2-treated straw as a silage additive: fermentation data on lucerne with particular reference to protein degradation 22(1988) 247

SO_2-treated straw as a silage additive: the participation of soluble and cell wall monosaccharide residues in lucerne silage fermentation 22(1988)255

Inoculation of corn silage and high-moisture corn with lactic acid bacteria and its effects on the subsequent fermentations and on feedlot performance of beef steers 25(1989) 23

Fertilizers

The quality of the green fraction of leaf protein concentrate from *Chenopodium quinoa* Willd. grown at different levels of fertilizer nitrogen 11(1984)239

The nutritional value of crop residue components from several wheat cultivars grown at different fertilizer levels 11(1984)301

Fractions of nitrogen, sulphur, phosphorus, calcium and magnesium in the herbage of perennial ryegrass as influenced by fertilizer nitrogen 14(1986)231

Fibre

A comparison of three fibre methods for predicting the metabolizable energy content of sorghum grain for poultry 2(1977)197

The digestibility and nutritive value for dairy heifers of rations containing cellulose fibre 2(1977)287

Cellulase insoluble fibre as a measure of unavailable organic matter in cattle compounds containing alkali-treated straw 3 (1978)227

Some remarks on the determination of acid-detergent fibre in feeds with cetyltrimethylammonium bromide versus determination of detergent fibre with alkylarylsulfonic acid 6(1981)91

The study of several modifications of the neutral detergent fibre procedure 9(1983)19

Chemical properties of fibre in relation to nutritive quality of ammonia-treated forages 10(1983/1984)155

The effect of level and source of protein, fibre and fat in the diet on the energy value of compounded pig feeds 11(1984)11

A new biological method for determining amino acid digestibility in poultry feedstuffs using a simple cannula, and the influence of dietary fibre on endogenous amino acid output 12(1984/1985)29

Effects of dietary concentrates on rumen digestion of fibrous feedstuffs 14(1986)193

The response of screened manure fibre to ozone and sodium hydroxide treatments 15 (1986)47

Influence of nutrient density on feed consumption, weight gain and gut capacity of broilers, Leghorns and turkeys reared to 26 days of age 16(1986/1987)129

The effects of different sources of nitrogen supplementation on the digestion of fibre components in the rumen of steers 17(1987) 65

Composition and fibre digestion in morphological components of an alfalfa–timothy sward 19(1988)135

Dietary fibre and nitrogen excretion and retention by pigs 19(1988)185

Fibre degradation in the rumens of animals receiving hay, untreated or ammonia-treated straw 19(1988)277

The effects of barley, unmolassed sugar-beet pulp and molasses supplements on organic matter, nitrogen and fibre digestion in the rumen of cattle given a silage diet 20(1988) 259

Effect of the nature of dietary fibre on transit time and faecal excretion in the growing pig 20(1988)327

Digestibility in vivo of ensiled grapevines (branches and leaves): influence of the system of analysis in the detergent fibre scheme on the prediction of digestibility 22 (1988)173

Measuring dietary fiber 23(1989)1

Determination of soluble fiber components: (1-<3; 1-<4)-β-D-glucans and pectins 23 (1989)15

Classification and measurement of plant polysaccharides 23(1989)27

Secretory response of the pig gut to non-starch polysaccharides 23(1989)55

Fibre-degrading microorganisms in the monogastric digestive tract 23(1989)91

In vitro fermentation of dietary fiber by human fecal organisms 23(1989)121

Approach to the energetic importance of fibre digestion in pigs. I. Importance of fermentation in the overall energy supply 23(1989) 141

Approach to the energetic importance of fibre digestion in pigs. II. An experimental approach to hindgut digestion 23(1989)169

Digestion and fermentation pattern of various dietary fiber sources in the rat 23(1989)195

Plant cell walls and monogastric diets 23 (1989)205

The rumen as a model of the microbiology of fibre digestion in the human intestine 23 (1989)227

The physiological effects of cellulose in the human large intestine 23(1989)241

Dietary fiber and energy balance: integration of the human ileostomy and in vitro fermentation models 23(1989)261

Comparative fiber degradation by mixed rumen fungi from Australian and U.S.A. cattle 23(1989)305

Response of high-producing dairy cows to high levels of dried corn gluten feed 24 (1989)191

The effect of a protein energy supplement on pasture protein and fibre digestion in the rumen of grazing steers 25(1989)39

Fish meal

Inhibition of ruminal deamination in vitro by formaldehyde treatment of sunflower-seed, soya bean and fish meals: response curves to protective treatment 2(1977)267

The involvement of trimethylamine oxide in fish meal in the production of egg taint 8 (1983)119

Substitution of finger millet (*Eleusine coracana*) and bulrush millet (*Pennisetum typhoides*) for maize in broiler feeds 10(1983/1984)291

Meat and bone and fish meals in balancer feeds for choice-fed broilers 11(1984)221

The amino acid and mineral composition of white fish meal containing enzyme-digested or untreated stickwater solids 14(1986)291

The effect of adding white fish meal containing enzyme digested or untreated stickwater solids to diets for broiler chickens 16 (1986/1987)253

Fish protein concentrates for pre-ruminant calves: effects of processing temperatures and source of fish on protein digestibility and biological value 18(1987)181

Fish protein hydrolates

Assessment of the production costs of fish protein hydrolysates 7(1982)147

Fish protein hydrolysates

General review of fish protein hydrolysates 7 (1982)113

Preparation of fish protein hydrolysates 7 (1982)125

Use of fish protein hydrolysate in milk replacers 7(1982)135

Performance of calves fed on milk replacers containing fish protein hydrolysate 7(1982) 141

Fish silage – preparation, properties and uses 7(1982)153

Fish silage

Nutritional value and mercury content of fish silage for growing pigs 9(1983)169

Thiaminase activity in fish silage and moist fish feed 13(1985)39

Effect of formaldehyde treatment on the degradation of acid-preserved fish silage protein in vitro 16(1986/1987)297

Fish silage as a protein supplement for early-weaned calves 17(1987)165

Examination of stability, and its effect on the nutritive value, of fish silage in diets for growing pigs 21(1988)43

Fodder beet roots

The chemical composition, digestibility and energy value of fodder-beet roots 18(1987) 225

Formaldehyde treatment

The digestion in the small intestine of young bulls of the protein of rapeseed meal treated or untreated with formaldehyde 7 (1982)225

Energy and nitrogen balance, intake and growth rate of lambs fed on unwilted grass silages treated with a formaldehyde–acid mixture or untreated 9(1983)107

Effect of formaldehyde treatment on the degradation of acid-preserved fish silage protein in vitro 16(1986/1987)297

Complexes of urea and formaldehyde as non-protein nitrogen compounds in ruminant rations: a review 20(1988)1

Effect of formaldehyde treatment on the rumen digestion of barley and wheat grain and its influence on hay intake and digestion by sheep 25(1989)55

Formalin

The composition of processed abattoir blood and its use in finishing-pig diets 2(1977)21

Fox

Slaughterhouse by-products preserved by *Lactobacillus plantarum* fermentation as feed for mink and foxes 20(1988)287

Fractionation

Possibilities for multipurpose fractionation and alternative uses of plant biomass 21 (1988)267

Fungi

Further thermotolerant fungi for the conversion of cassava starch to protein 2(1977)7

The value of 'Pekilo protein' for growing pigs 2(1977)161

Production of fungal protein from rasped fresh cassava roots using 200- and 3000-litre fermentors 8(1983)313

The role of ciliate protozoa and fungi in the rumen digestion of plant cell walls 10(1983/1984)121

Effects of *Drechslera sorokiniana* infection on the yield and quality of tall fescue forage 15(1986)41

Nutrient losses during deterioration of hay in relation to changes in biochemical composition and microbial growth 15(1986)149

Association of rumen fungi with various forage grasses 16(1986/1987)273

Colonization of thick-walled plant tissues by anaerobic fungi 19(1988)25

Fungal contamination of hydroponic forage 20 (1988)163

Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. I. Composition and nutritive evaluation 22(1988)1

Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. II. Effect of method of drying and response to amino acid supplementation 22(1988)13

Colonization of guinea grass by anaerobic rumen fungi in swamp buffalo and cattle 22 (1988)161

Comparative fiber degradation by mixed rumen fungi from Australian and U.S.A. cattle 23(1989)305

G

Garden orach
The chemical composition and nutritive value for ruminants of four possible energy crops and their residues of anaerobic fermentation 15(1986)197

Genetics
Genetic variation in the feeding value of cereal straw 21(1988)127

Germination
The role of germination in sorghum reconstitution 14(1986)243

Giant star grass
The effect of age of regrowth on content and release of manganese, iron, zinc and copper from four tropical forages incubated in sacco in rumen of sheep 20(1988)171

(1-<3; 1-<4)-β-D-glucans
Determination of soluble fiber components: (1-<3; 1-<4)-β-D-glucans and pectins 23 (1989)15

Glucose
The effects of inoculation, addition of glucose and mincing on fermentation and proteolysis in ryegrass ensiled in laboratory silos 19 (1988)85

Glucosinolate
Low-glucosinolate rapeseed products for pigs – a review 9(1983)239

Glutamic acid
Amino acid composition of maize silage produced in the United Kingdom 4(1979)163

Gluten
Evaluation of corn gluten meal with urea as a source of supplementary nitrogen for growing calves and lambs 7(1982)375

Goat
Seasonal changes in composition and intake of chaparral by Spanish goats 8(1983)47

Utilization of alkali-treated and neutralized wheat straw-based rations for growing goat kids 9(1983)155

The value of urea-treated straw in diets of lactating goats 11(1984)67

Performance and carcass characteristics of Sudan desert sheep and goats on high roughage diets with added fat 13(1985)147

Changes in nutrient intake and performance by goats fed coffee pulp-based diets followed by a commercial concentrate 14 (1986)1

The value of banana stalk as a feed for goats 15(1986)95

Effect of sodium hydroxide treatment of desert grass on carbohydrate digestibility by goats 17(1987)219

Cassava meal as supplement to napier grass diets for growing sheep and goats 18(1987) 197

Comparison of cottonseed cake and dried blood as nitrogen sources for growing Sudan Nubian goats 22(1988)333

The intake and digestibility in goats of *Leucaena leucocephala* supplemented with dehydrated palm oil mill effluent 24(1989)45

Upgrading barley straw for goats: the effectiveness of a sodium hydroxide and urea dip method 24(1989)57

Grain
High grain content maize silage in fattening diets of young male cattle 16(1986/1987)63

Gram
Nutritional evaluation of gram (*Cicer arietinum*) and arhar (*Cajanus cajan*) chunies for cattle 5(1980)315

Grapevine
Digestibility in vivo of ensiled grapevines (branches and leaves): influence of the system of analysis in the detergent fibre scheme on the prediction of digestibility 22 (1988)173

Grass, general
The accuracy of the pepsin–cellulase technique for estimating the dry matter digestibility in vivo of grasses and legumes 3(1978)277

Feed consumption and performance of dairy cows with alternate feedings of grass silage and maize silage 6(1981)337

Chemical composition and animal performance of grass forage of varying maturity stored as hay or silage 11(1984)35

Association of rumen fungi with various forage grasses 16(1986/1987)273

Grass, tropical
Bias when predicting crude protein, dry matter digestibility and voluntary intake of tropical grasses by near-infrared reflectance 9(1983)221

Effects of ensiling or drying on five forage species in western Kenya: *Zea mays* (maize stover), *Pennisetum purpureum* (Pakistan napier grass), *Pennisetum* sp. (bana grass), *Impomea batata* (sweet potato vines) and *Cajanus cajan* (pigeon pea leaves) 13(1985) 1

Grass, tropical pasture
Prediction of the dry matter digestibility of tropical grasses using a pepsin-cellulase assay 2(1977)247

Solubility of the protein of tropical pasture species and the rate of its digestion in the rumen 5(1980)183

Grass juice
Value of lucerne juice and grass juice as sources of protein for the growing pig 4 (1979)233

Grass silage
The use of effluent from grass silage in the diet of finishing pigs. I. Variation in composition of effluents 4(1979)263

The use of effluent from grass silage in the diet of finishing pigs. II. Assessment of nutritive value of fresh and stored effluent 4(1979) 275

Effect of soaking, ensilage and hydrogen peroxide treatment of barley straw on rumen degradability 22(1988)295

Green berseem
Ensiling characteristics of cattle waste fermented with wheat straw and green berseem (*Trifolium alexandrium*) 25(1989)149

Green panic
Preservation and storage of green panic (*Panicum maximum*) as moist hay with urea 20 (1988)87

Grinding
The influence of steam pelleting and grinding on the nutritive value of Canola rapeseed for poultry 8(1983)303

Ground mesquite
Pilot plant production of sulphur dioxide-treated mesquite wood. II. Digestibility and feedlot performance by lambs 22(1988) 61

Groundnut cake
Feeding value of sunflower and groundnut cakes for broilers 4(1979)143

Feeding value of sunflower- and groundnut-cakes for laying hens 6(1981)63

Groundnut hulls
The effect of grinding groundnut hulls either with or without alkali treatment on digestibility of diet and on ruminal and blood components 15(1986)69

Groundnut meal
Detoxification of aflatoxins in groundnut meal 2(1977)67

Growth promotors
Energy metabolism in lambs fed silage diets with sucrose or a growth promoter 16 (1986/1987)203

Guinea fowl
A comparison of metabolizable energy content of eight common feed ingredients determined with young guinea fowls (keets) and pullet chicks 15(1986)1

Guinea grass
The effect of age of regrowth on content and release of manganese, iron, zinc and copper from four tropical forages incubated in sacco in rumen of sheep 20(1988)171

Colonization of guinea grass by anaerobic rumen fungi in swamp buffalo and cattle 22 (1988)161

H

Hair
Utilization of processed pig hair in poultry diets 3(1978)243

Protein digestibility of hydrolyzed hog hair meal for pigs 3(1978)323

Harvesting time

Preservation of sorghum plant portions harvested, processed and ensiled at ten stages of maturity 7(1982)257

Influence of harvest date on the nutritive value and ensiling characteristics of maize stover 14(1986)11

The chemical composition of pea plant parts as related to harvesting time 16(1986/1987) 305

Yield, composition and feeding value of irrigated *Sesbania sesban* var. *Nubia* leaves harvested at latitude 38°N during a Mediterranean summer 18(1987)247

Treatment of whole-crop cereals with alkali. 1. The influence of sodium hydroxide and ensiling on the chemical composition and in vitro digestibility of rye, barley and wheat crops harvested at increasing maturity and dry matter content 18(1987)257

The nutritive value of ensiled forages cut at three stages of growth 22(1988)237

Hay

Apparent and true digestibility of the Weende components, cell content and cell wall of ryegrass 1(1976)607

Differences between cattle and sheep in their digestion and relative intake of a mature tropical grass hay 3(1978)41

Estimation of the digestibility of low-quality hays by cattle from measurements made with sheep 3(1978)51

Laboratory application of preservatives to hay and the effects of irregular distribution on mould development 6(1981)73

Chemicals for preventing moulding in damp hay 6(1981)323

Utilization of alkali-treated grain. 2. Utilization by steers of diets based on hay or straw and mixed with either NaOH-treated or rolled barley 6(1981)347

The effect of high levels of molasses in combinations with hay on digestibility of organic matter, microbial protein synthesis and volatile fatty acid production in vitro 7(1982)277

Laboratory tests of 8-quinolinol as an additive to volatile fatty acids for preserving hay and other feed-stuffs 7(1982)287

Ammoniation of wheat straw and native grass hay during baling of large round bales 8 (1983)35

The effects of sulphur supplementation on cellulose digestion in vitro and on nutrient digestion, nitrogen metabolism and rumen characteristics of lambs fed on good quality fescue and tropical star grass hays 8(1983) 129

Methods of improving the utilization of cereal straw by ruminants. I. Supplements of urea, molasses and dried grass and treatment with sodium hydroxide 8(1983)221

Methods of improving the utilization of cereal straw by ruminants. II. Further studies on dried grass as a supplement for lambs 8 (1983)229

Problems of testing novel chemicals for the preservation of damp hay 8(1983)283

Manipulation of rumen fluid pH and its influence on cellulolysis in sacco, dry matter degradation and the rumen microflora of sheep offered either hay or concentrate 10 (1983/1984)1

Associative effects of mixed feeds. II. The effect of dietary addition of bicarbonate salts on the voluntary intake and digestibility of diets containing various proportions of hay and barley 10(1983/1984)31

Chemical composition and animal performance of grass forage of varying maturity stored as hay or silage 11(1984)35

The effect of anhydrous ammonia and moisture content on the preservation and chemical composition of perennial ryegrass hay 11(1984)159

The effect of intake and hay: concentrate ratio upon digestibility and digesta passage 11 (1984)167

Digestibility and digesta retention in dairy cows receiving hay or silage at varying concentrate levels 11(1984)279

A comparison of laboratory methods involving single or two-stage cellulase and two-stage rumen inoculum–pepsin treatment for predicting digestibility and intake of hays in sheep 12(1984/1985)101

The voluntary intake and digestibility of combinations of cereal crop residues and legume hay for sheep 12(1984/1985)241

Effects of ensiling or drying on five forage species in western Kenya: *Zea mays* (maize stover), *Pennisetum purpureum* (Pakistan napier grass), *Pennisetum* sp. (bana grass), *Impomea batata* (sweet potato vines) and *Cajanus cajan* (pigeon pea leaves) 13(1985) 1

Storage of large round bales of hay harvested at different moisture concentrations and treated with sodium diacetate and/or covered with plastic 13(1985)69

Effect of ammonia level and time of exposure to ammonia on nutritional and preservatory characteristics of dry and high-moisture coastal bermuda grass hay 14 (1986)55

Dicoumarol concentrations in sweetclover hay treated with preservatives and in spoiled hay of high- and low-coumarin cultivars of sweetclover 14(1986)221

The effect of monensin on rumen fermentation in sheep fed on all-roughage and barley/roughage diets 15(1986)7

Nutrient losses during deterioration of hay in relation to changes in biochemical composition and microbial growth 15(1986)149

Estimating yield and nutrient losses due to rainfall on field-drying alfalfa hay 16(1986/1987)7

Digestibility of dry matter, nitrogen and energy by dairy cows fed whole or hammer-milled lupin grain in oaten hay or oaten pasture based diets 16(1986/1987)143

Microfloral and compositional changes in alfalfa hay treated with sodium diacetate and stored at different moisture contents 17(1987)45

Chemical composition and digestion in vitro of orchardgrass hay ammoniated by different techniques 17(1987)109

Effects of whole or ground corn with different forms of hay in 85% concentrate diets on digestion and passage rate in beef heifers 18(1987)151

The effect of grinding and pelleting hay on digestibility, fermentation rate, digesta passage and rumen and faecal particle size in cows 19(1988)145

Fibre degradation in the rumens of animals receiving hay, untreated or ammonia-treated straw 19(1988)277

Evaluation of mixed lucerne and caucasian bluestem hay diets fed to sheep 20(1988) 19

Nitrogen utilization, ruminal fermentation and abomasal nitrogen flow in sheep fed caucasian bluestem hay supplemented with lucerne or urea 20(1988)33

Preservation and storage of green panic (*Panicum maximum*) as moist hay with urea 20 (1988)87

Effect of level and frequency of maize supplementation on the utilization of low-quality grass hay by beef cows 24(1989)129

Oven and stack ammoniation of grass hays. 1. Changes in chemical composition in relation to digestibility in vitro and cell-wall degradability 24(1989)299

Oven and stack ammoniation of grass hays. 2. Voluntary intake, digestibility and faecal nitrogen excretion in sheep in relation to hay composition 24(1989)313

Effect of formaldehyde treatment on the rumen digestion of barley and wheat grain and its influence on hay intake and digestion by sheep 25(1989)55

Water-drinking frequency and quantity of water drunk by steers fed hay or fresh forage 25(1989)207

Hominy

Feeding and nutritive value of hominy and corn grits for poultry 19(1988)313

House fly larvae

House fly larvae meal grown on municipal organic waste as a source of protein in poultry diets 4(1979)227

Hydrilla

Utilization of aquatic plants *Elodea canadensis* and *Hydrilla verticillata* in broiler chick diets 20(1988)155

Hydrolysis

Estimation of lysine in compounded poultry diets after enzymic digestion 3(1978)57

Hydroponic forage

Fungal contamination of hydroponic forage 20 (1988)163

Hyperexcitability

Hyperexcitability in cattle fed ammoniated roughages 17(1987)121

I

Immunoglobulins

A continuous process for the production of purified porcine γ-globulin for use in pig-milk replacer supplements 17(1987)213

Intake

Nutritive value of thermoammoniated and steam-treated maize stover I. Intake, digestibility and nitrogen retention 4(1979) 177

Effect of dry-matter concentration of milk substitute and method of feeding on intake and performance by lambs 5(1980)321

Dry matter, protein, energy and fibre intake by dairy heifers grazing a Rhodes grass (*Chloris gayana*) pasture 6(1981)15

Intake and digestibility in sheep fed on alkali-treated straw and concentrates, as complete diets or separate feeds 7(1982)271

Seasonal changes in composition and intake of chaparral by Spanish goats 8(1983)47

Energy and nitrogen balance, intake and growth rate of lambs fed on unwilted grass silages treated with a formaldehyde–acid mixture or untreated 9(1983)107

Effect of intake on digestibility of plant cell walls and cell contents of complete diets for ruminants 9(1983)197

Bias when predicting crude protein, dry matter digestibility and voluntary intake of tropical grasses by near-infrared reflectance 9(1983)221

Associative effects of mixed feeds. II. The effect of dietary addition of bicarbonate salts on the voluntary intake and digestibility of diets containing various proportions of hay and barley 10(1983/1984)31

The effect of milling and sodium hydroxide treatment on intake and digestibility of wheat straw by sheep 11(1984)149

The effect of intake and hay: concentrate ratio upon digestibility and digesta passage 11 (1984)167

A comparison of laboratory methods involving single or two-stage cellulase and two-stage rumen inoculum–pepsin treatment for predicting digestibility and intake of hays in sheep 12(1984/1985)101

Improving the intake and nutritive value of wheat straw for beef cows 12(1984/1985) 125

The utilization of untreated and calcium hydroxide treated wheat straw by sheep 12 (1984/1985)141

The effect of supplementing grass silage with sodium hydroxide-treated chopped or milled straw on intake and digestibility by wether sheep 12(1984/1985)215

The voluntary intake and digestibility of combinations of cereal crop residues and legume hay for sheep 12(1984/1985)241

Methods of improving the utilization of cereal straw by ruminants. IV. The effect on intake of level of supplementation with urea and dried grass 13(1985)281

Changes in nutrient intake and performance by goats fed coffee pulp-based diets followed by a commercial concentrate 14 (1986)1

Aqueous ammonia treatment of wheat straw: voluntary intake and digestibility in cattle 14(1986)81

Effects of feeding alkali-treated olive cake on intake, digestibility and rumen liquor parameters 14(1986)139

Effects of supplementing a wheat straw diet with urea or a urea–molasses block and/or cottonseed meal on intake and liveweight change of lambs 16(1986/1987)25

The intake and digestibility by sheep of oat straw and maize stover offered with different levels of noug (*Guizotia abyssinica*) meal 16(1986/1987)99

Intake, digestibility and energy utilization of corrugated paper diets 17(1987)57

Treatment of whole-crop cereals with alkali. 2. Voluntary intake and digestibility by sheep of rye, barley and wheat crops ensiled with sodium hydroxide 18(1987)271

Treatment of whole-crop cereals with alkali. 3. Voluntary intake and digestibility studies in sheep given immature wheat ensiled with sodium hydroxide, urea or ammonia 18(1987)283

Treatment of whole-crop cereals with alkali. 4. Voluntary intake and growth in steers given wheat ensiled with sodium hydroxide, urea or ammonia 19(1988)55

The intake and digestion of maize silage-based diets by dairy cows and sheep 20 (1988)299

Feed intake and milk yield at increasing supplement of a palmitic and stearic acid-rich type of fat in comparison with animal fat 22(1988)179

Feed intake, milk yield and milk composition by replacing unprotected fat by Ca-soaps for dairy cows 22(1988)193

The intake and digestibility in goats of *Leucaena leucocephala* supplemented with dehydrated palm oil mill effluent 24(1989)45

Oven and stack ammoniation of grass hays. 2. Voluntary intake, digestibility and faecal nitrogen excretion in sheep in relation to hay composition 24(1989)313

Effect of formaldehyde treatment on the rumen digestion of barley and wheat grain and its influence on hay intake and digestion by sheep 25(1989)55

Intake, digestion and daily gain by cattle consuming bermudagrass (*Cynodon dactylon*) and supplemented with different combinations of ground corn, vegetable oil, urea and corn gluten and blood meals 25(1989)99

Iodine

Copper and iodine in pig diets with high glucosinolate rapeseed meal. I. Performance and thyroid hormone status of growing pigs fed on a diet with rapeseed meal treated with copper sulphate solution or untreated and supplements of iodine, copper or a quinoxaline derivative 22(1988)33

Copper and iodine in pig diets with high glucosinolate rapeseed meal. II. Influence of iodine supplements for rations with rapeseed meal untreated or treated with copper ions on performance and thyroid hormone status of growing pigs 22(1988)45

Iron

The effect of age of regrowth on content and release of manganese, iron, zinc and copper from four tropical forages incubated in sacco in rumen of sheep 20(1988)171

Irrigation
Cultivation of maize in irrigated areas for silage and grain production and their use by ruminants 1(1976)237

Isoacids
Isoacids in ruminant nutrition: their role in ruminal and intermediary metabolism and possible influences on performances – a review 18(1987)169

Isopropyl alcohol treatment
Effects of isopropyl alcohol treatment of soya bean meal on nitrogen utilization by ruminants 14(1986)127

J

Jackbean
Ripe beans of *Canavalia ensiformis* (jackbean) as feed ingredient for monogastric animals 20(1988)325

K

Kapok seed
The effect of nutrient supplementation on the feeding value of kapok seed for poultry 9 (1983)263

Karanja
The effect of kernels of karanja (*Pongamia glabra* Vent) on growth and feed efficiency in broiler chicks to 4 weeks of age 25(1989) 201

Keratin
Utilization of processed pig hair in poultry diets 3(1978)243

Protein digestibility of hydrolyzed hog hair meal for pigs 3(1978)323

Chemical preparation of NaOH–keratin hydrolysate for improving the nutritive value of wheat straw 13(1985)215

Kidney bean
The use of kidney beans (*Phaseolus vulgaris*) as protein supplements in diets for young rapidly growing beef steers 12(1984/1985) 1

L

Lactate
Effect of lactate on the in vitro digestion of *Agropyron elongatum* by rumen microorganisms 16(1986/1987)161

Lactobacillus
The chemical composition of grass silages prepared with and without pre-treatment with inoculants containing *Lactobacillus plantarum* 13(1985)269

Slaughterhouse by-products preserved by *Lactobacillus plantarum* fermentation as feed for mink and foxes 20(1988)287

Lactobacillus
Efficacy of virginiamycin and a commercially-available lactobacillus probiotic in swine diets 8(1983)69

Stability of viable lactobacillus products 11 (1984)261

Lantana seeds
Can lantana seeds be utilized in lamb rations? 9(1983)29

Leaf protein concentrate
Nutritive value of lupin-seed protein and leaf-protein concentrates supplemented with various sulphur sources 6(1981)189

Nutritive value of leaf-protein concentrate from potato haulm for rats and chicks 6 (1981)413

Yield, composition and nutritive value of a leaf protein concentrate obtained by fractionating the shelled aerial part of *Pisum savitum* L. 8(1983)155

The quality of the green fraction of leaf protein concentrate from *Chenopodium quinoa* Willd. grown at different levels of fertilizer nitrogen 11(1984)239

The effect of different precipitating agents on quality of leaf protein concentrate from lucerne 12(1984/1985)11

The effect of toxic substances present in some leaf protein concentrates on their nutritive value 24(1989)151

Legumes
The analytical and biological accuracy of estimating the dry matter digestibility of different legume species 1(1976)61

The accuracy of the pepsin–cellulase technique for estimating the dry matter digestibility in vivo of grasses and legumes 3(1978)277

Ensiling of leguminous green forages in combination with different dry roughages and molasses 12(1984/1985)133

The voluntary intake and digestibility of combinations of cereal crop residues and legume hay for sheep 12(1984/1985)241

Rumen degradability of nitrogen in typical cultivated grasses and leguminous fodders 22(1988)79

Leucaena leucocephala
A note on the nutrient and mineral composition of *Leucaena leucocephala* in Western Nigeria 4(1979)221

Growth performance of, and mimosine excretion by, young chicks fed on *Leucaena leucocephala* 7(1982)247

Investigations into the effects of processing on the retention of the carotenoid fractions of *Leucaena leucocephala* during storage, and the effects of processing on mimosine concentration 9(1983)307

Nutrient and amino acid composition of *Leucaena leucocephala* (Lam.) de Wit 15(1986) 183

The intake and digestibility in goats of *Leucaena leucocephala* supplemented with dehydrated palm oil mill effluent 24(1989)45

Lignification
Botanical composition and extent of lignification affecting digestibility of wheat and oat straw and paspalum hay 7(1982)71

Lignocellulose
Formation and utilization of end products of ligno cellulose degradation in ruminants 21 (1988)141

Production of ethanol from lignocellulosic materials 21(1988)175

Chemistry of lignocellulose: methods of analysis and consequences of structure 21(1988) 279

Biological structure of lignocellulose and its degradation in the rumen 21(1988)295

Listeria monocytogenes
Inhibition of *Listeria monocytogenes* by defined lactic microflora in gnotobiotic silages of lucerne, fescue, ryegrass and maize – influence of dry matter and temperature 2 (1977)297

Lucerne
Nutritive value of leaf protein coagulum and cereals mixtures 4(1979)91

The effect of different precipitating agents on quality of leaf protein concentrate from lucerne 12(1984/1985)11

The chemical composition and nutritive value for ruminants of four possible energy crops and their residues of anaerobic fermentation 15(1986)197

Nitrogen utilization, ruminal fermentation and abomasal nitrogen flow in sheep fed caucasian bluestem hay supplemented with lucerne or urea 20(1988)33

Lucerne juice
Value of lucerne juice and grass juice as sources of protein for the growing pig 4 (1979)233

Value of freshly produced lucerne juice as a source of supplemental protein for the growing pig 5(1980)215

Lucerne juice as a protein supplement for growing pigs: effects of mineral content of the diet and of the water supply 6(1981)35

Lucerne straw
SO_2-treated straw as a silage additive: the participation of soluble and cell wall monosaccharide residues in lucerne silage fermentation 22(1988)255

Lupin
Variability for protein and oil quality in *Lupinus albus* 9(1983)271

Fermentation of hammermilled barley, lupin, pea and faba bean grain in the rumen of dairy cows 16(1986/1987)261

Lupin grain
Milling barley and lupin grain in diets for cattle 12(1984/1985)57

Digestibility of dry matter, nitrogen and energy by dairy cows fed whole or hammermilled lupin grain in oaten hay or oaten pasture based diets 16(1986/1987)143

Lupin seed
The alkaloid content of sweet lupin seed used in feeding trials on pigs and rats 2(1977)59

Nutritive value of lupin-seed protein and leaf-protein concentrates supplemented with various sulphur sources 6(1981)189

Rumen degradation of protein in sunflower meal, rapeseed meal and lupin seed placed in nylon bags 11(1984)87

Digestibility and energy value of sweet lupin seed (*Lupinus albus* var. multolupa) in pigs 12(1984/1985)171

Protein and energy utilisation in rats of diets based on lupin seed (*Lupinus albus* var. multolupa) 15(1986)33

Lupin seed meal
A comparison between meals prepared from the seeds of different varieties of lupin as protein supplements to barley-based diets for growing pigs 2(1977)49

Lupinus angustifolius cv. Unicrop as a protein- and energy-source for weaner pigs 6(1981) 27

Lupin-seed meal (*Lupinus albus* cv. Hamburg) as a source of protein for growing pigs 6(1981)285

Lupin-seed meal
Lupin-seed meal (*Lupinus angustifolius* cv. Uniwhite) as a protein supplement to barley-based diets for growing pigs 1(1976) 631

Lysine
Adjustments of the protein level for slaughter pigs and supplementation with lysine and methionine 1(1976)619

The composition of processed abattoir blood and its use in finishing-pig diets 2(1977)21

Estimation of lysine in compounded poultry diets after enzymic digestion 3(1978)57

Meat meal in the diet of the early-weaned pig. II. Amino acid supplementation 4(1979)43

Amino acid composition of maize silage produced in the United Kingdom 4(1979)163

The determination of available lysine in barley using *Tetrahymena pyriformis* W. 5 (1980)33

A comparison of three methods for the determination of available lysine in barley 5 (1980)43

Meat meal in the diet of the early-weaned pig. IV. The supplementation of diets with tryptophane, lysine and methionine 7 (1982)27

Lysine requirements of imported European pigs in the humid lowland tropics from 23 to 62 kg liveweight 7(1982)233

Proximate analysis, amino acid composition and tannin content of improved Nigerian sorghum varieties and their potential in poultry feeds 7(1982)359

Evaluation of optimum lysine and threonine supplements to a wheat and barley-based diet in rats 8(1983)163

Lysine requirement of weanling European pigs fed on yellow maize and groundnut cake diets in the humid lowland tropics 8 (1983)197

The absorption of dietary L-lysine and extruded L-lysine in pigs 15(1986)141

The influence of oat groats and dietary level of lysine on the laying performance of crossbred hens 17(1987)271

The utilization by beef steers of a soft high lysine and a hard normal size maize cultivar in whole grain feedlot diets 21(1988)31

Lysinoalanine
Note on the lysinoalanine content of alkali-treated barley 7(1982)217

M

Magnesium
Fractions of nitrogen, sulphur, phosphorus, calcium and magnesium in the herbage of perennial ryegrass as influenced by fertilizer nitrogen 14(1986)231

Maize–cobs
Effect of level of sodium hydroxide treatment and volume of solution on the nutritive value of maize cobs 4(1979)1

Performance of heifers fed on diets based on NaOH-treated maize cobs and the effect of supplementary urea and source of carbohydrates 4(1979)97

Chemical composition, digestibility and feeding value of maize cobs 9(1983)121

High temperature and pressure processing of maize cobs. II. Animal response and effect of sodium metabisulphite 14(1986)161

Maize–ear
Maize grain or ears in concentrate diets for young fattening bulls 1(1976)393

Maize–general
Survey of the maize situation in Italy 1(1976) 109

Review of production and utilisation of maize for animal nutrition in Yugoslavia 1(1976) 115

Use of maize for ruminants and pig feeding in Greece 1(1976)125

Survey of the use of maize for livestock feeding in Spain 1(1976)129

Survey of the use of maize for livestock feeding in Portugal 1(1976)137

Maize and animal production in France 1 (1976)141

Survey of the use of maize for livestock feeding in Switzerland 1(1976)149

Survey of the use of maize for livestock feeding in Austria 1(1976)153

Use of maize for livestock feeding in the Federal Republic of Germany 1(1976)157

Survey of the use of maize for livestock feeding in Belgium 1(1976)167

Use of maize for livestock feeding in the Netherlands 1(1976)177

Prospects of growing maize in Finland 1 (1976)187

The current situation regarding the use of maize in Ireland 1(1976)195

The use of maize for livestock feeding in the United Kingdom 1(1976)199

Discussion on session I 1(1976)215

The use of techniques of maize cultivation to maximise feed for beef production 1(1976) 229

Cultivation of maize in irrigated areas for silage and grain production and their use by ruminants 1(1976)237

Selection objectives for maximum yields of feed with hybrid maize 1(1976)245

Effect of agronomic factors on the yield of maize and on its composition in relation to its use by ruminants 1(1976)251

Agronomic factors affecting the growth and composition of the maize plant 1(1976)263

Discussion on session 2 1(1976)273

Efficiency of harvesting and conservation 1 (1976)289

Discussion on session 3 1(1976)327

Maize silage with or without NPN, dehydrated whole-crop maize pellets or high moisture maize grain for finishing bulls 1 (1976)347

Utilising the maize crop in various forms for beef production 1(1976)417

Nutritive value of dehydrated whole-crop maize 1(1976)429

The digestion and metabolism of the energy and protein of maize 1(1976)455

Discussion on session 4 1(1976)485

For beef cattle, all forms of maize can be used 1(1976)521

Bull-beef production from the maize crop in Italy in large and small units 1(1976)531

Discussion on session 5 1(1976)545

Requirements for future research 1(1976)555

Abomasal and plasma nitrogen constituents of steers fed on different physical forms of shelled maize 2(1977)225

Waxy versus regular maize:energy value for growing pigs and utilization in piglet diets 4(1979)109

Preservation of high-moisture maize – a comparison of gaseous and liquid anhydrous ammonia with methylene-*bis*-propionate 5 (1980)337

Effects of milling and pelleting of maize, barley and wheat on their metabolizable energy value for cockerels and chicks 9(1983) 99

Measurement and prediction of digestibility of forage maize in The Netherlands 10 (1983/1984)301

The evaluation of fresh sweet liquid whey as a protein supplement to maize for growing pigs 14(1986)171

Response of different sorghum grain and maize varieties to reconstitution 15(1986) 231

The rate of dry matter disappearance from nylon bags of maize silage/grain diets as influenced by basal ration and processing of test samples 16(1986/1987)225

High-fat maize in diets for pigs and sows 17 (1987)201

The maize replacement value of cassava flour in rations for growing ewe lambs 20(1988) 233

Nutritive value of diets containing triticale and varying mixtures of triticale and maize for growing-finishing swine 22(1988)217

Effect of level and frequency of maize supplementation on the utilization of low-quality grass hay by beef cows 24(1989)129

Inoculation of corn silage and high-moisture corn with lactic acid bacteria and its effects on the subsequent fermentations and on feedlot performance of beef steers 25(1989) 23

Maize–grain

Maize grain or ears in concentrate diets for young fattening bulls 1(1976)393

Feeding value of ensiled high-moisture maize grain with cob for growing-finishing pigs 20 (1988)145

Maize–silage

Forage maize in the rations of dairy cows and fattening cattle in Bulgaria 1(1976)45

Maize as a forage crop in Denmark 1(1976) 183

Influence of ensiling maize ears and grain on chemical composition, conservation losses and digestibility 1(1976)301

Effect of maize silage harvest stage on yield, plant composition and fermentation losses 1(1976)313

Energy supplementation of maize silage harvested at different maturity stages 1(1976) 369

Factors influencing the composition and nutritive value of ensiled whole-crop maize 1 (1976)381

Rate of growth, feed utilisation and digestibility observed with diets containing ensiled maize grain or maize ears fed alone or in mixtures 1(1976)401

Voluntary intake and efficiency of utilisation of whole-crop maize silage 1(1976)441

Levels, sources and methods of nitrogen supplementation of maize silage for beef production 1(1976)465

The nutritive value (energy) of maize silage 1 (1976)481

Effect of weight at slaughter on the efficiency of fattening young cattle 1(1976)505

Economic aspects of high-plane feeding of bulls with maize silage 1(1976)513

Maize silage and whole maize plant pellets as the basic forages in complete feeds for milk cows 3(1978)23

A note on the effect on the aerobic deterioration of maize silage of the manipulation of the microflora by means of antibiotics 3 (1978)89

Laboratory evaluation of ensiled mixtures of alkali-treated cattle excreta and whole-crop maize 3(1978)335

Amino acid composition of maize silage produced in the United Kingdom 4(1979)163

Effect of wilting and the additives, straw, molasses and urea on the fermentation pattern of maize silage 9(1983)185

Effect of sodium acrylate and calcium and sodium hydroxides on the fermentation, digestibility, chemical composition and aerobic deterioration of whole-crop maize silage 12(1984/1985)91

The effect of nitrogen supplements to maize silage on the amount and composition of the nitrogen fraction entering the intestine of cattle 13(1985)25

High grain content maize silage in fattening diets of young male cattle 16(1986/1987)63

The intake and digestion of maize silage-based diets by dairy cows and sheep 20 (1988)299

Maize–solubles

Effects of dietary maize solubles and minerals on albumen quality of fresh and stored brown-shelled eggs 8(1983)177

Maize–starch

Effect of defaunation on the loss of energy in wethers fed different quantities of cellulose and normal or steamflaked maize starch 16 (1986/1987)233

Maize–stem

Colonization of thick-walled plant tissues by anaerobic fungi 19(1988)25

Maize–stover

Nutritive value of thermoammoniated and steam-treated maize stover I. Intake, digestibility and nitrogen retention 4(1979) 177

Nutritive value of single-cell protein produced by *Chaetomium cellulolyticum* grown on maize stover and pulp-mill sludge 11(1984) 211

Effects of ensiling or drying on five forage species in western Kenya: *Zea mays* (maize stover), *Pennisetum purpureum* (Pakistan napier grass), *Pennisetum* sp. (bana grass), *Impomea batata* (sweet potato vines) and *Cajanus cajan* (pigeon pea leaves) 13(1985) 1

Influence of harvest date on the nutritive value and ensiling characteristics of maize stover 14(1986)11

The intake and digestibility by sheep of oat straw and maize stover offered with different levels of noug (*Guizotia abyssinica*) meal 16(1986/1987)99

Effects of moisture level, treatment time and soya bean addition on the nutritive value of urea-treated maize stover 19(1988)67

Effects of ruminal degradability of the protein supplement on the utilization of maize-stover silage by growing beef cattle 20 (1988)241

Effects of added dietary urea on the utilization of maize stover silage by growing beef cattle 22(1988)227

Maize–whole

The utilization by beef steers of a soft high lysine and a hard normal size maize cultivar in whole grain feedlot diets 21(1988)31

Maize–whole stillage

Maize whole stillage as a supplemental protein source for growing lambs and finishing steers 11(1984)181

Maize silage

Effects of stage of maturity and addition of molasses on nutritive value of maize stover silage 12(1984/1985)65

Maltase

Proximate chemical analysis of final cane molasses and effect of feeding 30% molasses on intestinal sucrase and maltase activities in the rat 3(1978)15

Man

Assessing the importance of host-derived polysaccharides as carbon sources for bacteria growing in the human colon 23(1989) 109

In vitro fermentation of dietary fiber by human fecal organisms 23(1989)121

The physiological effects of cellulose in the human large intestine 23(1989)241

Dietary fiber and energy balance: integration of the human ileostomy and in vitro fermentation models 23(1989)261

Manganese
The effect of age of regrowth on content and release of manganese, iron, zinc and copper from four tropical forages incubated in sacco in rumen of sheep 20(1988)171

Manioc
The digestion of diets containing manioc by young growing pigs 12(1984/1985)119

Manure
Utilization by sheep of dried pig faeces containing a high concentration of copper 3 (1978)31

Digestibility by cattle of diets containing dried pig faeces 3(1978)155

Laboratory evaluation of ensiled mixtures of alkali-treated cattle excreta and whole-crop maize 3(1978)335

The effects of air dried poultry manure on performance, organ weights and haematological measurements of weaner pigs reared in the tropics 4(1979)23

Utilization of cattle manure for single-cell protein production with *Chaetomium cellulolyticum* 4(1979)199

Digestibility of wet cage layer excreta and an excreta-supplemented diet by cattle as determined by two methods 5(1980)77

Utilization of cattle manure for Torula yeast production from straw hydrolysates 5(1980) 175

Performance of fattening cattle and lactating beef cows on diets high in chemically treated wheat straw and poultry litter 5 (1980)299

Investigations of the treatment of sawdust for rabbit feeding. 2. Effect of boiling and incubation with poultry manure 6(1981)51

The determination of digesta partical size in some herbivores 7(1982)35

Fermentation characteristics and feeding value of ensiled poultry litter containing wheat straw, bagasse or sawdust 7(1982)341

Nutritive value of ensiled broiler litter for cattle 8(1983)19

Corrugated cardboard and dried poultry manure as dietary ingredients for sheep 14 (1986)265

The response of screened manure fibre to ozone and sodium hydroxide treatments 15 (1986)47

Digestibility and beef performance after feeding NaOH- or KOH-treated pig wastes 17 (1987)235

Evaluation of cardboard and dried poultry waste as feed ingredients for ruminants 18 (1987)209

Animal excreta as ruminant feed – scope and limitations under Indian conditions 19 (1988)1

Effect of the nature of dietary fibre on transit time and faecal excretion in the growing pig 20(1988)327

Nutritive value of rice straw (*Oryza sativa*) ensiled with animal excreta and rumen digesta 24(1989)69

Mastiction
The effect of method of preservation of forages on their resistance to reducion in particle size with an artificial masticator 24 (1989)185

Maturation
Whole-crop peas. I. Changes in botanical and chemical composition and rumen in vitro degradability during maturation 17(1987) 15

Meat and bone meal
Protein quality of mixtures of animal protein meals 9(1983)79

The digestion by sheep of barley and maize-based diets containing either meat and bone meal or soya bean meal 15(1986)301

Meat meal
Meat meal in the diet of the early-weaned pig. I. A comparison of meat meal and soya bean meal 4(1979)33

Meat meal in the diet of the early-weaned pig. II. Amino acid supplementation 4(1979)43

Meat meal in the diet of the early-weaned pig. III. Meat meal quality and the processing of meat meals 4(1979)53

Meat meal in the diet of the early-weaned pig. IV. The supplementation of diets with tryptophane, lysine and methionine 7 (1982)27

Meat and bone and fish meals in balancer feeds for choice-fed broilers 11(1984)221

Meat quality
Effect of avoparcin on performance, body composition and meat quality of finishing bulls 8(1983)237

Menadione
Comparison of the relative biological availability and stability of commercial sources of vitamin K activity 3(1978)329

Mercury
Nutritional value and mercury content of fish silage for growing pigs 9(1983)169

Metabolic faecal output
Digestibility and digesta retention in dairy cows receiving hay or silage at varying concentrate levels 11(1984)279

Methane digester effluent
Nitrogen balance and abomasal protein-nitrogen flow in growing ruminants fed methane digester effluent in combination with distillers dried grains 13(1985)47

Methionine
Adjustments of the protein level for slaughter pigs and supplementation with lysine and methionine 1(1976)619

Meat meal in the diet of the early-weaned pig. II. Amino acid supplementation 4(1979)43

Limiting amino acids in dried microbial cells given to young turkeys 4(1979)209

Meat meal in the diet of the early-weaned pig. IV. The supplementation of diets with tryptophane, lysine and methionine 7 (1982)27

Utilisation of L- and DL-amino acids by broiler chicks fed diets containing Pruteen 17(1987)1

Methylene-*bis*-propionate
Preservation of high-moisture maize – a comparison of gaseous and liquid anhydrous ammonia with methylene-*bis*-propionate 5 (1980)337

Micro-ingredients
The effect of dilution levels in premixes on micro-ingredient dispersion in animal feeds 8(1983)139

Microbial protein synthesis
Effect of dietary nitrogen source on microbial protein synthesis, dietary protein degradation and nutrient digestion in steers 14 (1986)117

Relative intakes, digestibility and bacterial protein synthesis by sheep and cattle fed high-roughage diets 16(1986/1987)75

Micronization
Effects of micronization on the digestibility of whole soya beans and rapeseeds for the growing pig 3(1978)179

Milk substitutes
The replacement of skim milk by dried microbial cells and whey in milk diets for calves 2(1977)85

Digestibility of protein and amino acids of a fermentation single-cell protein for veal calves 2(1977)377

Effect of dry-matter concentration of milk substitute and method of feeding on intake and performance by lambs 5(1980)321

Use of fish protein hydrolysate in milk replacers 7(1982)135

Performance of calves fed on milk replacers containing fish protein hydrolysate 7(1982) 141

The effects on feed intake, growth and digestibility of nutrients of including calcium soaps of fat in diets for young calves 14 (1986)103

A continuous process for the production of purified porcine γ-globulin for use in pig-milk replacer supplements 17(1987)213

Fish protein concentrates for pre-ruminant calves: effects of processing temperatures and source of fish on protein digestibility and biological value 18(1987)181

The effects of method of processing of starters, tallow inclusion and roughage supplementation on the performance of early-weaned calves 19(1988)231

An evaluation of skim milk powders subjected to different heating conditions during processing as dietary protein sources for the young pig 22(1988)203

Evaluation of pea protein concentrate as a source of protein in milk replacers for Holstein calves 24(1989)267

Milling
Effects of milling and pelleting of maize, barley and wheat on their metabolizable energy value for cockerels and chicks 9(1983) 99

The effect of milling and sodium hydroxide treatment on intake and digestibility of wheat straw by sheep 11(1984)149

Milling barley and lupin grain in diets for cattle 12(1984/1985)57

The effect of mechanical treatment on the nutritive value of high-moisture barley and the performance of growing-finishing pigs 24(1989)121

Mimosine
Growth performance of, and mimosine excretion by, young chicks fed on *Leucaena leucocephala* 7(1982)247

Investigations into the effects of processing on the retention of the carotenoid fractions of *Leucaena leucocephala* during storage, and the effects of processing on mimosine concentration 9(1983)307

Minerals, general
The use of whole barley diets fortified with solutions of urea, minerals and vitamins for lambs 2(1977)307

Soil chemistry and mineral problems in farm livestock. A review 5(1980)95

Lucerne juice as a protein supplement for growing pigs: effects of mineral content of the diet and of the water supply 6(1981)35

Effects of dietary maize solubles and minerals on albumen quality of fresh and stored brown-shelled eggs 8(1983)177

Mink

Slaughterhouse by-products preserved by *Lactobacillus plantarum* fermentation as feed for mink and foxes 20(1988)287

Modelling

Evaluation of mathematical models to describe neutral detergent residue in terms of its susceptibility to degradation in the rumen 15(1986)249

Sources of error in predicting digestible dry matter from the acid-detergent fiber content of forages 21(1988)205

Molasse

The nutritional value for sheep of molasses and condensed molasses solubles 2(1977) 143

Proximate chemical analysis of final cane molasses and effect of feeding 30% molasses on intestinal sucrase and maltase activities in the rat 3(1978)15

Studies on fermentation of sorghum silage during storage - effect of urea and molasses 3(1978)299

Distillery effluents as animal feed: the use of condensed beet molasses stillage (CBMS) in broiler feeding 5(1980)233

Methods of improving the utilization of cereal straw by ruminants. I. Supplements of urea, molasses and dried grass and treatment with sodium hydroxide 8(1983)221

Effect of wilting and the additives, straw, molasses and urea on the fermentation pattern of maize silage 9(1983)185

Effects of stage of maturity and addition of molasses on nutritive value of maize stover silage 12(1984/1985)65

Ensiling of leguminous green forages in combination with different dry roughages and molasses 12(1984/1985)133

Effects of molasses/urea and bacterial inoculant additives on silage quality, dry matter recovery, and feeding value for cattle 12 (1984/1985)205

Effect of sugarcane condensed molasses solubles, sugarcane molasses and monensin on performance and volatile fatty acid production in finishing steers 12(1984/1985)275

Differences in chemical composition and digestibility of beet and cane molasses 13 (1985)83

Methods of improving the uti'ization of cereal straw by ruminants. V. The effect of molasses and dried grass 13(1985)293

Effects of supplementing a wheat straw diet with urea or a urea–molasses block and/or cottonseed meal on intake and liveweight change of lambs 16(1986/1987)25

Sugar beet molasses for growing and fattening pigs 18(1987)131

Effect of sorghum straw alone or in combination with molasses and nitrogen sources on performance of Sudan desert lambs 19 (1988)351

A note on the use of sugar-beet pulp silage and molasses in the diet of lactating dairy cows 20(1988)13

The effects of barley, unmolassed sugar-beet pulp and molasses supplements on organic matter, nitrogen and fibre digestion in the rumen of cattle given a silage diet 20(1988) 259

Digestion of diets containing molassed or plain sugar-beet pulp by growing pigs 23 (1989)67

Utilization of sugar beet molasses and monensin for finishing dairy bullocks 25(1989) 11

Molybdenum

Reducing the potential copper toxity of concentrates to sheep by the use of molybdenum and sulphur supplements 2(1977)235

Monensin

Monensin-sodium as a performance-promoting additive for fattening bulls and its impact on carcass and meat quality characteristics 7(1982)401

Effect of sugarcane condensed molasses solubles, sugarcane molasses and monensin on performance and volatile fatty acid production in finishing steers 12(1984/1985)275

The effect of monensin on rumen fermentation in sheep fed on all-roughage and barley/roughage diets 15(1986)7

Utilization of sugar beet molasses and monensin for finishing dairy bullocks 25(1989) 11

Monocarbonyl compounds

Thermofractographic determination of monocarbonyl compounds in animal waste fats used as feed fat 13(1985)227

Mycotoxins

In vitro metabolism of mycotoxins by bacterial, protozoal and ovine ruminal fluid preparations 25(1989)169

N

Napier grass
Cassava meal as supplement to napier grass diets for growing sheep and goats 18(1987) 197

Navy bean
The dye absorption test as an indicator of the nutritive quality of heat-processed navy bean 9(1983)71

Neutral detergent
Effect of straw treatment and laboratory of incubation on degradation of neutral detergent residue in the rumen 18(1987)75

Neutral detergent fibre
The study of several modifications of the neutral detergent fibre procedure 9(1983)19

Neutral detergent residue
Evaluation of mathematical models to describe neutral detergent residue in terms of its susceptibility to degradation in the rumen 15(1986)249

Nitrogen
Effect on digestibility and nitrogen content of barley straw of different ammonia treatments 2(1977)131

Abomasal and plasma nitrogen constituents of steers fed on different physical forms of shelled maize 2(1977)225

Nitrogen levels in low-roughage diets for efficient rumen microbial protein production 2(1977)255

Nitrogen metabolism in broiler chickens consuming the bacterial strain *Alcaligenes eutrophus* 5(1980)241

Evaluation of corn gluten meal with urea as a source of supplementary nitrogen for growing calves and lambs 7(1982)375

The effects of sulphur supplementation on cellulose digestion in vitro and on nutrient digestion, nitrogen metabolism and rumen characteristics of lambs fed on good quality fescue and tropical star grass hays 8(1983) 129

The effect of prolonged storage on the digestibility and nitrogen content of ammonia-treated barley straw 8(1983)147

The effect of some pre-ensiling treatments on silage composition and nitrogen disappearance in the rumen 8(1983)259

Energy and nitrogen balance, intake and growth rate of lambs fed on unwilted grass silages treated with a formaldehyde–acid mixture or untreated 9(1983)107

Note on high-nitrate para grass (*Brachiaria mutica*) silage given to rabbits 9(1983)325

Digestion of organic matter and nitrogen in ammoniated barley straw 10(1983/1984)65

The effect of nitrogen supplements to maize silage on the amount and composition of the nitrogen fraction entering the intestine of cattle 13(1985)25

Nitrogen balance and abomasal protein-nitrogen flow in growing ruminants fed methane digester effluent in combination with distillers dried grains 13(1985)47

Effect of dietary nitrogen source on microbial protein synthesis, dietary protein degradation and nutrient digestion in steers 14 (1986)117

Effects of isopropyl alcohol treatment of soya bean meal on nitrogen utilization by ruminants 14(1986)127

Fractions of nitrogen, sulphur, phosphorus, calcium and magnesium in the herbage of perennial ryegrass as influenced by fertilizer nitrogen 14(1986)231

Effects of processing time and moisture content on amino acid composition and nitrogen characteristics of feather meal 14(1986) 279

Estimation of optimum supplements of essential amino acids to a wheat and barley-based diet in rats using a nitrogen balance technique 15(1986)21

The effect of β-glucanase on the utilization of starch and nitrogen by broiler chickens fed on barley of low- or high-viscosity 15(1986) 83

Digestibility of dry matter, nitrogen and energy by dairy cows fed whole or hammer-milled lupin grain in oaten hay or oaten pasture based diets 16(1986/1987)143

The effect of enzymatic treatment on amino acid content and nitrogen characteristics of feather meal 16(1986/1987)151

The apparent and true ileal digestibility of nitrogen and amino acids in lactic casein for the growing pig 16(1986/1987)169

The effects of different sources of nitrogen supplementation on the digestion of fibre components in the rumen of steers 17(1987) 65

Dietary fibre and nitrogen excretion and retention by pigs 19(1988)185

Effect of sorghum straw alone or in combination with molasses and nitrogen sources on performance of Sudan desert lambs 19 (1988)351

Complexes of urea and formaldehyde as non-protein nitrogen compounds in ruminant rations: a review 20(1988)1

The effects of barley, unmolassed sugar-beet pulp and molasses supplements on organic matter, nitrogen and fibre digestion in the rumen of cattle given a silage diet 20(1988) 259

Rumen degradability of nitrogen in typical cultivated grasses and leguminous fodders 22(1988)79

Comparison of cottonseed cake and dried blood as nitrogen sources for growing Sudan Nubian goats 22(1988)333

Distribution of liquid supplement injected into large bales of roughage and comparison of nitrogen supplements for stocker cattle fed low-quality roughages 24(1989)13

Oven and stack ammoniation of grass hays. 2. Voluntary intake, digestibility and faecal nitrogen excretion in sheep in relation to hay composition 24(1989)313

Effect of blood meal processing method on performance and nitrogen utilization by growing cattle 24(1989)337

Noug meal
The intake and digestibility by sheep of oat straw and maize stover offered with different levels of noug (*Guizotia abyssinica*) meal 16(1986/1987)99

Nutritive value
Genetic variation in the feeding value of cereal straw 21(1988)127

Nutritive value–age influence
Influence of age on the nutrient utilization of diets for broilers 6(1981)405

Nutritive value–almond
Nutritive value of almond hulls for sheep 8 (1983)63

Nutritive value–apple
Nutritive value of apple pomace for ruminants 16(1986/1987)311

Nutritive value–arhar
Nutritional evaluation of gram (*Cicer arietinum*) and arhar (*Cajanus cajan*) chunies for cattle 5(1980)315

Nutritive value–bagaruwa
Biochemical evaluation of bagaruwa seeds (*Acacia nilotica*) for use as livestock feed 11(1984)45

Nutritive value–banana
Nutritive value of banana stalk (*Musa cavendishi*) as a feed for sheep 22(1988)327

Nutritive value–barley
Nutritional value of waxy barleys for broiler chickens 8(1983)25

Feeding value of hydroponically sprouted barley for poultry and pigs 13(1985)183

Nutrient content of hydroponically sprouted barley 13(1985)191

The nutritive value of barley for monogastric animals and for ruminants as affected by artificial drying and alkali and anaerobic storage 15(1986)285

Nutritional value for chickens of dried distillers-spent-grain from barley and dehulled barley 17(1987)145

The nutritive value of hulless barley for swine 19(1988)191

The effect of mechanical treatment on the nutritive value of high-moisture barley and the performance of growing-finishing pigs 24(1989)121

Effect of enzyme supplementation on the nutritive value of hulless barley in chicken diets 24(1989)233

Nutritive value–bean, winged
The proximate composition and nutritive value of the winged bean *Psophocarpus tetragonolobus* (L.) DC for broilers 11 (1984)231

Nutritive value–brewer's dried grains
Evaluation of brewer's dried grains in the diets of laying hens 6(1981)169

Evaluation of brewer's dried grains in the diets of broiler chickens 6(1981)179

Nutritive value–buckwheat
A nutritional evaluation of buckwheat (*Fagopyrum esculentum*) 3(1978)95

Nutritive value–cabbage
Cabbage (*Brassica oleracea*) in the diet of growing–finishing pigs 5(1980)69

Nutritive value–canola meal
The nutritive value of canola meal for early-weaned pigs 18(1987)45

Nutritive value–Canola rapeseed
The influence of steam pelleting and grinding on the nutritive value of Canola rapeseed for poultry 8(1983)303

Nutritive value–cereals, general
Nutritional evaluation, using rats, of diets based on cereals and ethanol-grown yeast 6(1981)309

Nutritive value–cocoa
Studies on the feeding value of agro-industrial by-products. III. Digestibility of cocoa-pod and cocoa-pod-based diets by ruminants 13 (1985)249

Studies on the feeding value of agro-industrial by-products: strategies for improving the utilisation of cocoa-pod-based diets by ruminants 20(1988)189

Nutritive value–coffee pulp
A note on the nutritive value of dried instant coffee residue for broiler chickens and turkey poults 10(1983/1984)285

Nutritive value–cottonseed
A review of the nutritive value and utilization of whole cottonseed, cottonseed meal and associated by-products by dairy cattle 18 (1987)89

Nutritive value–fats
Investigations on the nutritional value for ruminants of various fats 19(1988)111

The nutritional value for broilers of fats characterized by short-chain fatty acids as affected by level of inclusion and age 22(1988) 105

Nutritive value–fibre
The digestibility and nutritive value for dairy heifers of rations containing cellulose fibre 2(1977)287

Nutritive value–fish silage
Examination of stability, and its effect on the nutritive value, of fish silage in diets for growing pigs 21(1988)43

Nutritive value–forages, general
A new technique for upgrading the nutritive value of roughages under field conditions 3 (1978)201

Principles governing the use of chemical methods for assessing the nutritive value of forages: a review 3(1978)265

Predicting the energetic value of forage from digestibility of organic matter in vitro or solubility in potassium hydroxide 6(1981) 235

Different sources of ammonia for improving the nutritive value of low quality roughages 6(1981)377

Chemical properties of fibre in relation to nutritive quality of ammonia-treated forages 10(1983/1984)155

Drying treatment drastically affects feed evaluation and feed quality with certain tropical forage species 22(1988)69

The nutritive value of ensiled forages cut at three stages of growth 22(1988)237

Nutritive value–fungal protein
Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. I. Composition and nutritive evaluation 22(1988)1

Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. III. Nutritive value for chicks and piglets 22(1988)23

Nutritive value–gram
Nutritional evaluation of gram (*Cicer arietinum*) and arhar (*Cajanus cajan*) chunies for cattle 5(1980)315

Nutritive value–groundnut
Feeding value of sunflower and groundnut cakes for broilers 4(1979)143

Feeding value of sunflower- and groundnut-cakes for laying hens 6(1981)63

Nutritive value–hominy
Feeding and nutritive value of hominy and corn grits for poultry 19(1988)313

Nutritive value–kapok seed
The effect of nutrient supplementation on the feeding value of kapok seed for poultry 9 (1983)263

Nutritive value–leaf protein
Nutritive value of leaf protein coagulum and cereals mixtures 4(1979)91

Nutritive value–leaf protein concentrate
Nutritive value of lupin-seed protein and leaf-protein concentrates supplemented with various sulphur sources 6(1981)189

Nutritive value of leaf-protein concentrate from potato haulm for rats and chicks 6 (1981)413

The effect of toxic substances present in some leaf protein concentrates on their nutritive value 24(1989)151

Nutritive value–*Leucaena leucocephala*
A note on the nutrient and mineral composition of *Leucaena leucocephala* in Western Nigeria 4(1979)221

Nutritive value–lupin seed
Nutritive value of lupin-seed protein and leaf-protein concentrates supplemented with various sulphur sources 6(1981)189

Nutritive value–maize
Factors influencing the composition and nutritive value of ensiled whole-crop maize 1 (1976)381

Nutritive value of dehydrated whole-crop maize 1(1976)429

The nutritive value (energy) of maize silage 1 (1976)481

Effect of level of sodium hydroxide treatment and volume of solution on the nutritive value of maize cobs 4(1979)1

Nutritive value of thermoammoniated and steam-treated maize stover I. Intake, digestibility and nitrogen retention 4(1979) 177

Nutritive value of thermoammoniated and steam-treated maize stover II. Rumen metabolites and rate of passage 4(1979)187

Chemical composition, digestibility and feeding value of maize cobs 9(1983)121

Influence of harvest date on the nutritive value and ensiling characteristics of maize stover 14(1986)11

Effects of moisture level, treatment time and soya bean addition on the nutritive value of urea-treated maize stover 19(1988)67

Feeding value of ensiled high-moisture maize grain with cob for growing-finishing pigs 20 (1988)145

Nutritive value–manure

Fermentation characteristics and feeding value of ensiled poultry litter containing wheat straw, bagasse or sawdust 7(1982)341

Nutritive value of ensiled broiler litter for cattle 8(1983)19

Nutritive value–molasse

The nutritional value for sheep of molasses and condensed molasses solubles 2(1977) 143

Nutritive value–navy bean

The dye absorption test as an indicator of the nutritive quality of heat-processed navy bean 9(1983)71

Nutritive value–oat

The effects of cereal particle size and pelleting on the nutritive value of oat-based diets for the growing pig 8(1983)91

Nutritive value of irradiated and β-glucanase-treated wild oat groats (*Avena fatua* L.) for broiler chickens 16(1986/1987)243

Nutritive value of naked oats for ruminants 18(1987)83

Nutritive value–olive pulp

Nutritional value for rabbits of olive pulp and the effects on their visceral organs 25(1989) 79

Nutritive value–para-rubber seed

Chemical composition and nutritional value of para-rubber seed and its products for chickens 10(1983/1984)257

Nutritive value–pea

Yield, composition and nutritive value of a leaf protein concentrate obtained by fractionating the shelled aerial part of *Pisum savitum* L. 8(1983)155

Nutritional and growth studies with pea-crop meals and peas for growing–finishing pigs 16(1986/1987)119

The nutritive value of white- and dark-flowered cultivars of pea for growing-finishing pigs 17(1987)245

Dry matter yield and chemical composition of pigeon pea (*C. cajan*) leaf meal and the nutritive value of pigeon pea leaf meal and grain meal for laying hens 24(1989)111

Nutritive value–peanut

Effect of sodium hydroxide treatment and/or extrusion cooking on the nutritive value of peanut hulls 12(1984/1985)187

Nutritive value–pigeon pea

Nutritional value of pigeonpea (*Cajanus cajan*) meal in poultry diets 25(1989)123

Nutritive value–potato

Liquid potato feed for finishing pigs: feeding value, inclusion rate and storage properties 15(1986)129

Nutritive value–proteins and amino acids

Composition and nutritive value of single-cell protein (SCP) 1(1976)9

Nutritive value–pungam cake

Nutritive value of pungam (*Pongamia glabra* Vent) cake for sheep 22(1988)321

Nutritive value–rapeseed

The nutritive value of tower rapeseed meal for swine 3(1978)235

Influence of steam pelleting on the feeding value of tower and candle rapeseed products for broiler chickens 3(1978)251

Nutritive value–rubber seed

Nutritive value of Nigerian rubber seed (*Hevea brasiliensis*) meal and oil. III. Performance characteristics, relative organ weights, hematocrit and plasma metabolites of growing female rats fed corn diets containing rubber seed meal, soya bean meal or casein 20(1988)125

Nutritive value–rumen contents

Nutritive value of rumen contents for monogastric animals 2(1977)351

Nutritive value–rye grain

Feeding value of alkali-treated whole rye (*Secale cereale* L.) grain for lactating cows and its digestibility for sheep 10(1983/1984)77

Nutritive value–*Sesbania sesban*
Yield, composition and feeding value of irrigated *Sesbania sesban* var. *Nubia* leaves harvested at latitude 38°N during a Mediterranean summer 18(1987)247

Nutritive value–silage
A calorimetric investigation of the nutritive value of formic acid- and formaldehyde–formic acid-treated grass silages 6(1981)245

The effect of silage additives containing formaldehyde on the fermentation of ryegrass ensiled at different dry matter levels and on the nutritive value of direct-cut silage 7(1982)303

Nutritional value and mercury content of fish silage for growing pigs 9(1983)169

Chemical composition and animal performance of grass forage of varying maturity stored as hay or silage 11(1984)35

Nutritive value–silage effluent
The use of effluent from grass silage in the diet of finishing pigs. II. Assessment of nutritive value of fresh and stored effluent 4(1979) 275

Nutritive value–single-cell protein
Nutritive value of single-cell protein produced by *Chaetomium cellulolyticum* grown on maize stover and pulp-mill sludge 11(1984) 211

Nutritive value–sorghum
Comparison of the nutritional value of low and medium tannin sorghum grains for pigs 8(1983)85

Nutritive value–soya bean
Nutrient and trypsin inhibitor content of hydroponically sprouted soya beans 13(1985) 203

Nutritive value–spruce
Nutritive value of spruce muka (foliage) for the growing chick 3(1978)63

Nutritive value–straw
Ensiled alkali-treated straw. II. The nutritive value for young beef cattle of mixtures of ensiled or frozen alkali-treated straw and ryegrass silage 3(1978)133

Akali treatment of straw for ruminants. II. Nutritive value of straw ensiled after alkali treatment. 3(1978)289

The effect of ammoniation on the intake and nutritive value of straw 4(1979)169

The effect of urease enzyme on treatment time and the nutritive value of straw treated with ammonia as urea 8(1983)271

The value of urea-treated straw in diets of lactating goats 11(1984)67

Improving the intake and nutritive value of wheat straw for beef cows 12(1984/1985) 125

A comparison of alkali treatment methods to improve the nutritive value of straw. I. Digestibility and metabolizability 12(1984/ 1985)295

Nutritive evaluation of uromol–wheat straw as a basal ration for buffaloes 13(1985)103

Chemical preparation of NaOH–keratin hydrolysate for improving the nutritive value of wheat straw 13(1985)215

Urea as a source of ammonia for improving the nutritive value of wheat straw 14(1986) 67

A comparison of alkali treatment methods used to improve the nutritive value of straw. II. In sacco and in vitro degradation relative to in vivo digestibility 14(1986)215

Evaluation of the increase in feeding value of barley straw following mild treatment with sulphur dioxide 15(1986)13

Yield and potential feeding value of straws of grain sorghum cultivars developed in Nigeria 17(1987)285

The effect of ammoniation on the nutritive value of wheat, barley and oat straws. I. Changes in chemical composition in relation to digestibility in vitro and cell wall degradability 19(1988)159

The effect of ammoniation on the nutritive value of wheat, barley and oat straws. II. Digestibility and energy value measurements in vivo and their prediction from laboratory measurements 19(1988)173

The effect of variation in the water content of wheat straw before ammoniation on its nutritive value for beef cattle 20(1988)111

The feeding value of ammoniated flax straw, wheat straw and wheat chaff for beef cattle 21(1988)57

Nutritive value of rice straw (*Oryza sativa*) ensiled with animal excreta and rumen digesta 24(1989)69

Nutritive value–sugar cane
Nutritive value for ruminants of sugar cane bagasse ensiled after spray treatment with different levels of NaOH 9(1983)1

Nutritive value–sunflower
Feeding value of sunflower and groundnut cakes for broilers 4(1979)143

Feeding value of sunflower- and groundnut-cakes for laying hens 6(1981)63

Nutritive value–tamarind
The nutritive value of tamarind seeds for broiler chicks 22(1988)285

Nutritive value–triticale
A nutritional evaluation of triticale with pigs 9(1983)49

Nutritive value of three cultivars of triticale for sheep, pigs and poultry 18(1987)19

Nutritional value of triticale (Cariman) for broiler diets 22(1988)119

Nutritive value of diets containing triticale and varying mixtures of triticale and maize for growing-finishing swine 22(1988)217

Nutritive value–vinasse
Protein and energy value of vinasse for pigs 5 (1980)221

Nutritive value–wheat
Observations on the chemical and nutritive content of white winter and spring wheats contaminated with deoxynivalenol (vomitoxin) 11(1984)293

Chemical composition and nutritive value of Canadian winter wheats (*Triticum aestivum* L.) given to laboratory rats 12(1984/1985)75

Feeding value of sprouted wheat (*Triticum aestivum*) for beef cattle finishing diets 15 (1986)113

Nutritive value–wheat residue
The nutritional value of crop residue components from several wheat cultivars grown at different fertilizer levels 11(1984)301

Nutritive value–whey
The nutritive value of liquid whey, either sour or sweet, when given in restricted amounts to the growing pig 3(1978)163

Nutritive value–yeast
Nutritional evaluation, using rats, of diets based on cereals and ethanol-grown yeast 6(1981)309

O

Oak browse
Composition of kermes oak browse as affected by shade and stage of maturity 21(1988)1

Oat grain
The effects of cereal particle size and pelleting on the nutritive value of oat-based diets for the growing pig 8(1983)91

Oat groats
The influence of oat groats and dietary level of lysine on the laying performance of crossbred hens 17(1987)271

Oat straw
Studies on ammonia-treated straw. I. The effects of type and level of ammonia, moisture content and treatment time on the digestibility in vitro and enzyme soluble organic matter of oat straw 7(1982)45

Botanical composition and extent of lignification affecting digestibility of wheat and oat straw and paspalum hay 7(1982)71

The effect of calcium hydroxide treatment on the digestibility of chopped oat straw supplemented with a solution containing urea, calcium, phosphorus, sodium, trace elements and vitamins 7(1982)93

The intake and digestibility by sheep of oat straw and maize stover offered with different levels of noug (*Guizotia abyssinica*) meal 16(1986/1987)99

The effect of ammoniation on the nutritive value of wheat, barley and oat straws. I. Changes in chemical composition in relation to digestibility in vitro and cell wall degradability 19(1988)159

The effect of ammoniation on the nutritive value of wheat, barley and oat straws. II. Digestibility and energy value measurements in vivo and their prediction from laboratory measurements 19(1988)173

Oats
Utilization of alkali-treated grain. 1. Alkali-treated grain in complete diets for steers and lambs 6(1981)273

The feeding value for ruminants of straw and whole-crop barley and oats treated with anhydrous or aqueous ammonia or urea 8 (1983)247

Anaerobically stored high moisture cereals for growing pigs 12(1984/1985)159

Digestibility of dry matter, nitrogen and energy by dairy cows fed whole or hammer-milled lupin grain in oaten hay or oaten pasture based diets 16(1986/1987)143

Nutritive value of irradiated and β-glucanase-treated wild oat groats (*Avena fatua* L.) for broiler chickens 16(1986/1987)243

Nutritive value of naked oats for ruminants 18(1987)83

The effect of β-glucanase supplementation on nutrient digestibility and growth in broilers given diets containing barley, oat groats or wheat 25(1989)193

Oil palm slurry
Oil palm slurry (OPS) as a partial replacement for maize in the diets of broiler chickens 17(1987)157

Olaquindox
Olaquindox as performance-promoting feed additive for growing pigs 4(1979)117

Olive cake
Effects of feeding alkali-treated olive cake on intake, digestibility and rumen liquor parameters 14(1986)139

Olive pulp
Nutritional value for rabbits of olive pulp and the effects on their visceral organs 25(1989) 79

Orange
Effect of urea on the ensiling process of orange pulp 12(1984/1985)233

The effect of treatment with urea, sorbic acid, or dehydration on orange peel silage 20 (1988)335

Organic matter, general
Comparison of laboratory methods for predicting the organic matter digestibility of forages 2(1977)337

Cellulase insoluble fibre as a measure of unavailable organic matter in cattle compounds containing alkali-treated straw 3 (1978)227

Ozone
The response of screened manure fibre to ozone and sodium hydroxide treatments 15 (1986)47

Effects of ozonation on the chemical composition and in vitro degradation of triticale internodes and red clover stems 24(1989) 159

P

Palm kernel cake
Rumen fermentation, urease activity and performance of cattle given palm kernel cake-based diet 20(1988)79

Palm kernel meal
Palm kernel meal as a feed for poultry. 1. Composition of palm kernel meal and availability of its amino acids to chicks 16(1986/ 1987)179

Palm kernel meal as a feed for poultry. 2. Diets containing palm kernel meal for starter and grower pullets 16(1986/1987)187

Palm kernel meal as a feed for poultry. 3. Replacement of groundnut cake by palm kernel meal in broiler diets 16(1986/1987) 195

Palm kernel meal as a feed for poultry. 4. Use of palm kernel meal by laying birds 20 (1988)279

Palm oil mill effluent
The intake and digestibility in goats of *Leucaena leucocephala* supplemented with dehydrated palm oil mill effluent 24(1989)45

Palmitic acid
Feed intake and milk yield at increasing supplement of a palmitic and stearic acid-rich type of fat in comparison with animal fat 22(1988)179

Paper diets
Intake, digestibility and energy utilization of corrugated paper diets 17(1987)57

Para grass
Note on high-nitrate para grass (*Brachiaria mutica*) silage given to rabbits 9(1983)325

Para-rubber seed
The influence of processing and storage on hydrogen cyanide and tannin contents of para-rubber seed and its products 9(1983) 319

Chemical composition and nutritional value of para-rubber seed and its products for chickens 10(1983/1984)257

Particle size
The effect of method of preservation of forages on their resistance to reducion in particle size with an artificial masticator 24 (1989)185

Particle size distribution
Particle size distribution of a feed premix containing amprolium 3(1978)211

The effects of cereal particle size and pelleting on the nutritive value of oat-based diets for the growing pig 8(1983)91

Pasture plants
The relationship between the intake of herbage grazed by dairy cows and its digestibility 1 (1976)53

Pea
Yield, composition and nutritive value of a leaf protein concentrate obtained by fractionating the shelled aerial part of *Pisum savitum* L. 8(1983)155

Effects of ensiling or drying on five forage species in western Kenya: *Zea mays* (maize stover), *Pennisetum purpureum* (Pakistan

napier grass), *Pennisetum* sp. (bana grass), *Impomea batata* (sweet potato vines) and *Cajanus cajan* (pigeon pea leaves) 13(1985) 1

Fermentation of hammermilled barley, lupin, pea and faba bean grain in the rumen of dairy cows 16(1986/1987)261

The chemical composition of pea plant parts as related to harvesting time 16(1986/1987) 305

Whole-crop peas. I. Changes in botanical and chemical composition and rumen in vitro degradability during maturation 17(1987) 15

Whole-crop peas. II. Digestion of early- and late-harvested crops in the gastrointestinal tract of pigs 17(1987)33

The nutritive value of white- and dark-flowered cultivars of pea for growing-finishing pigs 17(1987)245

Utilization of white-flowered smooth-seeded spring peas (*Pisum sativum hortense*, CV Amino) by the breeding sow 22(1988)91

Pea-crop meals

Nutritional and growth studies with pea-crop meals and peas for growing–finishing pigs 16(1986/1987)119

Peanut hulls

Effect of sodium hydroxide treatment and/or extrusion cooking on the nutritive value of peanut hulls 12(1984/1985)187

Pectin

Determination of soluble fiber components: (1-<3; 1-<4)-β-D-glucans and pectins 23 (1989)15

The effects of starch and pectin-rich diets on quantitative aspects of digestion in sheep 24(1989)289

'Pekilo protein'

The value of 'Pekilo protein' for growing pigs 2(1977)161

Pellet quality

Effect of pellet size and pellet quality on pig performance 10(1983/1984)277

Pelleting

Influence of steam pelleting on the feeding value of tower and candle rapeseed products for broiler chickens 3(1978)251

The effects of cereal particle size and pelleting on the nutritive value of oat-based diets for the growing pig 8(1983)91

The influence of steam pelleting and grinding on the nutritive value of Canola rapeseed for poultry 8(1983)303

Effects of milling and pelleting of maize, barley and wheat on their metabolizable energy value for cockerels and chicks 9(1983) 99

Effects of pelleting, added sodium bentonite and fat in a wheat-based diet on performance and carcass characteristics of small white turkeys 12(1984/1985)223

The effect of pelleting on the utilisation of yeast grown on n-alkanes by broiler chicks 12(1984/1985)285

Effect of ammonia treatment and processing of whole cotton plants as sole source of roughage in complete feeds for growing cross-bred calves 13(1985)93

The functional properties of feed raw materials and their effect on the production and quality of feed pellets 18(1987)1

Pentosanase

Effects of enzyme supplementation of diets based on wheat, rye or triticale on their productive value for broiler chickens 20 (1988)313

Pepsin

Prediction of the digestibility of ruminant feeds from their solubility in enzyme solutions 2(1977)153

Prediction of the dry matter digestibility of tropical grasses using a pepsin-cellulase assay 2(1977)247

Comparison of forage digestibility in vitro with enzymic solubility 11(1984)125

A comparison of laboratory methods involving single or two-stage cellulase and two-stage rumen inoculum–pepsin treatment for predicting digestibility and intake of hays in sheep 12(1984/1985)101

The use of an enzymatic technique to predict digestibility, metabolizable and net energy of compound feedstuffs for ruminants 14 (1986)203

Pepsin–cellulase technique

The accuracy of the pepsin–cellulase technique for estimating the dry matter digestibility in vivo of grasses and legumes 3(1978)277

Pharmacology

Long term feeding of concentrate fumigated with carbon tetrachloride to sheep 2(1977) 77

A note on the effect on the aerobic deterioration of maize silage of the manipulation of the microflora by means of antibiotics 3 (1978)89

Particle size distribution of a feed premix containing amprolium 3(1978)211

Efficacy of virginiamycin and a commercially-available lactobacillus probiotic in swine diets 8(1983)69

Phosphorus

Solubility and flow of calcium, magnesium and phosphorus in the digestive tract of sheep given maize or alfalfa silages 9(1983) 131

Biological availability of phosphorus from basic steel slag 11(1984)75

The relative availability of phosphorus in inorganic feed phosphates for young turkeys and pigs 13(1985)117

Fractions of nitrogen, sulphur, phosphorus, calcium and magnesium in the herbage of perennial ryegrass as influenced by fertilizer nitrogen 14(1986)231

Compositional quality of certain uncoventional calcium and phosphorus sources in India for use as mineral supplements for livestock 23(1989)343

Pig–finishing

The composition of processed abattoir blood and its use in finishing-pig diets 2(1977)21

Treatment of the haem fraction of abattoir blood with formic acid to produce a solid state, low dry matter protein supplement for use in finishing-pig diets 2(1977)367

The use of effluent from grass silage in the diet of finishing pigs. I. Variation in composition of effluents 4(1979)263

The use of effluent from grass silage in the diet of finishing pigs. II. Assessment of nutritive value of fresh and stored effluent 4(1979) 275

Liquid potato feed for finishing pigs: feeding value, inclusion rate and storage properties 15(1986)129

A comparison of wheat- or barley-based diets given ad libitum as meal or pellets to finishing pigs 22(1988)263

Pig–general

The influence of feeding level and level of dietary inclusion on the digestibility of barley meal in the pig 2(1977)41

The potato (*Solanum tuberosum*) as a source of nutrients for pigs, calves and fowl – a review 2(1977)171

Influence of pericarp and endosperm colour and type on digestibility of grain sorghum by pigs 2(1977)219

Relationships between tristimulus colour values and digestibility of grain sorghum by swine 2(1977)327

Use of unextracted sunflower seeds as a protein source 5(1980)51

The energy systems used in the EEC countries: their application to 24 commercial pig-feeds and relationships between them 6(1981)211

Prediction of the energy value of commercial pig feeds, according to the various energy systems used in the European Community, that are based on simple analytical parameters of the feed 6(1981)367

Energy evaluation of feeds and compounded diets for pigs – a review 7(1982)387

Efficacy of virginiamycin and a commercially-available lactobacillus probiotic in swine diets 8(1983)69

Comparison of bioavailable energy values of diets measured with poultry and swine 8 (1983)191

Effect of storage and processing of barley on its utilization by pigs and rats 9(1983)89

Low-glucosinolate rapeseed products for pigs – a review 9(1983)239

Digestibility and energy value of sweet lupin seed (*Lupinus albus* var. multolupa) in pigs 12(1984/1985)171

Waxy and normal grain sorghums with varying tannin contents in diets for young pigs 12(1984/1985)179

Feeding value of hydroponically sprouted barley for poultry and pigs 13(1985)183

Whole-crop peas. II. Digestion of early- and late-harvested crops in the gastrointestinal tract of pigs 17(1987)33

Nutritive value of three cultivars of triticale for sheep, pigs and poultry 18(1987)19

Secretory response of the pig gut to non-starch polysaccharides 23(1989)55

Characterization of the flora of the large bowel of pigs: a status report 23(1989)79

Approach to the energetic importance of fibre digestion in pigs. I. Importance of fermentation in the overall energy supply 23(1989) 141

Approach to the energetic importance of fibre digestion in pigs. II. An experimental approach to hindgut digestion 23(1989)169

Factors effecting the digestibility by pigs of energy and protein in wheat, barley and sorghum diets supplemented with canola meal 24(1989)253

Effect of graded levels of raw and processed rapeseed on feed digestibility and nutrient utilization in young pigs 24(1989)275

Pig–growing

Utilization of yeast grown on $_n$-paraffins in the feeding of heavy pigs, from weaning to slaughter 1(1976)25

Adjustments of the protein level for slaughter pigs and supplementation with lysine and methionine 1(1976)619

Lupin-seed meal (*Lupinus angustifolius* cv. Uniwhite) as a protein supplement to barley-based diets for growing pigs 1(1976) 631

Swedes (*Brassica napus*) for growing pigs: chemical composition and use as a replacement for barley in the diet 2(1977)31

A comparison between meals prepared from the seeds of different varieties of lupin as protein supplements to barley-based diets for growing pigs 2(1977)49

The alkaloid content of sweet lupin seed used in feeding trials on pigs and rats 2(1977)59

The value of 'Pekilo protein' for growing pigs 2(1977)161

The effect of supplementary protein on the performance of pigs given swedes (*Brassica napus*) as a replacement for barley in the diet 2(1977)277

The nutritive value of liquid whey, either sour or sweet, when given in restricted amounts to the growing pig 3(1978)163

Effects of micronization on the digestibility of whole soya beans and rapeseeds for the growing pig 3(1978)179

Blood meal as a source of protein for grower/finisher pigs 3(1978)191

Protein digestibility of hydrolyzed hog hair meal for pigs 3(1978)323

Waxy versus regular maize:energy value for growing pigs and utilization in piglet diets 4(1979)109

Olaquindox as performance-promoting feed additive for growing pigs 4(1979)117

Value of lucerne juice and grass juice as sources of protein for the growing pig 4 (1979)233

The effect of different patterns of thermal processing of potatoes on their digestibility by growing pigs 4(1979)295

Cabbage (*Brassica oleracea*) in the diet of growing–finishing pigs 5(1980)69

Digestibility of a fully hydrogenated tallow for growing pigs 5(1980)87

Value of freshly produced lucerne juice as a source of supplemental protein for the growing pig 5(1980)215

Protein and energy value of vinasse for pigs 5 (1980)221

Lucerne juice as a protein supplement for growing pigs: effects of mineral content of the diet and of the water supply 6(1981)35

Lupin-seed meal (*Lupinus albus* cv. Hamburg) as a source of protein for growing pigs 6(1981)285

Effects of different handling systems on pig feed quality 7(1982)11

Lysine requirements of imported European pigs in the humid lowland tropics from 23 to 62 kg liveweight 7(1982)233

Comparison of the nutritional value of low and medium tannin sorghum grains for pigs 8(1983)85

The effects of cereal particle size and pelleting on the nutritive value of oat-based diets for the growing pig 8(1983)91

Lysine requirement of weanling European pigs fed on yellow maize and groundnut cake diets in the humid lowland tropics 8 (1983)197

Cassava meal and cassava peel meal in diets for growing pigs 8(1983)211

A nutritional evaluation of triticale with pigs 9(1983)49

Nutritional value and mercury content of fish silage for growing pigs 9(1983)169

Effect of pellet size and pellet quality on pig performance 10(1983/1984)277

The effect of level and source of protein, fibre and fat in the diet on the energy value of compounded pig feeds 11(1984)11

The digestion of diets containing manioc by young growing pigs 12(1984/1985)119

Anaerobically stored high moisture cereals for growing pigs 12(1984/1985)159

The relative availability of phosphorus in inorganic feed phosphates for young turkeys and pigs 13(1985)117

Effect of dietary protein and fat levels on the utilization of the fat in sunflower seeds by the young pig 13(1985)159

Tryptophan content of feeds, limitations in diets and requirement for growing pigs 13 (1985)171

The evaluation of fresh sweet liquid whey as a protein supplement to maize for growing pigs 14(1986)171

Apparent ileal and faecal digestibility of amino acids in barley-based diets supplemented with soya bean meal or canola meal for growing pigs 14(1986)183

The absorption of dietary L-lysine and extruded L-lysine in pigs 15(1986)141

The utilisation by growing pigs of ruminant offal hydrolysed in formic acid 15(1986)273

Nutritional and growth studies with pea-crop meals and peas for growing–finishing pigs 16(1986/1987)119

The apparent and true ileal digestibility of nitrogen and amino acids in lactic casein for the growing pig 16(1986/1987)169

The prediction of the energy value of compounded pig foods from chemical analysis 17(1987)81

High-fat maize in diets for pigs and sows 17 (1987)201

The nutritive value of white- and dark-flowered cultivars of pea for growing-finishing pigs 17(1987)245

Canola meal as a protein supplement for growing–finishing pigs 18(1987)37

Sugar beet molasses for growing and fattening pigs 18(1987)131

Comparative bioavailable energy values of five ground cereal grains measured with growing rats and pigs 18(1987)143

Evaluation of Greek cottonseed meal. 2. Use in rations for fattening pigs 18(1987)303

A comparison of bioavailable energy values of ground cereal grains measured with adult cockerels and growing pigs 19(1988)105

Dietary fibre and nitrogen excretion and retention by pigs 19(1988)185

The nutritive value of hulless barley for swine 19(1988)191

Feeding value of ensiled high-moisture maize grain with cob for growing-finishing pigs 20 (1988)145

Effect of the nature of dietary fibre on transit time and faecal excretion in the growing pig 20(1988)327

Examination of stability, and its effect on the nutritive value, of fish silage in diets for growing pigs 21(1988)43

Copper and iodine in pig diets with high glucosinolate rapeseed meal. I. Performance and thyroid hormone status of growing pigs fed on a diet with rapeseed meal treated with copper sulphate solution or untreated and supplements of iodine, copper or a quinoxaline derivative 22(1988)33

Copper and iodine in pig diets with high glucosinolate rapeseed meal. II. Influence of iodine supplements for rations with rapeseed meal untreated or treated with copper ions on performance and thyroid hormone status of growing pigs 22(1988)45

Nutritive value of diets containing triticale and varying mixtures of triticale and maize for growing-finishing swine 22(1988)217

Digestion of diets containing molassed or plain sugar-beet pulp by growing pigs 23 (1989)67

The effect of mechanical treatment on the nutritive value of high-moisture barley and the performance of growing-finishing pigs 24(1989)121

Body composition and growth of young pigs as affected by protein source and a growth promoter 24(1989)219

Pig–sow

The nutritive value of tower rapeseed meal for swine 3(1978)235

Pruteen in the diet of breeding pigs: reproductive performance 6(1981)297

High-fat maize in diets for pigs and sows 17 (1987)201

Utilization of white-flowered smooth-seeded spring peas (*Pisum sativum hortense*, CV Amino) by the breeding sow 22(1988)91

Observations on the feeding of non-lactating sows on graded levels of liquid whey 24 (1989)327

Pig–starter

The effect of diet complexity and physical form on the performance of starter pigs and feed processing efficiency 8(1983)99

A continuous process for the production of purified porcine γ-globulin for use in pig-milk replacer supplements 17(1987)213

Pig–weaner

The effects of air dried poultry manure on performance, organ weights and haematological measurements of weaner pigs reared in the tropics 4(1979)23

Meat meal in the diet of the early-weaned pig. I. A comparison of meat meal and soya bean meal 4(1979)33

Meat meal in the diet of the early-weaned pig. II. Amino acid supplementation 4(1979)43

Meat meal in the diet of the early-weaned pig. III. Meat meal quality and the processing of meat meals 4(1979)53

Lupinus angustifolius cv. Unicrop as a protein- and energy-source for weaner pigs 6(1981) 27

Meat meal in the diet of the early-weaned pig. IV. The supplementation of diets with tryptophane, lysine and methionine 7 (1982)27

Effect of processing on the utilization by young pigs of the fat in soya beans and sunflower seeds 12(1984/1985)267

The nutritive value of canola meal for early-weaned pigs 18(1987)45

Evaluation of triticale given to pigs, poultry and rats 19(1988)359

Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. III. Nutritive value for chicks and piglets 22(1988)23

An evaluation of skim milk powders subjected to different heating conditions during processing as dietary protein sources for the young pig 22(1988)203

Alkali-treated soya flour for piglets 24(1989)1

Pigeon pea

Variation in metabolizable energy content of raw or autoclaved white and brown varieties of three tropical grain legumes 13 (1985)141

Dry matter yield and chemical composition of pigeon pea (*C. cajan*) leaf meal and the nutritive value of pigeon pea leaf meal and grain meal for laying hens 24(1989)111

Nutritional value of pigeonpea (*Cajanus cajan*) meal in poultry diets 25(1989)123

Pimaricin

A note on the effect on the aerobic deterioration of maize silage of the manipulation of the microflora by means of antibiotics 3 (1978)89

Plant age

Cassava for animal feeding: effect of variety and plant age on production of leaves and roots 11(1984)49

Effect of variety and plant age on the cyanide content of whole-root cassava chips and its reduction by sun-drying 11(1984)57

Plant variety

Cassava for animal feeding: effect of variety and plant age on production of leaves and roots 11(1984)49

Effect of variety and plant age on the cyanide content of whole-root cassava chips and its reduction by sun-drying 11(1984)57

Plantain sheath

In vitro studies and short-term feeding trial in lambs to evaluate plantain sheath (*Musa sapientum*) as a feed for ruminants 20 (1988)343

Polysaccharides

Secretory response of the pig gut to non-starch polysaccharides 23(1989)55

Fibre-degrading microorganisms in the monogastric digestive tract 23(1989)91

Assessing the importance of host-derived polysaccharides as carbon sources for bacteria growing in the human colon 23(1989) 109

Potassium hydroxide

Predicting the energetic value of forage from digestibility of organic matter in vitro or solubility in potassium hydroxide 6(1981) 235

Potato

The potato (*Solanum tuberosum*) as a source of nutrients for pigs, calves and fowl – a review 2(1977)171

The effect of different patterns of thermal processing of potatoes on their digestibility by growing pigs 4(1979)295

Nutritive value of leaf-protein concentrate from potato haulm for rats and chicks 6 (1981)413

Liquid potato feed for finishing pigs: feeding value, inclusion rate and storage properties 15(1986)129

Poultry–general

Nutritive value of three cultivars of triticale for sheep, pigs and poultry 18(1987)19

Preservatives

Laboratory application of preservatives to hay and the effects of irregular distribution on mould development 6(1981)73

Chemicals for preventing moulding in damp hay 6(1981)323

Laboratory tests of 8-quinolinol as an additive to volatile fatty acids for preserving hay and other feed-stuffs 7(1982)287

Problems of testing novel chemicals for the preservation of damp hay 8(1983)283

Losses of beta-carotene in unwilted forage crops during silage-making and feeding 9 (1983)63

Dicoumarol concentrations in sweetclover hay treated with preservatives and in spoiled hay of high- and low-coumarin cultivars of sweetclover 14(1986)221

Efficacy of sulfur dioxide as a grain preservative 19(1988)203

Pressure treatment

High temperature and pressure processing of maize cobs. II. Animal response and effect of sodium metabisulphite 14(1986)161

Proline

Amino acid composition of maize silage produced in the United Kingdom 4(1979)163

Propionic acid

Problems of testing novel chemicals for the preservation of damp hay 8(1983)283

Protein–pig

Canola meal as a protein supplement for growing–finishing pigs 18(1987)37

Protein-nitrogen flow

Nitrogen balance and abomasal protein-nitrogen flow in growing ruminants fed methane digester effluent in combination with distillers dried grains 13(1985)47

Proteins–cattle

Methanol-fermentation protein in veal calf nutrition 1(1976)33

Levels, sources and methods of nitrogen supplementation of maize silage for beef production 1(1976)465

Solubility of the protein of tropical pasture species and the rate of its digestion in the rumen 5(1980)183

Dry matter, protein, energy and fibre intake by dairy heifers grazing a Rhodes grass (*Chloris gayana*) pasture 6(1981)15

Rumen and post-rumen fermentation of spent tea leaf protein and other protein sources studied by the nylon bag method 7(1982) 221

The digestion in the small intestine of young bulls of the protein of rapeseed meal treated or untreated with formaldehyde 7 (1982)225

The use of kidney beans (*Phaseolus vulgaris*) as protein supplements in diets for young rapidly growing beef steers 12(1984/1985) 1

Effect of dietary nitrogen source on microbial protein synthesis, dietary protein degradation and nutrient digestion in steers 14 (1986)117

Utilization of whole rape (Canola) seed and sunflower seeds as sources of energy and protein in calf starter diets 15(1986)101

Fish silage as a protein supplement for early-weaned calves 17(1987)165

Fish protein concentrates for pre-ruminant calves: effects of processing temperatures and source of fish on protein digestibility and biological value 18(1987)181

Fresh and dried brewer's grains as protein supplements to barley straw diets given to pregnant beef cows 19(1988)33

Effects of ruminal degradability of the protein supplement on the utilization of maize-stover silage by growing beef cattle 20 (1988)241

Evaluation of pea protein concentrate as a source of protein in milk replacers for Holstein calves 24(1989)267

Effect of blood meal processing method on performance and nitrogen utilization by growing cattle 24(1989)337

The effect of a protein energy supplement on pasture protein and fibre digestion in the rumen of grazing steers 25(1989)39

Proteins–chicken

House fly larvae meal grown on municipal organic waste as a source of protein in poultry diets 4(1979)227

Rapeseed meal and its use in poultry diets. A review 5(1980)255

The utilization of torula yeast in diets for White Leghorn birds during growth and laying periods 7(1982)185

Proteins–fungal

Production of fungal protein from rasped fresh cassava roots using 200- and 3000-litre fermentors 8(1983)313

Proteins–general

The digestion and metabolism of the energy and protein of maize 1(1976)455

Further thermotolerant fungi for the conversion of cassava starch to protein 2(1977)7

Nitrogen levels in low-roughage diets for efficient rumen microbial protein production 2(1977)255

Inhibition of ruminal deamination in vitro by formaldehyde treatment of sunflower-seed, soya bean and fish meals: response curves to protective treatment 2(1977)267

Utilization of cattle manure for single-cell protein production with *Chaetomium cellulolyticum* 4(1979)199

Amino acid composition of two varieties of sorghum grain 5(1980)169

Nutritive value of lupin-seed protein and leaf-protein concentrates supplemented with various sulphur sources 6(1981)189

Protein quality of mixtures of animal protein meals 9(1983)79

Variability for protein and oil quality in *Lupinus albus* 9(1983)271

The quality of the green fraction of leaf protein concentrate from *Chenopodium quinoa* Willd. grown at different levels of fertilizer nitrogen 11(1984)239

In vitro effects of avoparcin on protein degradability and rumen fermentation 15 (1986)215

The effect of enzymatic treatment on amino acid content and nitrogen characteristics of feather meal 16(1986/1987)151

Effect of formaldehyde treatment on the degradation of acid-preserved fish silage protein in vitro 16(1986/1987)297

Yield and composition of protein concentrate, press cake, green juice and solubles concentrate from wet fractionation of *Sophora japonica* L. foliage 20(1988)177

Chemical and biological evaluation of Nigerian cottonseed meal protein quality 20 (1988)251

Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. I. Composition and nutritive evaluation 22(1988)1

Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. II. Effect of method of drying and response to amino acid supplementation 22(1988)13

Proteins–pig

Adjustments of the protein level for slaughter pigs and supplementation with lysine and methionine 1(1976)619

Lupin-seed meal (*Lupinus angustifolius* cv. Uniwhite) as a protein supplement to barley-based diets for growing pigs 1(1976) 631

The composition of processed abattoir blood and its use in finishing-pig diets 2(1977)21

A comparison between meals prepared from the seeds of different varieties of lupin as protein supplements to barley-based diets for growing pigs 2(1977)49

The value of 'Pekilo protein' for growing pigs 2(1977)161

The effect of supplementary protein on the performance of pigs given swedes (*Brassica napus*) as a replacement for barley in the diet 2(1977)277

Treatment of the haem fraction of abattoir blood with formic acid to produce a solid state, low dry matter protein supplement for use in finishing-pig diets 2(1977)367

Blood meal as a source of protein for grower/finisher pigs 3(1978)191

Protein digestibility of hydrolyzed hog hair meal for pigs 3(1978)323

Value of lucerne juice and grass juice as sources of protein for the growing pig 4 (1979)233

Use of unextracted sunflower seeds as a protein source 5(1980)51

Value of freshly produced lucerne juice as a source of supplemental protein for the growing pig 5(1980)215

Protein and energy value of vinasse for pigs 5 (1980)221

Lupinus angustifolius cv. Unicrop as a protein- and energy-source for weaner pigs 6(1981) 27

Lucerne juice as a protein supplement for growing pigs: effects of mineral content of the diet and of the water supply 6(1981)35

Lupin-seed meal (*Lupinus albus* cv. Hamburg) as a source of protein for growing pigs 6(1981)285

Pruteen in the diet of breeding pigs: reproductive performance 6(1981)297

The effect of level and source of protein, fibre and fat in the diet on the energy value of compounded pig feeds 11(1984)11

Effect of dietary protein and fat levels on the utilization of the fat in sunflower seeds by the young pig 13(1985)159

An evaluation of skim milk powders subjected to different heating conditions during processing as dietary protein sources for the young pig 22(1988)203

Body composition and growth of young pigs as affected by protein source and a growth promoter 24(1989)219

Proteins–rat

Protein quality studies on rats fed on cassava-root meal enriched with *Cephalosporium eichhorniae* 152 or with conventional plant-protein supplements 6(1981)197

Protein and energy utilisation in rats of diets based on lupin seed (*Lupinus albus* var. multolupa) 15(1986)33

Proteins–sheep

Effects of treatment of barley straw with anhydrous ammonia and supplementation with heat-treated protein meals on feed intake and liveweight performance of growing lambs 6(1981)145

Rumen degradation of protein in sunflower meal, rapeseed meal and lupin seed placed in nylon bags 11(1984)87

Value of mechanically extracted rape and safflower oilseed meals as protein supplements for growing lambs 11(1984)269

Degradation of crude protein and quality of undegradable protein in untreated or formaldehyde-treated rapeseed meal 12 (1984/1985)19

Proteins–turkey

Turkey feeding regimes with different ages of introduction of a choice of foods 4(1979)125

Protozoa

The role of ciliate protozoa and fungi in the rumen digestion of plant cell walls 10(1983/1984)121

Effect of absence of ciliate protoza on rumen fluid volume, flow rate and bacterial populations in sheep 10(1983/1984)145

Fibre-degrading microorganisms in the monogastric digestive tract 23(1989)91

In vitro metabolism of mycotoxins by bacterial, protozoal and ovine ruminal fluid preparations 25(1989)169

Pruteen
Digestibility of protein and amino acids of a fermentation single-cell protein for veal calves 2(1977)377

Pruteen in the diet of breeding pigs: reproductive performance 6(1981)297

Utilisation of L- and DL-amino acids by broiler chicks fed diets containing Pruteen 17(1987)1

Publishing, general
Titles of scientific papers 2(1977)1

Citation decay in animal science 2(1977)195

Pungam cake
Nutritive value of pungam (*Pongamia glabra* Vent) cake for sheep 22(1988)321

Q

Quinoxaline
Copper and iodine in pig diets with high glucosinolate rapeseed meal. I. Performance and thyroid hormone status of growing pigs fed on a diet with rapeseed meal treated with copper sulphate solution or untreated and supplements of iodine, copper or a quinoxaline derivative 22(1988)33

R

Rabbit
Citrus pulp in diets for fattening rabbits 5 (1980)23

Investigations of the treatment of sawdust for rabbit feeding. 1. Effect of sodium hydroxide treatment 6(1981)43

Investigations of the treatment of sawdust for rabbit feeding. 2. Effect of boiling and incubation with poultry manure 6(1981)51

Note on high-nitrate para grass (*Brachiaria mutica*) silage given to rabbits 9(1983)325

Studies on the nutritive value of roots, cabbage and grass silage for growing commercial rabbits 13(1985)299

Fat supplementation of diets for growing rabbits 16(1986/1987)109

Nutritional value for rabbits of olive pulp and the effects on their visceral organs 25(1989) 79

Rainfall
Estimating yield and nutrient losses due to rainfall on field-drying alfalfa hay 16(1986/1987)7

Rape seed
Utilization of whole rape (Canola) seed and sunflower seeds as sources of energy and protein in calf starter diets 15(1986)101

Rapeseed
Effects of micronization on the digestibility of whole soya beans and rapeseeds for the growing pig 3(1978)179

Influence of steam pelleting on the feeding value of tower and candle rapeseed products for broiler chickens 3(1978)251

Rapeseed meal and its use in poultry diets. A review 5(1980)255

Effect of graded levels of raw and processed rapeseed on feed digestibility and nutrient utilization in young pigs 24(1989)275

Rapeseed meal
The nutritive value of tower rapeseed meal for swine 3(1978)235

Rapeseed meal tannins and egg taint 6(1981) 421

The digestion in the small intestine of young bulls of the protein of rapeseed meal treated or untreated with formaldehyde 7 (1982)225

Low-glucosinolate rapeseed products for pigs – a review 9(1983)239

Rumen degradation of protein in sunflower meal, rapeseed meal and lupin seed placed in nylon bags 11(1984)87

Value of mechanically extracted rape and safflower oilseed meals as protein supplements for growing lambs 11(1984)269

Degradation of crude protein and quality of undegradable protein in untreated or formaldehyde-treated rapeseed meal 12 (1984/1985)19

Copper and iodine in pig diets with high glucosinolate rapeseed meal. I. Performance and thyroid hormone status of growing pigs fed on a diet with rapeseed meal treated with copper sulphate solution or untreated and supplements of iodine, copper or a quinoxaline derivative 22(1988)33

Copper and iodine in pig diets with high glucosinolate rapeseed meal. II. Influence of iodine supplements for rations with rapeseed meal untreated or treated with copper ions on performance and thyroid hormone status of growing pigs 22(1988)45

Rat
The alkaloid content of sweet lupin seed used in feeding trials on pigs and rats 2(1977)59

Proximate chemical analysis of final cane molasses and effect of feeding 30% molasses on

intestinal sucrase and maltase activities in the rat 3(1978)15

Protein quality studies on rats fed on cassava-root meal enriched with *Cephalosporium eichhorniae* 152 or with conventional plant-protein supplements 6(1981)197

Nutritional evaluation, using rats, of diets based on cereals and ethanol-grown yeast 6(1981)309

Nutritive value of leaf-protein concentrate from potato haulm for rats and chicks 6 (1981)413

Evaluation of optimum lysine and threonine supplements to a wheat and barley-based diet in rats 8(1983)163

Effect of storage and processing of barley on its utilization by pigs and rats 9(1983)89

Chemical composition and nutritive value of Canadian winter wheats (*Triticum aestivum* L.) given to laboratory rats 12(1984/1985)75

Estimation of optimum supplements of essential amino acids to a wheat and barley-based diet in rats using a nitrogen balance technique 15(1986)21

Protein and energy utilisation in rats of diets based on lupin seed (*Lupinus albus* var. multolupa) 15(1986)33

Comparative bioavailable energy values of five ground cereal grains measured with growing rats and pigs 18(1987)143

Evaluation of triticale given to pigs, poultry and rats 19(1988)359

Nutritive value of Nigerian rubber seed (*Hevea brasiliensis*) meal and oil. III. Performance characteristics, relative organ weights, hematocrit and plasma metabolites of growing female rats fed corn diets containing rubber seed meal, soya bean meal or casein 20(1988)125

Digestion and fermentation pattern of various dietary fiber sources in the rat 23(1989)195

Reconstitution

The role of germination in sorghum reconstitution 14(1986)243

Response of different sorghum grain and maize varieties to reconstitution 15(1986) 231

Red clover

Effects of ozonation on the chemical composition and in vitro degradation of triticale internodes and red clover stems 24(1989) 159

Rhizopus chinensis

Further thermotolerant fungi for the conversion of cassava starch to protein 2(1977)7

Rhodes grass

Dry matter, protein, energy and fibre intake by dairy heifers grazing a Rhodes grass (*Chloris gayana*) pasture 6(1981)15

Rice bran

Trypsin inhibitor, mineral availability, and performance of broiler chickens fed on diets based on rice bran 4(1979)133

Rice straw

Effect of alkali-treated rice straw supplemented with spent tea leaf and thyroprotein on milk yield, milk composition and certain physiological parameters of dairy cows 7(1982)201

Supplementation of rice straw with crop by-products. I. Legume straw supplementation 19(1988)43

Nutritive value of rice straw (*Oryza sativa*) ensiled with animal excreta and rumen digesta 24(1989)69

Effect of urea treatment on rumen degradation characteristics of rice straws 24(1989) 83

Rubber-seed meal

Nutritive value of Nigerian rubber seed (*Hevea brasiliensis*) meal and oil. III. Performance characteristics, relative organ weights, hematocrit and plasma metabolites of growing female rats fed corn diets containing rubber seed meal, soya bean meal or casein 20(1988)125

Rumen contents

Nutritive value of rumen contents for monogastric animals 2(1977)351

Rumen degradation

Rumen degradation of protein in sunflower meal, rapeseed meal and lupin seed placed in nylon bags 11(1984)87

Rumen degredation

Evaluation of mathematical models to describe neutral detergent residue in terms of its susceptibility to degradation in the rumen 15(1986)249

Rumen fluid

Effect of absence of ciliate protoza on rumen fluid volume, flow rate and bacterial populations in sheep 10(1983/1984)145

Effect of sodium silicate on forage digestion with rumen fluid of goats or cellulase using culture solutions adjusted for pH 24(1989) 173

Rumen ingesta kinetics

Rumen ingesta kinetics of cell wall components in dairy cows 25(1989)89

Rumen liquor
Effects of feeding alkali-treated olive cake on intake, digestibility and rumen liquor parameters 14(1986)139

Rumen microbiology
Nitrogen levels in low-roughage diets for efficient rumen microbial protein production 2(1977)255

Manipulation of rumen fluid pH and its influence on cellulolysis in sacco, dry matter degradation and the rumen microflora of sheep offered either hay or concentrate 10 (1983/1984)1

Effect of absence of ciliate protoza on rumen fluid volume, flow rate and bacterial populations in sheep 10(1983/1984)145

Influence of heat sterilization and ammoniation on straw composition and degradation by pure cultures of cellulolytic rumen bacteria 12(1984/1985)195

Effect of lactate on the in vitro digestion of *Agropyron elongatum* by rumen microorganisms 16(1986/1987)161

Population changes of fibrolytic rumen bacteria in the presence of phenolic acids and plant extracts 19(1988)261

The establishment of rumen microbial communities 21(1988)69

Possibilities of the use of recombinant-DNA techniques with rumen micro-organisms 21 (1988)161

Consecutive batch culture – a novel technique for the in vitro study of mixed microbial populations from the rumen 21(1988)183

Factors affecting the formation of polysaccharide-degrading enzymes by rumen microorganisms 21(1988)191

Use of the rumen simulation technique (RUSITEC) to compare microbial digestion of various by-products 21(1988)197

Lignin-polysaccharide complexes of the plant cell wall and their effect on microbial degradation in the rumen 21(1988)219

Effect of defaunating the rumen 21(1988)229

Biological structure of lignocellulose and its degradation in the rumen 21(1988)295

Colonization of guinea grass by anaerobic rumen fungi in swamp buffalo and cattle 22 (1988)161

Influence of stage of forage maturity and ammoniation of wheat straw on ruminal degradation of wheat forage tissues 24 (1989)201

Evaluation of various methods for protecting soya bean protein from degradation by rumen bacteria 25(1989)111

Rumen pH
Manipulation of rumen fluid pH and its influence on cellulolysis in sacco, dry matter degradation and the rumen microflora of sheep offered either hay or concentrate 10 (1983/1984)1

Associative effects of mixed feeds. I. Effects of type and level of supplementation and the influence of the rumen fluid pH on cellulolysis in vivo and dry matter digestion of various roughages 10(1983/1984)15

Use of fat in diets of sheep in hot environments. II. Effect on rumen metabolism 19 (1988)343

Efficacy of supplemental dietary neutralizing agents for lactating dairy cows. A review 23(1989)277

Rye grain
Feeding value of alkali-treated whole rye (*Secale cereale* L.) grain for lactating cows and its digestibility for sheep 10(1983/1984)77

Ryegrass
Apparent and true digestibility of the Weende components, cell content and cell wall of ryegrass 1(1976)607

Ensiled alkali-treated straw. II. The nutritive value for young beef cattle of mixtures of ensiled or frozen alkali-treated straw and ryegrass silage 3(1978)133

The effect of silage additives containing formaldehyde on the fermentation of ryegrass ensiled at different dry matter levels and on the nutritive value of direct-cut silage 7(1982)303

The effect of urea on the preservation and digestibility in vitro of perennial ryegrass 10(1983/1984)49

The effect of anhydrous ammonia and moisture content on the preservation and chemical composition of perennial ryegrass hay 11(1984)159

Effect of prolonged wilting in poor conditions on the fermentation quality, metabolisability and net energy value of silage given to sheep 12(1984/1985)109

The chemical composition of grass silages prepared with and without pre-treatment with inoculants containing *Lactobacillus plantarum* 13(1985)269

Fractions of nitrogen, sulphur, phosphorus, calcium and magnesium in the herbage of perennial ryegrass as influenced by fertilizer nitrogen 14(1986)231

Treatment of whole-crop cereals with alkali. 1. The influence of sodium hydroxide and ensiling on the chemical composition and in-vitro digestibility of rye, barley and wheat

crops harvested at increasing maturity and dry matter content 18(1987)257

Treatment of whole-crop cereals with alkali. 2. Voluntary intake and digestibility by sheep of rye, barley and wheat crops ensiled with sodium hydroxide 18(1987)271

The effects of inoculation, addition of glucose and mincing on fermentation and proteolysis in ryegrass ensiled in laboratory silos 19 (1988)85

Effects of enzyme supplementation of diets based on wheat, rye or triticale on their productive value for broiler chickens 20 (1988)313

S

Safflower oilseed meal
Value of mechanically extracted rape and safflower oilseed meals as protein supplements for growing lambs 11(1984)269

Sal seed meal
Tannins in sal seed (*Shorea robusta*) and sal seed meal limit their utilization as livestock feeds 7(1982)161

Sawdust
Investigations of the treatment of sawdust for rabbit feeding. 1. Effect of sodium hydroxide treatment 6(1981)43

Investigations of the treatment of sawdust for rabbit feeding. 2. Effect of boiling and incubation with poultry manure 6(1981)51

Seasonality
Seasonal changes in composition and intake of chaparral by Spanish goats 8(1983)47

Chemical composition and animal performance of grass forage of varying maturity stored as hay or silage 11(1984)35

Seed dehulling
A simple device for dehulling seeds and grain 3(1978)109

Selenium
Treatment of dairy sheep with soluble glass boluses containing copper, cobalt and selenium 19(1988)79

The effect of soluble-glass boluses containing copper, cobalt and selenium on the blood composition of ewes 21(1988)23

Sesbania
Canavanine content and toxicity of sesbania leaf meal for growing chicks 25(1989)137

Sesbania sesban
Yield, composition and feeding value of irrigated *Sesbania sesban* var. *Nubia* leaves harvested at latitude 38°N during a Mediterranean summer 18(1987)247

Sheep–adult
Long term feeding of concentrate fumigated with carbon tetrachloride to sheep 2(1977) 77

The nutritional value for sheep of molasses and condensed molasses solubles 2(1977) 143

Utilization by sheep of dried pig faeces containing a high concentration of copper 3 (1978)31

Differences between cattle and sheep in their digestion and relative intake of a mature tropical grass hay 3(1978)41

Estimation of the digestibility of low-quality hays by cattle from measurements made with sheep 3(1978)51

Spent tea leaf as a ruminant feed 3(1978)219

Chemical composition of culinary wastes and their potential as a feed for ruminants 5 (1980)205

Intake and digestibility in sheep fed on alkali-treated straw and concentrates, as complete diets or separate feeds 7(1982)271

Solubility and flow of calcium, magnesium and phosphorus in the digestive tract of sheep given maize or alfalfa silages 9(1983) 131

Effect of intake on digestibility of plant cell walls and cell contents of complete diets for ruminants 9(1983)197

Manipulation of rumen fluid pH and its influence on cellulolysis in sacco, dry matter degradation and the rumen microflora of sheep offered either hay or concentrate 10 (1983/1984)1

Digestion of organic matter and nitrogen in ammoniated barley straw 10(1983/1984)65

Feeding value of alkali-treated whole rye (*Secale cereale* L.) grain for lactating cows and its digestibility for sheep 10(1983/1984)77

The effect of milling and sodium hydroxide treatment on intake and digestibility of wheat straw by sheep 11(1984)149

Feeding value of apple pomace silage for sheep 11(1984)189

Degradation of crude protein and quality of undegradable protein in untreated or formaldehyde-treated rapeseed meal 12 (1984/1985)19

A comparison of laboratory methods involving single or two-stage cellulase and two-stage rumen inoculum–pepsin treatment

for predicting digestibility and intake of hays in sheep 12(1984/1985)101

Effect of prolonged wilting in poor conditions on the fermentation quality, metabolisability and net energy value of silage given to sheep 12(1984/1985)109

The utilization of untreated and calcium hydroxide treated wheat straw by sheep 12 (1984/1985)141

The effect of supplementing grass silage with sodium hydroxide-treated chopped or milled straw on intake and digestibility by wether sheep 12(1984/1985)215

The voluntary intake and digestibility of combinations of cereal crop residues and legume hay for sheep 12(1984/1985)241

Performance and carcass characteristics of Sudan desert sheep and goats on high roughage diets with added fat 13(1985)147

Studies on the feeding value of agro-industrial by-products. III. Digestibility of cocoa-pod and cocoa-pod-based diets by ruminants 13 (1985)249

Corrugated cardboard and dried poultry manure as dietary ingredients for sheep 14 (1986)265

The effect of monensin on rumen fermentation in sheep fed on all-roughage and barley/roughage diets 15(1986)7

The effect of grinding groundnut hulls either with or without alkali treatment on digestibility of diet and on ruminal and blood components 15(1986)69

The digestion by sheep of barley and maize-based diets containing either meat and bone meal or soya bean meal 15(1986)301

Comparison between two methods of feeding concentrate to sheep given a diet high or low in concentrate with or without buffering substances 16(1986/1987)37

Relative intakes, digestibility and bacterial protein synthesis by sheep and cattle fed high-roughage diets 16(1986/1987)75

Composition of vine-branches with leaves of *Vitis vinifera* L. fresh and ensiled using different additives (previous results) 16(1986/1987)89

The intake and digestibility by sheep of oat straw and maize stover offered with different levels of noug (*Guizotia abyssinica*) meal 16(1986/1987)99

Effect of defaunation on the loss of energy in wethers fed different quantities of cellulose and normal or steamflaked maize starch 16 (1986/1987)233

Effect of formaldehyde-treated soya bean meal and urea in starters on nitrogen quality, degradibility in sacco, sheep digestibility and calf performance 16(1986/1987)287

Nutritive value of three cultivars of triticale for sheep, pigs and poultry 18(1987)19

The effect of dietary barley on carbohydrate digestibility of sulphur dioxide-treated wheat straw by sheep 18(1987)55

Nutritive value of naked oats for ruminants 18(1987)83

Treatment of whole-crop cereals with alkali. 2. Voluntary intake and digestibility by sheep of rye, barley and wheat crops ensiled with sodium hydroxide 18(1987)271

Treatment of whole-crop cereals with alkali. 3. Voluntary intake and digestibility studies in sheep given immature wheat ensiled with sodium hydroxide, urea or ammonia 18(1987)283

Treatment of dairy sheep with soluble glass boluses containing copper, cobalt and selenium 19(1988)79

Use of fat in diets of sheep in hot environments. II. Effect on rumen metabolism 19 (1988)343

Evaluation of mixed lucerne and caucasian bluestem hay diets fed to sheep 20(1988) 19

The effect of age of regrowth on content and release of manganese, iron, zinc and copper from four tropical forages incubated in sacco in rumen of sheep 20(1988)171

The intake and digestion of maize silage-based diets by dairy cows and sheep 20 (1988)299

The effect of dietary inclusion of starch, sucrose and xylose on the utilization of dietary energy in sheep 21(1988)11

The effect of soluble-glass boluses containing copper, cobalt and selenium on the blood composition of ewes 21(1988)23

Amino acids flow to and absorption from the small intestine of sheep fed barley and sulfur dioxide-treated straw at different ratios 22(1988)147

Digestibility in vivo of ensiled grapevines (branches and leaves): influence of the system of analysis in the detergent fibre scheme on the prediction of digestibility 22 (1988)173

The nutritive value of ensiled forages cut at three stages of growth 22(1988)237

Effect of soaking, ensilage and hydrogen peroxide treatment of barley straw on rumen degradability 22(1988)295

Effect of ammoniation and energy supplementation on the utilization of wheat straw by sheep 22(1988)305

Nutritive value of pungam (*Pongamia glabra* Vent) cake for sheep 22(1988)321

The effects of starch and pectin-rich diets on quantitative aspects of digestion in sheep 24(1989)289

Oven and stack ammoniation of grass hays. 2. Voluntary intake, digestibility and faecal nitrogen excretion in sheep in relation to hay composition 24(1989)313

Effect of formaldehyde treatment on the rumen digestion of barley and wheat grain and its influence on hay intake and digestion by sheep 25(1989)55

By-products from the canning industry as feedstuff for ruminants: digestibility and its prediction from chemical composition and laboratory bioassays 25(1989)67

Digestion rates in and outflow rates from the rumen of sheep fed untreated or calcium hydroxide-treated wheat straw 25(1989) 179

Sheep–general

Raw and steam-pelleted cassava, sweet potato and yam *cayenensis* as starch sources for ruminant and chicken diets 3(1978)73

Comparative digestibility of feedstuffs by sheep and cows 12(1984/1985)47

The effect of five different supplements on the degradation of straw in sheep given untreated barley straw 19(1988)289

Use of fat in diets of sheep in hot environments. I. Effects on performance, carcass characteristics and lipid composition of plasma 19(1988)327

Sheep–lamb

Poly-unsaturated fat in young ruminants 2 (1977)93

Reducing the potential copper toxity of concentrates to sheep by the use of molybdenum and sulphur supplements 2(1977)235

The use of whole barley diets fortified with solutions of urea, minerals and vitamins for lambs 2(1977)307

The effect of ammoniation on the intake and nutritive value of straw 4(1979)169

Nutritive value of thermoammoniated and steam-treated maize stover I. Intake, digestibility and nitrogen retention 4(1979) 177

Nutritive value of thermoammoniated and steam-treated maize stover II. Rumen metabolites and rate of passage 4(1979)187

Citrus pulp in diets for fattening lambs 5 (1980)11

Effect of dry-matter concentration of milk substitute and method of feeding on intake and performance by lambs 5(1980)321

Effects of treatment of barley straw with anhydrous ammonia and supplementation with heat-treated protein meals on feed intake and liveweight performance of growing lambs 6(1981)145

Utilization of alkali-treated grain. 1. Alkali-treated grain in complete diets for steers and lambs 6(1981)273

Use of processing to explore the factors affecting the digestion of forage cell walls 6 (1981)387

Use of fish protein hydrolysate in milk replacers 7(1982)135

Evaluation of corn gluten meal with urea as a source of supplementary nitrogen for growing calves and lambs 7(1982)375

Nutritive value of almond hulls for sheep 8 (1983)63

The effects of sulphur supplementation on cellulose digestion in vitro and on nutrient digestion, nitrogen metabolism and rumen characteristics of lambs fed on good quality fescue and tropical star grass hays 8(1983) 129

Methods of improving the utilization of cereal straw by ruminants. II. Further studies on dried grass as a supplement for lambs 8 (1983)229

Can lantana seeds be utilized in lamb rations? 9(1983)29

Energy and nitrogen balance, intake and growth rate of lambs fed on unwilted grass silages treated with a formaldehyde–acid mixture or untreated 9(1983)107

Chemical composition, digestibility and feeding value of maize cobs 9(1983)121

The use of Balanites kernel cake in a diet for fattening sheep 9(1983)301

Effect of absence of ciliate protoza on rumen fluid volume, flow rate and bacterial populations in sheep 10(1983/1984)145

Biological availability of phosphorus from basic steel slag 11(1984)75

Maize whole stillage as a supplemental protein source for growing lambs and finishing steers 11(1984)181

Nitrogen balance and abomasal protein-nitrogen flow in growing ruminants fed methane digester effluent in combination with distillers dried grains 13(1985)47

High temperature and pressure processing of maize cobs. II. Animal response and effect of sodium metabisulphite 14(1986)161

Effects of supplementing a wheat straw diet with urea or a urea–molasses block and/or cottonseed meal on intake and liveweight change of lambs 16(1986/1987)25

Energy metabolism in lambs fed silage diets with sucrose or a growth promoter 16 (1986/1987)203

Cassava meal as supplement to napier grass diets for growing sheep and goats 18(1987) 197

SO_2 treatment for converting straw into a concentrate-like feed: a growth study with lambs 19(1988)219

Effect of sorghum straw alone or in combination with molasses and nitrogen sources on performance of Sudan desert lambs 19 (1988)351

Nitrogen utilization, ruminal fermentation and abomasal nitrogen flow in sheep fed caucasian bluestem hay supplemented with lucerne or urea 20(1988)33

The maize replacement value of cassava flour in rations for growing ewe lambs 20(1988) 233

In vitro studies and short-term feeding trial in lambs to evaluate plantain sheath (*Musa sapientum*) as a feed for ruminants 20 (1988)343

Pilot plant production of sulphur dioxide-treated mesquite wood. II. Digestibility and feedlot performance by lambs 22(1988) 61

Nutritive value of banana stalk (*Musa cavendishi*) as a feed for sheep 22(1988)327

Fermentation and utilization by lambs of sugarcane harvested fresh and ensiled with and without NaOH. 4. Ruminal kinetics 23 (1989)323

Silage–alfalfa

Solubility and flow of calcium, magnesium and phosphorus in the digestive tract of sheep given maize or alfalfa silages 9(1983) 131

Silage–animal waste

Ensiling of rumen contents and blood with wheat straw 18(1987)67

Silage–apple pomace

Feeding value of apple pomace silage for sheep 11(1984)189

Some properties of apple pomace ensiled with and without additives 15(1986)57

Silage–barley

Ensiled alkali-treated straw. I. Effect of level and type of alkali on the composition and digestibility in vitro of ensiled barley straw 3(1978)117

Ensiled alkali-treated straw. II. The nutritive value for young beef cattle of mixtures of ensiled or frozen alkali-treated straw and ryegrass silage 3(1978)133

Akali treatment of straw for ruminants. II. Nutritive value of straw ensiled after alkali treatment. 3(1978)289

Effect of soaking, ensilage and hydrogen peroxide treatment of barley straw on rumen degradability 22(1988)295

Silage–beet

Distillery effluents as animal feed: the use of condensed beet molasses stillage (CBMS) in broiler feeding 5(1980)233

Silage–clover

The enzymic nature of the degradation of beta-carotene in red clover and in other forage crops during silage-making with acid additives 5(1980)59

Silage–fish

Nutritional value and mercury content of fish silage for growing pigs 9(1983)169

Thiaminase activity in fish silage and moist fish feed 13(1985)39

Effect of formaldehyde treatment on the degradation of acid-preserved fish silage protein in vitro 16(1986/1987)297

Fish silage as a protein supplement for early-weaned calves 17(1987)165

Examination of stability, and its effect on the nutritive value, of fish silage in diets for growing pigs 21(1988)43

Silage–forage, general

Losses of beta-carotene in unwilted forage crops during silage-making and feeding 9 (1983)63

Silage–general

Treatment of whole-crop cereals with alkali. 1. The influence of sodium hydroxide and ensiling on the chemical composition and in vitro digestibility of rye, barley and wheat crops harvested at increasing maturity and dry matter content 18(1987)257

Treatment of whole-crop cereals with alkali. 2. Voluntary intake and digestibility by sheep of rye, barley and wheat crops ensiled with sodium hydroxide 18(1987)271

The nutritive value of ensiled forages cut at three stages of growth 22(1988)237

Silage–grass

The intake of steers and heifers fed on mixtures of alkali-treated or untreated straw with grass silage 2(1977)315

The use of effluent from grass silage in the diet of finishing pigs. I. Variation in composition of effluents 4(1979)263

The use of effluent from grass silage in the diet of finishing pigs. II. Assessment of nutritive value of fresh and stored effluent 4(1979) 275

A calorimetric investigation of the nutritive value of formic acid- and formaldehyde–formic acid-treated grass silages 6(1981)245

Feed consumption and performance of dairy cows with alternate feedings of grass silage and maize silage 6(1981)337

The effect of silage additives containing formaldehyde on the fermentation of ryegrass ensiled at different dry matter levels and on the nutritive value of direct-cut silage 7(1982)303

Investigations on the effect of sodium bentonite and other moisture absorbents on the production of effluent from grass silage 8(1983)107

Energy and nitrogen balance, intake and growth rate of lambs fed on unwilted grass silages treated with a formaldehyde–acid mixture or untreated 9(1983)107

Chemical composition and animal performance of grass forage of varying maturity stored as hay or silage 11(1984)35

Digestibility and digesta retention in dairy cows receiving hay or silage at varying concentrate levels 11(1984)279

The effect of supplementing grass silage with sodium hydroxide-treated chopped or milled straw on intake and digestibility by wether sheep 12(1984/1985)215

The chemical composition of grass silages prepared with and without pre-treatment with inoculants containing *Lactobacillus plantarum* 13(1985)269

Studies on the nutritive value of roots, cabbage and grass silage for growing commercial rabbits 13(1985)299

The substitution of grass silage by a straw and concentrate mixture for dairy cows 20 (1988)135

The digestibility and metabolisable energy content of grass silage and their prediction from laboratory measurements 24(1989)27

Silage–legume

Ensiling of leguminous green forages in combination with different dry roughages and molasses 12(1984/1985)133

Silage–lucerne

SO_2-treated straw as a silage additive: fermentation data on lucerne with particular reference to protein degradation 22(1988) 247

Silage–maize

Forage maize in the rations of dairy cows and fattening cattle in Bulgaria 1(1976)45

Cultivation of maize in irrigated areas for silage and grain production and their use by ruminants 1(1976)237

Influence of ensiling maize ears and grain on chemical composition, conservation losses and digestibility 1(1976)301

Effect of maize silage harvest stage on yield, plant composition and fermentation losses 1(1976)313

Maize silage with or without NPN, dehydrated whole-crop maize pellets or high moisture maize grain for finishing bulls 1 (1976)347

Energy supplementation of maize silage harvested at different maturity stages 1(1976) 369

Factors influencing the composition and nutritive value of ensiled whole-crop maize 1 (1976)381

Rate of growth, feed utilisation and digestibility observed with diets containing ensiled maize grain or maize ears fed alone or in mixtures 1(1976)401

Voluntary intake and efficiency of utilisation of whole-crop maize silage 1(1976)441

Levels, sources and methods of nitrogen supplementation of maize silage for beef production 1(1976)465

The nutritive value (energy) of maize silage 1 (1976)481

Effect of weight at slaughter on the efficiency of fattening young cattle 1(1976)505

Economic aspects of high-plane feeding of bulls with maize silage 1(1976)513

Maize silage and whole maize plant pellets as the basic forages in complete feeds for milk cows 3(1978)23

A note on the effect on the aerobic deterioration of maize silage of the manipulation of the microflora by means of antibiotics 3 (1978)89

Amino acid composition of maize silage produced in the United Kingdom 4(1979)163

Feed consumption and performance of dairy cows with alternate feedings of grass silage and maize silage 6(1981)337

Solubility and flow of calcium, magnesium and phosphorus in the digestive tract of sheep given maize or alfalfa silages 9(1983) 131

Effect of wilting and the additives, straw, molasses and urea on the fermentation pattern of maize silage 9(1983)185

Effects of stage of maturity and addition of molasses on nutritive value of maize stover silage 12(1984/1985)65

Effect of sodium acrylate and calcium and sodium hydroxides on the fermentation, digestibility, chemical composition and aerobic deterioration of whole-crop maize silage 12(1984/1985)91

The effect of nitrogen supplements to maize silage on the amount and composition of the nitrogen fraction entering the intestine of cattle 13(1985)25

Influence of harvest date on the nutritive value and ensiling characteristics of maize stover 14(1986)11

High grain content maize silage in fattening diets of young male cattle 16(1986/1987)63

Feeding value of ensiled high-moisture maize grain with cob for growing-finishing pigs 20 (1988)145

The intake and digestion of maize silage-based diets by dairy cows and sheep 20 (1988)299

Effects of added dietary urea on the utilization of maize stover silage by growing beef cattle 22(1988)227

Inoculation of corn silage and high-moisture corn with lactic acid bacteria and its effects on the subsequent fermentations and on feedlot performance of beef steers 25(1989) 23

Silage–manure–maize

Laboratory evaluation of ensiled mixtures of alkali-treated cattle excreta and whole-crop maize 3(1978)335

Silage–orange peel

The effect of treatment with urea, sorbic acid, or dehydration on orange peel silage 20 (1988)335

Silage–orange pulp

Effect of urea on the ensiling process of orange pulp 12(1984/1985)233

Silage–para grass

Note on high-nitrate para grass (*Brachiaria mutica*) silage given to rabbits 9(1983)325

Silage–quality assessment

A titration method for silage assessment 22 (1988)341

Silage–ryegrass

The effect of some pre-ensiling treatments on silage composition and nitrogen disappearance in the rumen 8(1983)259

Effect of prolonged wilting in poor conditions on the fermentation quality, metabolisability and net energy value of silage given to sheep 12(1984/1985)109

Energy metabolism in lambs fed silage diets with sucrose or a growth promoter 16 (1986/1987)203

The effects of inoculation, addition of glucose and mincing on fermentation and proteolysis in ryegrass ensiled in laboratory silos 19 (1988)85

Silage–sorghum

Studies on fermentation of sorghum silage during storage - effect of urea and molasses 3(1978)299

Preservation of sorghum plant portions harvested, processed and ensiled at ten stages of maturity 7(1982)257

Studies on fermentation of sorghum silage during storage, and its effect on milch animals 9(1983)143

Effects of molasses/urea and bacterial inoculant additives on silage quality, dry matter recovery, and feeding value for cattle 12 (1984/1985)205

Silage–straw

Methods of improving the utilization of cereal straw by ruminants. III. A note on the effect of ensiling straw treated with urea 9(1983)181

Ensiling characteristics of cattle waste fermented with wheat straw and green berseem (*Trifolium alexandrium*) 25(1989)149

Silage–sugar beet

A note on the use of sugar-beet pulp silage and molasses in the diet of lactating dairy cows 20(1988)13

Counteracting structure loss in pressed sugar beet pulp silage 24(1989)97

Silage–sugar cane

Nutritive value for ruminants of sugar cane bagasse ensiled after spray treatment with different levels of NaOH 9(1983)1

Fermentation and utilization by lambs of sugarcane harvested fresh and ensiled with and without NaOH. 4. Ruminal kinetics 23 (1989)323

Silage–tropical forages

Effects of ensiling or drying on five forage species in western Kenya: *Zea mays* (maize stover), *Pennisetum purpureum* (Pakistan napier grass), *Pennisetum* sp. (bana grass), *Impomea batata* (sweet potato vines) and *Cajanus cajan* (pigeon pea leaves) 13(1985) 1

Silage–vine
Composition of vine-branches with leaves of *Vitis vinifera* L. fresh and ensiled using different additives (previous results) 16(1986/1987)89

Silage–wheat
Wheat silage for fattening young male cattle 12(1984/1985)253

Treatment of whole-crop cereals with alkali. 4. Voluntary intake and growth in steers given wheat ensiled with sodium hydroxide, urea or ammonia 19(1988)55

Silage disease
Inhibition of *Listeria monocytogenes* by defined lactic microflora in gnotobiotic silages of lucerne, fescue, ryegrass and maize – influence of dry matter and temperature 2(1977)297

Silage effluent
Investigations on the effect of sodium bentonite and other moisture absorbents on the production of effluent from grass silage 8(1983)107

Silica
Effects of the addition of silica to the diet of the laying hen, with particular reference to egg-shell quality 6(1981)157

Single cell protein
Utilization of cattle manure for Torula yeast production from straw hydrolysates 5(1980)175

Utilisation of L- and DL-amino acids by broiler chicks fed diets containing Pruteen 17(1987)1

Single-cell protein
Composition and nutritive value of single-cell protein (SCP) 1(1976)9

Digestibility of protein and amino acids of a fermentation single-cell protein for veal calves 2(1977)377

Nutritive value of single-cell protein produced by *Chaetomium cellulolyticum* grown on maize stover and pulp-mill sludge 11(1984)211

Sisal
Feed value of Tanzanian sisal wastes: chemical composition and digestibility in vitro 14(1986)255

Sodium acrylate
The effect of calcium and sodium hydroxides and of sodium acrylate on the fermentation and digestibility in vitro of ensiled whole-crop wheat and barley harvested at different stages of maturity 9(1983)37

Effect of sodium acrylate and calcium and sodium hydroxides on the fermentation, digestibility, chemical composition and aerobic deterioration of whole-crop maize silage 12(1984/1985)91

Sodium bentonite
Investigations on the effect of sodium bentonite and other moisture absorbents on the production of effluent from grass silage 8(1983)107

Effects of pelleting, added sodium bentonite and fat in a wheat-based diet on performance and carcass characteristics of small white turkeys 12(1984/1985)223

Sodium bicarbonate
Effects of sodium chloride and sodium bicarbonate in the diet on the performance of calves 5(1980)309

Efficacy of supplemental dietary neutralizing agents for lactating dairy cows. A review 23(1989)277

Sodium chloride
Effects of sodium chloride and sodium bicarbonate in the diet on the performance of calves 5(1980)309

Sodium diacetate
Storage of large round bales of hay harvested at different moisture concentrations and treated with sodium diacetate and/or covered with plastic 13(1985)69

Microfloral and compositional changes in alfalfa hay treated with sodium diacetate and stored at different moisture contents 17(1987)45

Sodium hydroxide
The effect of calcium and sodium hydroxides and of sodium acrylate on the fermentation and digestibility in vitro of ensiled whole-crop wheat and barley harvested at different stages of maturity 9(1983)37

Ammonia treatment of straw via the hydrolysis of urea. II. Additions of soya bean (urease), sodium hydroxide and molasses; effects on the digestibility of urea-treated straw 11(1984)115

The effect of milling and sodium hydroxide treatment on intake and digestibility of wheat straw by sheep 11(1984)149

Sodium metabisulphite
High temperature and pressure processing of maize cobs. II. Animal response and effect of sodium metabisulphite 14(1986)161

Sodium silicate
Effect of sodium silicate on forage digestion with rumen fluid of goats or cellulase using culture solutions adjusted for pH 24(1989)173

Soil chemistry
Soil chemistry and mineral problems in farm livestock. A review 5(1980)95

Sophora japonica
Yield and composition of protein concentrate, press cake, green juice and solubles concentrate from wet fractionation of *Sophora japonica* L. foliage 20(1988)177

Sorbic acid
The effect of treatment with urea, sorbic acid, or dehydration on orange peel silage 20 (1988)335

Sorghum
Influence of pericarp and endosperm colour and type on digestibility of grain sorghum by pigs 2(1977)219

Relationships between tristimulus colour values and digestibility of grain sorghum by swine 2(1977)327

Preservation of sorghum plant portions harvested, processed and ensiled at ten stages of maturity 7(1982)257

Proximate analysis, amino acid composition and tannin content of improved Nigerian sorghum varieties and their potential in poultry feeds 7(1982)359

Waxy and normal grain sorghums with varying tannin contents in diets for young pigs 12(1984/1985)179

The role of germination in sorghum reconstitution 14(1986)243

Yield and potential feeding value of straws of grain sorghum cultivars developed in Nigeria 17(1987)285

Effects of wood ash extract treatment on the feeding value and utilization of high-tannin sorghums by broiler chicks 22(1988)131

Factors effecting the digestibility by pigs of energy and protein in wheat, barley and sorghum diets supplemented with canola meal 24(1989)253

Sorghum grain
A comparison of three fibre methods for predicting the metabolizable energy content of sorghum grain for poultry 2(1977)197

Amino acid composition of two varieties of sorghum grain 5(1980)169

Comparison of the nutritional value of low and medium tannin sorghum grains for pigs 8(1983)85

Response of different sorghum grain and maize varieties to reconstitution 15(1986) 231

Treatment of low-tannin sorghum grain for broiler feed 20(1988)69

Sorghum silage
Studies on fermentation of sorghum silage during storage - effect of urea and molasses 3(1978)299

Studies on fermentation of sorghum silage during storage, and its effect on milch animals 9(1983)143

Sorghum straw
Effect of sorghum straw alone or in combination with molasses and nitrogen sources on performance of Sudan desert lambs 19 (1988)351

Soya bean
Effects of micronization on the digestibility of whole soya beans and rapeseeds for the growing pig 3(1978)179

Effect of processing on the utilization by young pigs of the fat in soya beans and sunflower seeds 12(1984/1985)267

Nutrient and trypsin inhibitor content of hydroponically sprouted soya beans 13(1985) 203

Effects of moisture level, treatment time and soya bean addition on the nutritive value of urea-treated maize stover 19(1988)67

The energy content of full-fat soya beans in meal and pelleted feeds for adult cocks and broilers 19(1988)97

Alkali-treated soya flour for piglets 24(1989)1

Evaluation of various methods for protecting soya bean protein from degradation by rumen bacteria 25(1989)111

Soya bean hulls
Colonization of thick-walled plant tissues by anaerobic fungi 19(1988)25

Soya bean meal
Inhibition of ruminal deamination in vitro by formaldehyde treatment of sunflower-seed, soya bean and fish meals: response curves to protective treatment 2(1977)267

Alkali treatment of straw for ruminants. I. Utilization of complete diets containing straw by beef cattle 3(1978)143

Meat meal in the diet of the early-weaned pig. I. A comparison of meat meal and soya bean meal 4(1979)33

Ammonia treatment of straw via the hydrolysis of urea. II. Additions of soya bean (urease), sodium hydroxide and molasses; effects on the digestibility of urea-treated straw 11(1984)115

Effects of isopropyl alcohol treatment of soya bean meal on nitrogen utilization by ruminants 14(1986)127

Apparent ileal and faecal digestibility of amino acids in barley-based diets supplemented with soya bean meal or canola meal for growing pigs 14(1986)183

The digestion by sheep of barley and maize-based diets containing either meat and bone meal or soya bean meal 15(1986)301

Effect of formaldehyde-treated soya bean meal and urea in starters on nitrogen quality, degradibility in sacco, sheep digestibility and calf performance 16(1986/1987)287

Nutritive value of Nigerian rubber seed (*Hevea brasiliensis*) meal and oil. III. Performance characteristics, relative organ weights, hematocrit and plasma metabolites of growing female rats fed corn diets containing rubber seed meal, soya bean meal or casein 20(1988)125

Spruce muka

Nutritive value of spruce muka (foliage) for the growing chick 3(1978)63

Starch

Further thermotolerant fungi for the conversion of cassava starch to protein 2(1977)7

Raw and steam-pelleted cassava, sweet potato and yam *cayenensis* as starch sources for ruminant and chicken diets 3(1978)73

The effect of β-glucanase on the utilization of starch and nitrogen by broiler chickens fed on barley of low- or high-viscosity 15(1986) 83

The effect of dietary inclusion of starch, sucrose and xylose on the utilization of dietary energy in sheep 21(1988)11

The effects of starch and pectin-rich diets on quantitative aspects of digestion in sheep 24(1989)289

Steam treatment

Effect of high-pressure steam treatment on chemical composition and digestibility in vitro of roughages 7(1982)61

Stearic acid

Feed intake and milk yield at increasing supplement of a palmitic and stearic acid-rich type of fat in comparison with animal fat 22(1988)179

Stockfeed, general

Energetics of stockfeed production 3(1978)1

Storage

Studies on fermentation of sorghum silage during storage - effect of urea and molasses 3(1978)299

Effect of time of storage and method of drying on the digestibility in vitro of alkali-treated barley straw 3(1978)381

Effects of different handling systems on pig feed quality 7(1982)11

The effect of prolonged storage on the digestibility and nitrogen content of ammonia-treated barley straw 8(1983)147

Effects of dietary maize solubles and minerals on albumen quality of fresh and stored brown-shelled eggs 8(1983)177

Effect of storage and processing of barley on its utilization by pigs and rats 9(1983)89

Studies on fermentation of sorghum silage during storage, and its effect on milch animals 9(1983)143

Investigations into the effects of processing on the retention of the carotenoid fractions of *Leucaena leucocephala* during storage, and the effects of processing on mimosine concentration 9(1983)307

The influence of processing and storage on hydrogen cyanide and tannin contents of para-rubber seed and its products 9(1983) 319

Anaerobically stored high moisture cereals for growing pigs 12(1984/1985)159

Storage of large round bales of hay harvested at different moisture concentrations and treated with sodium diacetate and/or covered with plastic 13(1985)69

Liquid potato feed for finishing pigs: feeding value, inclusion rate and storage properties 15(1986)129

Nutrient losses during deterioration of hay in relation to changes in biochemical composition and microbial growth 15(1986)149

The nutritive value of barley for monogastric animals and for ruminants as affected by artificial drying and alkali and anaerobic storage 15(1986)285

Microfloral and compositional changes in alfalfa hay treated with sodium diacetate and stored at different moisture contents 17(1987)45

Influence of processing methods and storage time on the cyanide potential of cassava leaf meal 17(1987)227

Microbial dynamics in barley grain stored under controlled atmosphere 19(1988)299

Preservation and storage of green panic (*Panicum maximum*) as moist hay with urea 20 (1988)87

The effect of method of preservation of forages on their resistance to reducion in particle size with an artificial masticator 24 (1989)185

Straw–barley

Effect on digestibility and nitrogen content of barley straw of different ammonia treatments 2(1977)131

Ensiled alkali-treated straw. I. Effect of level and type of alkali on the composition and digestibility in vitro of ensiled barley straw 3(1978)117

Ensiled alkali-treated straw. II. The nutritive value for young beef cattle of mixtures of ensiled or frozen alkali-treated straw and ryegrass silage 3(1978)133

Alkali treatment of straw for ruminants. I. Utilization of complete diets containing straw by beef cattle 3(1978)143

Akali treatment of straw for ruminants. II. Nutritive value of straw ensiled after alkali treatment. 3(1978)289

Effect of time of storage and method of drying on the digestibility in vitro of alkali-treated barley straw 3(1978)381

Effects of treatment of barley straw with anhydrous ammonia and supplementation with heat-treated protein meals on feed intake and liveweight performance of growing lambs 6(1981)145

Studies on ammonia-treated straw. II. Fixation of ammonia in treated straw by spraying with acids 7(1982)53

The feeding value for ruminants of straw and whole-crop barley and oats treated with anhydrous or aqueous ammonia or urea 8 (1983)247

The effect of urease enzyme on treatment time and the nutritive value of straw treated with ammonia as urea 8(1983)271

Digestion of organic matter and nitrogen in ammoniated barley straw 10(1983/1984)65

A comparison between degradation in vitro and in sacco of constituents of untreated and ammonia-treated barley straw 10(1983/1984)199

Digestibility studies on ammonia-treated straw 10(1983/1984)213

The digestion of untreated and ammonia-treated barley straw in an artificial rumen 10(1983/1984)229

Ammonia-treated barley straw and rolled barley offered either together, in a mixed ration, or successively to beef steers 10 (1983/1984)247

A comparison of alkali treatment methods to improve the nutritive value of straw. I. Digestibility and metabolizability 12(1984/1985)295

Treatment of barley straw with ammonia and sulphur dioxide gases under laboratory conditions 14(1986)41

A comparison of alkali treatment methods used to improve the nutritive value of straw. II. In sacco and in vitro degradation relative to in vivo digestibility 14(1986)215

Evaluation of the increase in feeding value of barley straw following mild treatment with sulphur dioxide 15(1986)13

Near infrared reflectance spectroscopy of untreated and ammoniated barley straw 15 (1986)189

Effect of straw treatment and laboratory of incubation on degradation of neutral detergent residue in the rumen 18(1987)75

Fresh and dried brewer's grains as protein supplements to barley straw diets given to pregnant beef cows 19(1988)33

Effects of ammonia and sulphur dioxide gases on the composition and digestion of barley straw 19(1988)121

Fibre degradation in the rumens of animals receiving hay, untreated or ammonia-treated straw 19(1988)277

The effect of five different supplements on the degradation of straw in sheep given untreated barley straw 19(1988)289

Effect of soaking, ensilage and hydrogen peroxide treatment of barley straw on rumen degradability 22(1988)295

Upgrading barley straw for goats: the effectiveness of a sodium hydroxide and urea dip method 24(1989)57

Straw–cotton

Effects of ammonia treatment and processing of cotton straw as the sole source of roughage in complete feeds on rumen characteristics with Murrah buffaloes 14(1986) 95

Straw–general

The effect on digestibility of a new technique for alkali treatment of straw 1(1976)73

Review article: the alkali treatment of straws 2(1977)105

Enzyme solubility – a method for evaluating the digestibility of alkali-treated straw 2 (1977)205

The intake of steers and heifers fed on mixtures of alkali-treated or untreated straw with grass silage 2(1977)315

A new technique for upgrading the nutritive value of roughages under field conditions 3 (1978)201

Cellulase insoluble fibre as a measure of unavailable organic matter in cattle compounds containing alkali-treated straw 3 (1978)227

The effect of ammoniation on the intake and nutritive value of straw 4(1979)169

Utilization of alkali-treated grain. 2. Utilization by steers of diets based on hay or straw and mixed with either NaOH-treated or rolled barley 6(1981)347

Intake and digestibility in sheep fed on alkali-treated straw and concentrates, as complete diets or separate feeds 7(1982)271

The effect of prolonged storage on the digestibility and nitrogen content of ammonia-treated barley straw 8(1983)147

Methods of improving the utilization of cereal straw by ruminants. I. Supplements of urea, molasses and dried grass and treatment with sodium hydroxide 8(1983)221

Methods of improving the utilization of cereal straw by ruminants. II. Further studies on dried grass as a supplement for lambs 8 (1983)229

Methods of improving the utilization of cereal straw by ruminants. III. A note on the effect of ensiling straw treated with urea 9(1983)181

A note on the OECD collaborative experiment on ammonia-treated straw 10(1983/1984)89

Effect of defaunation on straw digestion in sacco and on particle retention in the rumen 10(1983/1984)165

Ammonia treatment of straw: methods for treatment and feeding experience in Norway 10(1983/1984)173

Ammonia versus sodium hydroxide treatment of straw for growing cattle 10(1983/1984) 189

Commercial experience of treating straw with ammonia 10(1983/1984)223

Anhydrous ammonia-treated cereal straw for animal feeding. Some results from the Mediterranean area 10(1983/1984)239

The value of urea-treated straw in diets of lactating goats 11(1984)67

Ammonia treatment of straw via the hydrolysis of urea. I. Effects of dry matter and urea concentrations on the rate of hydrolysis of urea 11(1984)103

Ammonia treatment of straw via the hydrolysis of urea. II. Additions of soya bean (urease), sodium hydroxide and molasses; effects on the digestibility of urea-treated straw 11(1984)115

The effect of milling and sodium hydroxide treatment on intake and digestibility of wheat straw by sheep 11(1984)149

Influence of heat sterilization and ammoniation on straw composition and degradation by pure cultures of cellulolytic rumen bacteria 12(1984/1985)195

The effect of supplementing grass silage with sodium hydroxide-treated chopped or milled straw on intake and digestibility by wether sheep 12(1984/1985)215

Methods of improving the utilization of cereal straw by ruminants. IV. The effect on intake of level of supplementation with urea and dried grass 13(1985)281

Methods of improving the utilization of cereal straw by ruminants. V. The effect of molasses and dried grass 13(1985)293

The use of barley straw as a feed 14(1986)29

Urea as a source of ammonia for improving the nutritive value of wheat straw 14(1986) 67

Alkaline hydrogen peroxide treatments of crop residues to increase ruminal dry matter disappearance in sacco 17(1987)179

Yield and potential feeding value of straws of grain sorghum cultivars developed in Nigeria 17(1987)285

The effect of ammoniation on the nutritive value of wheat, barley and oat straws. I. Changes in chemical composition in relation to digestibility in vitro and cell wall degradability 19(1988)159

The effect of ammoniation on the nutritive value of wheat, barley and oat straws. II. Digestibility and energy value measurements in vivo and their prediction from laboratory measurements 19(1988)173

The substitution of grass silage by a straw and concentrate mixture for dairy cows 20 (1988)135

The feeding value of ammoniated flax straw, wheat straw and wheat chaff for beef cattle 21(1988)57

Genetic variation in the feeding value of cereal straw 21(1988)127

Near infra-red spectra of ammonia-treated straw and of isolated cell walls 21(1988)209

Upgrading of straw by ammoniation 21(1988) 287

SO_2-treated straw as a silage additive: the participation of soluble and cell wall monosaccharide residues in lucerne silage fermentation 22(1988)255

Straw–grass

Titratable groups and soluble phenolic compounds as indicators of the digestibility of chemically treated roughages 6(1981)123

Straw–grass seed

Composition and enzymatic digestibility of oregon grass straws 9(1983)283

Straw–oat

Studies on ammonia-treated straw. I. The effects of type and level of ammonia, moisture content and treatment time on the digestibility in vitro and enzyme soluble organic matter of oat straw 7(1982)45

Botanical composition and extent of lignification affecting digestibility of wheat and oat straw and paspalum hay 7(1982)71

The effect of calcium hydroxide treatment on the digestibility of chopped oat straw supplemented with a solution containing urea, calcium, phosphorus, sodium, trace elements and vitamins 7(1982)93

The intake and digestibility by sheep of oat straw and maize stover offered with different levels of noug (*Guizotia abyssinica*) meal 16(1986/1987)99

Straw–rice

Effect of alkali-treated rice straw supplemented with spent tea leaf and thyroprotein on milk yield, milk composition and certain physiological parameters of dairy cows 7(1982)201

Supplementation of rice straw with crop by-products. I. Legume straw supplementation 19(1988)43

Nutritive value of rice straw (*Oryza sativa*) ensiled with animal excreta and rumen digesta 24(1989)69

Effect of urea treatment on rumen degradation characteristics of rice straws 24(1989) 83

Straw–sorghum

Effect of sorghum straw alone or in combination with molasses and nitrogen sources on performance of Sudan desert lambs 19 (1988)351

Straw–wheat

Performance of fattening cattle and lactating beef cows on diets high in chemically treated wheat straw and poultry litter 5 (1980)299

Ammonia and sodium hydroxide treatment of wheat straw in diets for fattening steers 7 (1982)1

Botanical composition and extent of lignification affecting digestibility of wheat and oat straw and paspalum hay 7(1982)71

Effect of treatment of wheat straw with ammonia and peracetic acid on digestibility in vitro and cell wall composition 7(1982)325

Digestibility of diets containing increasing levels of NaOH-treated or untreated wheat straw 7(1982)331

Ammoniation of wheat straw and native grass hay during baling of large round bales 8 (1983)35

Utilization of alkali-treated and neutralized wheat straw-based rations for growing goat kids 9(1983)155

Effect of wilting and the additives, straw, molasses and urea on the fermentation pattern of maize silage 9(1983)185

The response of wheat straw varieties to mild sulphur dioxide treatment 10(1983/1984) 269

Improving the intake and nutritive value of wheat straw for beef cows 12(1984/1985) 125

The utilization of untreated and calcium hydroxide treated wheat straw by sheep 12 (1984/1985)141

Nutritive evaluation of uromol–wheat straw as a basal ration for buffaloes 13(1985)103

Effects of ethanolamine and ethylenediamine treatments on the composition and degradability of wheat straw 13(1985)131

Chemical preparation of NaOH–keratin hydrolysate for improving the nutritive value of wheat straw 13(1985)215

Aqueous ammonia treatment of wheat straw: voluntary intake and digestibility in cattle 14(1986)81

Effects of supplementing a wheat straw diet with urea or a urea–molasses block and/or cottonseed meal on intake and liveweight change of lambs 16(1986/1987)25

Colonization of thick-walled plant tissues by anaerobic fungi 19(1988)25

SO_2 treatment for converting straw into a concentrate-like feed: a growth study with lambs 19(1988)219

Amino acids flow to and absorption from the small intestine of sheep fed barley and sulfur dioxide-treated straw at different ratios 22(1988)147

SO_2-treated straw as a silage additive: fermentation data on lucerne with particular reference to protein degradation 22(1988) 247

Effect of ammoniation and energy supplementation on the utilization of wheat straw by sheep 22(1988)305

Influence of stage of forage maturity and ammoniation of wheat straw on ruminal degradation of wheat forage tissues 24 (1989)201

Ensiling characteristics of cattle waste fermented with wheat straw and green berseem (*Trifolium alexandrium*) 25(1989)149

Digestion rates in and outflow rates from the rumen of sheep fed untreated or calcium hydroxide-treated wheat straw 25(1989) 179

Straw hydrolysates
Utilization of cattle manure for Torula yeast production from straw hydrolysates 5(1980) 175

Sucrase
Proximate chemical analysis of final cane molasses and effect of feeding 30% molasses on intestinal sucrase and maltase activities in the rat 3(1978)15

Sucrose
Energy metabolism in lambs fed silage diets with sucrose or a growth promoter 16 (1986/1987)203

The effect of dietary inclusion of starch, sucrose and xylose on the utilization of dietary energy in sheep 21(1988)11

Sugar beet
Sugar beet molasses for growing and fattening pigs 18(1987)131

Utilization of sugar beet molasses and monensin for finishing dairy bullocks 25(1989) 11

Sugar beet pulp
Dried suger beet pulp as a high energy feed for beef cattle 1(1976)643

Sugar beet pulp – a review 8(1983)1

Sugar beet pulp products for dairy cows 15 (1986)123

A note on the use of sugar-beet pulp silage and molasses in the diet of lactating dairy cows 20(1988)13

The effects of barley, unmolassed sugar-beet pulp and molasses supplements on organic matter, nitrogen and fibre digestion in the rumen of cattle given a silage diet 20(1988) 259

The effect of dietary inclusion of starch, sucrose and xylose on the utilization of dietary energy in sheep 21(1988)11

Digestion of diets containing molassed or plain sugar-beet pulp by growing pigs 23 (1989)67

Sugar beet pulp silage
Counteracting structure loss in pressed sugar beet pulp silage 24(1989)97

Sugarcane
Studies on the fermentation of chopped sugarcane 7(1982)411

The effects of ammonia on the fermentation of chopped sugarcane 9(1983)291

Effect of sugarcane condensed molasses solubles, sugarcane molasses and monensin on performance and volatile fatty acid production in finishing steers 12(1984/1985)275

Fermentation and utilization by lambs of sugarcane harvested fresh and ensiled with and without NaOH. 4. Ruminal kinetics 23 (1989)323

Sugarcane bagasse
Nutritive value for ruminants of sugar cane bagasse ensiled after spray treatment with different levels of NaOH 9(1983)1

Sulphur
Reducing the potential copper toxity of concentrates to sheep by the use of molybdenum and sulphur supplements 2(1977)235

Nutritive value of lupin-seed protein and leaf-protein concentrates supplemented with various sulphur sources 6(1981)189

The effects of sulphur supplementation on cellulose digestion in vitro and on nutrient digestion, nitrogen metabolism and rumen characteristics of lambs fed on good quality fescue and tropical star grass hays 8(1983) 129

Fractions of nitrogen, sulphur, phosphorus, calcium and magnesium in the herbage of perennial ryegrass as influenced by fertilizer nitrogen 14(1986)231

Sulphur dioxide
The response of wheat straw varieties to mild sulphur dioxide treatment 10(1983/1984) 269

Treatment of barley straw with ammonia and sulphur dioxide gases under laboratory conditions 14(1986)41

Evaluation of the increase in feeding value of barley straw following mild treatment with sulphur dioxide 15(1986)13

The effect of dietary barley on carbohydrate digestibility of sulphur dioxide-treated wheat straw by sheep 18(1987)55

Effects of ammonia and sulphur dioxide gases on the composition and digestion of barley straw 19(1988)121

Efficacy of sulfur dioxide as a grain preservative 19(1988)203

SO_2 treatment for converting straw into a concentrate-like feed: a growth study with lambs 19(1988)219

Pilot plant production of sulphur dioxide-treated mesquite wood. II. Digestibility and feedlot performance by lambs 22(1988) 61

Amino acids flow to and absorption from the small intestine of sheep fed barley and sulfur dioxide-treated straw at different ratios 22(1988)147

SO_2-treated straw as a silage additive: fermentation data on lucerne with particular reference to protein degradation 22(1988) 247

Sunflower cake
Feeding value of sunflower and groundnut cakes for broilers 4(1979)143

Feeding value of sunflower- and groundnut-cakes for laying hens 6(1981)63

Sunflower meal
Rumen degradation of protein in sunflower meal, rapeseed meal and lupin seed placed in nylon bags 11(1984)87

Sunflower seed
Inhibition of ruminal deamination in vitro by formaldehyde treatment of sunflower-seed, soya bean and fish meals: response curves to protective treatment 2(1977)267

Use of unextracted sunflower seeds as a protein source 5(1980)51

Effect of processing on the utilization by young pigs of the fat in soya beans and sunflower seeds 12(1984/1985)267

Effect of dietary protein and fat levels on the utilization of the fat in sunflower seeds by the young pig 13(1985)159

Utilization of whole rape (Canola) seed and sunflower seeds as sources of energy and protein in calf starter diets 15(1986)101

Supplementation–vine
Composition of vine-branches with leaves of *Vitis vinifera* L. fresh and ensiled using different additives (previous results) 16(1986/1987)89

Swedes
Swedes (*Brassica napus*) for growing pigs: chemical composition and use as a replacement for barley in the diet 2(1977)31

The effect of supplementary protein on the performance of pigs given swedes (*Brassica napus*) as a replacement for barley in the diet 2(1977)277

Sweet potato
Raw and steam-pelleted cassava, sweet potato and yam *cayenensis* as starch sources for ruminant and chicken diets 3(1978)73

Effects of ensiling or drying on five forage species in western Kenya: *Zea mays* (maize stover), *Pennisetum purpureum* (Pakistan napier grass), *Pennisetum* sp. (bana grass), *Impomea batata* (sweet potato vines) and *Cajanus cajan* (pigeon pea leaves) 13(1985)1

Sweetclover
Dicoumarol concentrations in sweetclover hay treated with preservatives and in spoiled hay of high- and low-coumarin cultivars of sweetclover 14(1986)221

T

Tall fescue
Effects of *Drechslera sorokiniana* infection on the yield and quality of tall fescue forage 15(1986)41

Tallow
Digestibility of a fully hydrogenated tallow for growing pigs 5(1980)87

Tamarind seeds
The nutritive value of tamarind seeds for broiler chicks 22(1988)285

Tannin
Rapeseed meal tannins and egg taint 6(1981)421

Tannins in sal seed (*Shorea robusta*) and sal seed meal limit their utilization as livestock feeds 7(1982)161

Proximate analysis, amino acid composition and tannin content of improved Nigerian sorghum varieties and their potential in poultry feeds 7(1982)359

Comparison of the nutritional value of low and medium tannin sorghum grains for pigs 8(1983)85

The influence of processing and storage on hydrogen cyanide and tannin contents of para-rubber seed and its products 9(1983)319

Waxy and normal grain sorghums with varying tannin contents in diets for young pigs 12(1984/1985)179

Utilization of tannin-containing and tannin-free faba beans (*Vicia faba*) by young chicks: effects of pelleting feeds on energy, protein and starch digestibility 20(1988)59

Treatment of low-tannin sorghum grain for broiler feed 20(1988)69

Effects of wood ash extract treatment on the feeding value and utilization of high-tannin sorghums by broiler chicks 22(1988)131

Tea
Spent tea leaf as a ruminant feed 3(1978)219

Effect of alkali-treated rice straw supplemented with spent tea leaf and thyroprotein on milk yield, milk composition and certain physiological parameters of dairy cows 7(1982)201

Rumen and post-rumen fermentation of spent tea leaf protein and other protein sources studied by the nylon bag method 7(1982)221

Techniques–amino acid evaluation
The determination of available lysine in barley using *Tetrahymena pyriformis* W. 5 (1980)33

A comparison of three methods for the determination of available lysine in barley 5 (1980)43

Estimation of optimum supplements of essential amino acids to a wheat and barley-based diet in rats using a nitrogen balance technique 15(1986)21

Techniques–ammonia treatment evaluation
Near infra-red spectra of ammonia-treated straw and of isolated cell walls 21(1988)209

Techniques–compound feed evaluation
The use of an enzymatic technique to predict digestibility, metabolizable and net energy of compound feedstuffs for ruminants 14 (1986)203

Techniques–delignification evaluation
Titratable groups and soluble phenolic compounds as indicators of the digestibility of chemically treated roughages 6(1981)123

Techniques–digestibility
Comparison of methods for predicting digestibility of feeds 20(1988)203

Techniques–digestibility evaluation
The analytical and biological accuracy of estimating the dry matter digestibility of different legume species 1(1976)61

Enzyme solubility – a method for evaluating the digestibility of alkali-treated straw 2 (1977)205

Prediction of the dry matter digestibility of tropical grasses using a pepsin-cellulase assay 2(1977)247

Principles governing the use of chemical methods for assessing the nutritive value of forages: a review 3(1978)265

Some remarks on the analytical procedure of Van Soest for the prediction of forage digestibility 3(1978)309

A note on Onozuka 3S cellulase as a replacement for Onozuka SS (P1500) cellulase when estimating forage digestibility in vitro 5(1980)347

Accuracy of predicting digestibility by the cellulase technique; the effect of pretreatment of forage samples with neutral detergent or acid pepsin 7(1982)83

Suitability of cell-wall constituents as predictors of organic matter digestibility in some tropical and subtropical by-products 7(1982)191

A new biological method for determining amino acid digestibility in poultry feedstuffs using a simple cannula, and the influence of dietary fibre on endogenous amino acid output 12(1984/1985)29

A comparison of laboratory methods involving single or two-stage cellulase and two-stage rumen inoculum–pepsin treatment for predicting digestibility and intake of hays in sheep 12(1984/1985)101

Improving the reliability of the two-stage in vitro assay for ruminant feed digestibility by calibration against in vivo data from a wide range of sources 18(1987)233

The use of a cellulase technique to predict digestibility, metabolizable and net energy of forages 19(1988)247

Comparison of methods for predicting digestibility of feeds 20(1988)203

A comparison of microbial, enzymatic, chemical and near-infrared reflectance spectroscopy methods in forage evaluation 20 (1988)219

Techniques–dry matter evaluation
Sources of error in predicting digestible dry matter from the acid-detergent fiber content of forages 21(1988)205

Techniques–dual flow continuous culture system
Evaluation of a dual flow continuous culture system for estimating bacterial fermentation in vivo of mixed diets containing various soya bean products 16(1986/1987)51

Techniques–dye absorption test
The dye absorption test as an indicator of the nutritive quality of heat-processed navy bean 9(1983)71

Techniques–energy availability evaluation
A comparison between fermentation heat of forages and organic matter digestibility determined by in vitro incubation with rumen fluid 23(1989)333

Techniques–fat and protein evaluation
Discriminating power of the nylon bag technique and the pepsin-cellulase method to detect less commonly used components included in concentrates for dairy cows 25 (1989)1

Techniques–fibre determination
A comparison of three fibre methods for predicting the metabolizable energy content of sorghum grain for poultry 2(1977)197

Some remarks on the determination of acid-detergent fibre in feeds with cetyltrimethylammonium bromide versus determination of detergent fibre with alkylarylsulfonic acid 6(1981)91

Techniques–forage digestibility prediction
Prediction of forage digestibility by principal component analysis of near infrared reflectance spectra 16(1986/1987)215

Techniques–monocarbonyl determination
Thermofractographic determination of monocarbonyl compounds in animal waste fats used as feed fat 13(1985)227

Techniques–near-infrared reflectance
Bias when predicting crude protein, dry matter digestibility and voluntary intake of tropical grasses by near-infrared reflectance 9(1983)221

Techniques–neutral detergent fibre evaluation
The study of several modifications of the neutral detergent fibre procedure 9(1983)19

Techniques–nutritive value
A new technique for upgrading the nutritive value of roughages under field conditions 3 (1978)201

Techniques–organic matter prediction
Comparison of laboratory methods for predicting the organic matter digestibility of forages 2(1977)337

Techniques–polysaccharide measurement
Classification and measurement of plant polysaccharides 23(1989)27

Techniques–preservative application
Laboratory application of preservatives to hay and the effects of irregular distribution on mould development 6(1981)73

Techniques–rumen microbiology
Possibilities of the use of recombinant-DNA techniques with rumen micro-organisms 21 (1988)161

Consecutive batch culture – a novel technique for the in vitro study of mixed microbial populations from the rumen 21(1988)183

Techniques–seed dehulling
A simple device for dehulling seeds and grain 3(1978)109

Techniques–silage assessment
A titration method for silage assessment 22 (1988)341

Techniques–straw spectroscopy
Near infrared reflectance spectroscopy of untreated and ammoniated barley straw 15 (1986)189

Tetrahymena pyriformis
The determination of available lysine in barley using *Tetrahymena pyriformis* W. 5 (1980)33

Thermal treatment
Nutritive value of thermoammoniated and steam-treated maize stover I. Intake, digestibility and nitrogen retention 4(1979) 177

Nutritive value of thermoammoniated and steam-treated maize stover II. Rumen metabolites and rate of passage 4(1979)187

The effect of different patterns of thermal processing of potatoes on their digestibility by growing pigs 4(1979)295

Investigations of the treatment of sawdust for rabbit feeding. 2. Effect of boiling and incubation with poultry manure 6(1981)51

The effect of steam-conditioning rate on the pelleting process 6(1981)83

The effect of some pre-ensiling treatments on silage composition and nitrogen disappearance in the rumen 8(1983)259

The influence of steam pelleting and grinding on the nutritive value of Canola rapeseed for poultry 8(1983)303

The dye absorption test as an indicator of the nutritive quality of heat-processed navy bean 9(1983)71

High temperature and pressure processing of maize cobs. I. Digestibility in vitro of processed cobs 12(1984/1985)151

Influence of heat sterilization and ammoniation on straw composition and degradation by pure cultures of cellulolytic rumen bacteria 12(1984/1985)195

Influence of supplemental fat and steam pelleting on the response of broilers to diets varying in energy density 13(1985)57

High temperature and pressure processing of maize cobs. II. Animal response and effect of sodium metabisulphite 14(1986)161

Thiaminase
Thiaminase activity in fish silage and moist fish feed 13(1985)39

Threonine
Evaluation of optimum lysine and threonine supplements to a wheat and barley-based diet in rats 8(1983)163

Thyroprotein
Effect of alkali-treated rice straw supplemented with spent tea leaf and thyroprotein on milk yield, milk composition and certain physiological parameters of dairy cows 7(1982)201

Timothy
Composition and fibre digestion in morphological components of an alfalfa–timothy sward 19(1988)135

Torula
Utilization of cattle manure for Torula yeast production from straw hydrolysates 5(1980) 175

The utilization of torula yeast in diets for White Leghorn birds during growth and laying periods 7(1982)185

Toxicology
Detoxification of aflatoxins in groundnut meal 2(1977)67

Long term feeding of concentrate fumigated with carbon tetrachloride to sheep 2(1977) 77

Reducing the potential copper toxity of concentrates to sheep by the use of molybdenum and sulphur supplements 2(1977)235

Utilization by sheep of dried pig faeces containing a high concentration of copper 3 (1978)31

Soil chemistry and mineral problems in farm livestock. A review 5(1980)95

Can lantana seeds be utilized in lamb rations? 9(1983)29

Variability for protein and oil quality in *Lupinus albus* 9(1983)271

The use of Balanites kernel cake in a diet for fattening sheep 9(1983)301

The influence of processing and storage on hydrogen cyanide and tannin contents of para-rubber seed and its products 9(1983) 319

Note on high-nitrate para grass (*Brachiaria mutica*) silage given to rabbits 9(1983)325

Effect of cassava-based diets varying in cyanide content on the performance and physiopathology of the African giant rat (*Cricetomys gambianus* Waterhouse) 11 (1984)1

Observations on the chemical and nutritive content of white winter and spring wheats contaminated with deoxynivalenol (vomitoxin) 11(1984)293

Hyperexcitability in cattle fed ammoniated roughages 17(1987)121

The effect of toxic substances present in some leaf protein concentrates on their nutritive value 24(1989)151

Canavanine content and toxicity of sesbania leaf meal for growing chicks 25(1989)137

In vitro metabolism of mycotoxins by bacterial, protozoal and ovine ruminal fluid preparations 25(1989)169

Trimethylamine oxide
The involvement of trimethylamine oxide in fish meal in the production of egg taint 8 (1983)119

Triticale
A nutritional evaluation of tritical 9(1983)49

Nutritive value of three cultivars for sheep, pigs and poultry 18(1

Evaluation of triticale given to pi and rats 19(1988)359

Effects of enzyme supplementati based on wheat, rye or tritica productive value for broiler c (1988)313

Nutritional value of triticale (C broiler diets 22(1988)119

Nutritive value of diets containi and varying mixtures of triticale for growing-finishing swine 22(1

The effect of fat level and type lization of triticale (cultivar C broiler chicks 22(1988)273

Effects of ozonation on the chem sition and in vitro degradation internodes and red clover sten 159

Trypsin
Trypsin inhibitor, mineral avail performance of broiler chickens based on rice bran 4(1979)133

Nutrient and trypsin inhibitor co droponically sprouted soya bea 203

Tryptophan
Meat meal in the diet of the early- II. Amino acid supplementatio

Meat meal in the diet of the e pig. IV. The supplementatic with tryptophane, lysine and n (1982)27

Tryptophan content of feeds, li diets and requirement for grow (1985)171

Utilisation of L- and DL-amin broiler chicks fed diets contain 17(1987)1

Turkey
Turkey feeding regime with diffe introduction of a choice of foods

Limiting amino acids in dried mi given to young turkeys 4(1979)

A note on the nutritive value stant coffee residue for broiler turkey poults 10(1983/1984)28

Effects of pelleting, added sodiu and fat in a wheat-based die mance and carcass characteris white turkeys 12(1984/1985)2

Torula
Utilization of cattle manure for Torula yeast production from straw hydrolysates 5(1980) 175

The utilization of torula yeast in diets for White Leghorn birds during growth and laying periods 7(1982)185

Toxicology
Detoxification of aflatoxins in groundnut meal 2(1977)67

Long term feeding of concentrate fumigated with carbon tetrachloride to sheep 2(1977) 77

Reducing the potential copper toxity of concentrates to sheep by the use of molybdenum and sulphur supplements 2(1977)235

Utilization by sheep of dried pig faeces containing a high concentration of copper 3 (1978)31

Soil chemistry and mineral problems in farm livestock. A review 5(1980)95

Can lantana seeds be utilized in lamb rations? 9(1983)29

Variability for protein and oil quality in *Lupinus albus* 9(1983)271

The use of Balanites kernel cake in a diet for fattening sheep 9(1983)301

The influence of processing and storage on hydrogen cyanide and tannin contents of para-rubber seed and its products 9(1983) 319

Note on high-nitrate para grass (*Brachiaria mutica*) silage given to rabbits 9(1983)325

Effect of cassava-based diets varying in cyanide content on the performance and physiopathology of the African giant rat (*Cricetomys gambianus* Waterhouse) 11 (1984)1

Observations on the chemical and nutritive content of white winter and spring wheats contaminated with deoxynivalenol (vomitoxin) 11(1984)293

Hyperexcitability in cattle fed ammoniated roughages 17(1987)121

The effect of toxic substances present in some leaf protein concentrates on their nutritive value 24(1989)151

Canavanine content and toxicity of sesbania leaf meal for growing chicks 25(1989)137

In vitro metabolism of mycotoxins by bacterial, protozoal and ovine ruminal fluid preparations 25(1989)169

Trimethylamine oxide
The involvement of trimethylamine oxide in fish meal in the production of egg taint 8 (1983)119

Triticale
A nutritional evaluation of triticale with pigs 9(1983)49

Nutritive value of three cultivars of triticale for sheep, pigs and poultry 18(1987)19

Evaluation of triticale given to pigs, poultry and rats 19(1988)359

Effects of enzyme supplementation of diets based on wheat, rye or triticale on their productive value for broiler chickens 20 (1988)313

Nutritional value of triticale (Carman) for broiler diets 22(1988)119

Nutritive value of diets containing triticale and varying mixtures of triticale and maize for growing-finishing swine 22(1988)217

The effect of fat level and type on the utilization of triticale (cultivar Carman) by broiler chicks 22(1988)273

Effects of ozonation on the chemical composition and in vitro degradation of triticale internodes and red clover stems 24(1989) 159

Trypsin
Trypsin inhibitor, mineral availability, and performance of broiler chickens fed on diets based on rice bran 4(1979)133

Nutrient and trypsin inhibitor content of hydroponically sprouted soya beans 13(1985) 203

Tryptophan
Meat meal in the diet of the early-weaned pig. II. Amino acid supplementation 4(1979)43

Meat meal in the diet of the early-weaned pig. IV. The supplementation of diets with tryptophane, lysine and methionine 7 (1982)27

Tryptophan content of feeds, limitations in diets and requirement for growing pigs 13 (1985)171

Utilisation of L- and DL-amino acids by broiler chicks fed diets containing Pruteen 17(1987)1

Turkey
Turkey feeding regimes with different ages of introduction of a choice of foods 4(1979)125

Limiting amino acids in dried microbial cells given to young turkeys 4(1979)209

A note on the nutritive value of dried instant coffee residue for broiler chickens and turkey poults 10(1983/1984)285

Effects of pelleting, added sodium bentonite and fat in a wheat-based diet on performance and carcass characteristics of small white turkeys 12(1984/1985)223

The relative availability of phosphorus in inorganic feed phosphates for young turkeys and pigs 13(1985)117

Influence of nutrient density on feed consumption, weight gain and gut capacity of broilers, Leghorns and turkeys reared to 26 days of age 16(1986/1987)129

U

Urea

The use of whole barley diets fortified with solutions of urea, minerals and vitamins for lambs 2(1977)307

Studies on fermentation of sorghum silage during storage - effect of urea and molasses 3(1978)299

Performance of heifers fed on diets based on NaOH-treated maize cobs and the effect of supplementary urea and source of carbohydrates 4(1979)97

Evaluation of corn gluten meal with urea as a source of supplementary nitrogen for growing calves and lambs 7(1982)375

Methods of improving the utilization of cereal straw by ruminants. I. Supplements of urea, molasses and dried grass and treatment with sodium hydroxide 8(1983)221

The effect of urea on the preservation and digestibility in vitro of perennial ryegrass 10(1983/1984)49

The value of urea-treated straw in diets of lactating goats 11(1984)67

Ammonia treatment of straw via the hydrolysis of urea. I. Effects of dry matter and urea concentrations on the rate of hydrolysis of urea 11(1984)103

Effects of molasses/urea and bacterial inoculant additives on silage quality, dry matter recovery, and feeding value for cattle 12 (1984/1985)205

Effect of urea on the ensiling process of orange pulp 12(1984/1985)233

Effects of supplementing a wheat straw diet with urea or a urea–molasses block and/or cottonseed meal on intake and liveweight change of lambs 16(1986/1987)25

Effect of formaldehyde-treated soya bean meal and urea in starters on nitrogen quality, degradibility in sacco, sheep digestibility and calf performance 16(1986/1987)287

Complexes of urea and formaldehyde as non-protein nitrogen compounds in ruminant rations: a review 20(1988)1

Nitrogen utilization, ruminal fermentation and abomasal nitrogen flow in sheep fed caucasian bluestem hay supplemented with lucerne or urea 20(1988)33

Preservation and storage of green panic (*Panicum maximum*) as moist hay with urea 20 (1988)87

The effect of treatment with urea, sorbic acid, or dehydration on orange peel silage 20 (1988)335

Effects of added dietary urea on the utilization of maize stover silage by growing beef cattle 22(1988)227

Urease

The effect of urease enzyme on treatment time and the nutritive value of straw treated with ammonia as urea 8(1983)271

Rumen fermentation, urease activity and performance of cattle given palm kernel cake-based diet 20(1988)79

Uromol

Nutritive evaluation of uromol–wheat straw as a basal ration for buffaloes 13(1985)103

V

Vinasse

Protein and energy value of vinasse for pigs 5 (1980)221

Vine

Composition of vine-branches with leaves of *Vitis vinifera* L. fresh and ensiled using different additives (previous results) 16(1986/1987)89

Virginiamycin

Efficacy of virginiamycin and a commercially-available lactobacillus probiotic in swine diets 8(1983)69

Vitamin C

Effect of dietary ascorbic acid (vitamin C) supplementation on the performance of broiler chickens in a tropical environment 16(1986/1987)17

Vitamin K

Comparison of the relative biological availability and stability of commercial sources of vitamin K activity 3(1978)329

Vitamins, general

The use of whole barley diets fortified with solutions of urea, minerals and vitamins for lambs 2(1977)307

W

Water supply

Lucerne juice as a protein supplement for growing pigs: effects of mineral content of the diet and of the water supply 6(1981)35

Weeds

Utilization of aquatic plants *Elodea canadensis* and *Hydrilla verticillata* in broiler chick diets 20(1988)155

Weende components

Apparent and true digestibility of the Weende components, cell content and cell wall of ryegrass 1(1976)607

Wheat

Nutritive value of leaf protein coagulum and cereals mixtures 4(1979)91

Nutritional evaluation, using rats, of diets based on cereals and ethanol-grown yeast 6(1981)309

Evaluation of optimum lysine and threonine supplements to a wheat and barley-based diet in rats 8(1983)163

The effect of calcium and sodium hydroxides and of sodium acrylate on the fermentation and digestibility in vitro of ensiled whole-crop wheat and barley harvested at different stages of maturity 9(1983)37

Effects of milling and pelleting of maize, barley and wheat on their metabolizable energy value for cockerels and chicks 9(1983) 99

The effect of milling and sodium hydroxide treatment on intake and digestibility of wheat straw by sheep 11(1984)149

Observations on the chemical and nutritive content of white winter and spring wheats contaminated with deoxynivalenol (vomitoxin) 11(1984)293

Chemical composition and nutritive value of Canadian winter wheats (*Triticum aestivum* L.) given to laboratory rats 12(1984/1985)75

Effects of pelleting, added sodium bentonite and fat in a wheat-based diet on performance and carcass characteristics of small white turkeys 12(1984/1985)223

Estimation of optimum supplements of essential amino acids to a wheat and barley-based diet in rats using a nitrogen balance technique 15(1986)21

Feeding value of sprouted wheat (*Triticum aestivum*) for beef cattle finishing diets 15 (1986)113

Treatment of whole-crop cereals with alkali. 1. The influence of sodium hydroxide and ensiling on the chemical composition and in vitro digestibility of rye, barley and wheat crops harvested at increasing maturity and dry matter content 18(1987)257

Treatment of whole-crop cereals with alkali. 2. Voluntary intake and digestibility by sheep of rye, barley and wheat crops ensiled with sodium hydroxide 18(1987)271

Treatment of whole-crop cereals with alkali. 3. Voluntary intake and digestibility studies in sheep given immature wheat ensiled with sodium hydroxide, urea or ammonia 18(1987)283

Effects of enzyme supplementation of diets based on wheat, rye or triticale on their productive value for broiler chickens 20 (1988)313

A comparison of wheat- or barley-based diets given ad libitum as meal or pellets to finishing pigs 22(1988)263

Effect of soft white wheat addition to alfalfa-grass forage on heifer gain, diet digestibility and in vitro digestion kinetics 24(1989)141

Factors effecting the digestibility by pigs of energy and protein in wheat, barley and sorghum diets supplemented with canola meal 24(1989)253

Effect of formaldehyde treatment on the rumen digestion of barley and wheat grain and its influence on hay intake and digestion by sheep 25(1989)55

The effect of β-glucanase supplementation on nutrient digestibility and growth in broilers given diets containing barley, oat groats or wheat 25(1989)193

Wheat grain

Turkey feeding regimes with different ages of introduction of a choice of foods 4(1979)125

Wheat silage

Wheat silage for fattening young male cattle 12(1984/1985)253

Treatment of whole-crop cereals with alkali. 4. Voluntary intake and growth in steers given wheat ensiled with sodium hydroxide, urea or ammonia 19(1988)55

Wheat straw

Performance of fattening cattle and lactating beef cows on diets high in chemically treated wheat straw and poultry litter 5 (1980)299

Ammonia and sodium hydroxide treatment of wheat straw in diets for fattening steers 7 (1982)1

Botanical composition and extent of lignification affecting digestibility of wheat and oat straw and paspalum hay 7(1982)71

Effect of treatment of wheat straw with ammonia and peracetic acid on digestibility in vitro and cell wall composition 7(1982)325

Digestibility of diets containing increasing levels of NaOH-treated or untreated wheat straw 7(1982)331

Ammoniation of wheat straw and native grass hay during baling of large round bales 8 (1983)35

Utilization of alkali-treated and neutralized wheat straw-based rations for growing goat kids 9(1983)155

Effect of wilting and the additives, straw, molasses and urea on the fermentation pattern of maize silage 9(1983)185

The response of wheat straw varieties to mild sulphur dioxide treatment 10(1983/1984) 269

Improving the intake and nutritive value of wheat straw for beef cows 12(1984/1985) 125

The utilization of untreated and calcium hydroxide treated wheat straw by sheep 12 (1984/1985)141

Nutritive evaluation of uromol–wheat straw as a basal ration for buffaloes 13(1985)103

Effects of ethanolamine and ethylenediamine treatments on the composition and degradability of wheat straw 13(1985)131

Chemical preparation of NaOH–keratin hydrolysate for improving the nutritive value of wheat straw 13(1985)215

Aqueous ammonia treatment of wheat straw: voluntary intake and digestibility in cattle 14(1986)81

Effects of supplementing a wheat straw diet with urea or a urea–molasses block and/or cottonseed meal on intake and liveweight change of lambs 16(1986/1987)25

Colonization of thick-walled plant tissues by anaerobic fungi 19(1988)25

The effect of ammoniation on the nutritive value of wheat, barley and oat straws. I. Changes in chemical composition in relation to digestibility in vitro and cell wall degradability 19(1988)159

The effect of ammoniation on the nutritive value of wheat, barley and oat straws. II. Digestibility and energy value measurements in vivo and their prediction from laboratory measurements 19(1988)173

SO_2 treatment for converting straw into a concentrate-like feed: a growth study with lambs 19(1988)219

The effect of variation in the water content of wheat straw before ammoniation on its nutritive value for beef cattle 20(1988)111

Upgrading of straw by ammoniation 21(1988) 287

Amino acids flow to and absorption from the small intestine of sheep fed barley and sulfur dioxide-treated straw at different ratios 22(1988)147

SO_2-treated straw as a silage additive: fermentation data on lucerne with particular reference to protein degradation 22(1988) 247

SO_2-treated straw as a silage additive: the participation of soluble and cell wall monosaccharide residues in lucerne silage fermentation 22(1988)255

Effect of ammoniation and energy supplementation on the utilization of wheat straw by sheep 22(1988)305

Influence of stage of forage maturity and ammoniation of wheat straw on ruminal degradation of wheat forage tissues 24 (1989)201

Ensiling characteristics of cattle waste fermented with wheat straw and green berseem (*Trifolium alexandrium*) 25(1989)149

Digestion rates in and outflow rates from the rumen of sheep fed untreated or calcium hydroxide-treated wheat straw 25(1989) 179

Whey

The replacement of skim milk by dried microbial cells and whey in milk diets for calves 2(1977)85

The nutritive value of liquid whey, either sour or sweet, when given in restricted amounts to the growing pig 3(1978)163

The evaluation of fresh sweet liquid whey as a protein supplement to maize for growing pigs 14(1986)171

Observations on the feeding of non-lactating sows on graded levels of liquid whey 24 (1989)327

Wilting

Effect of prolonged wilting in poor conditions on the fermentation quality, metabolisability and net energy value of silage given to sheep 12(1984/1985)109

Influence of processing methods and storage time on the cyanide potential of cassava leaf meal 17(1987)227

Wood ash extract

Effects of wood ash extract treatment on the feeding value and utilization of high-tannin sorghums by broiler chicks 22(1988)131

X

Xanthophylls
Investigations into the effects of processing on the retention of the carotenoid fractions of *Leucaena leucocephala* during storage, and the effects of processing on mimosine concentration 9(1983)307

Xylose
The effect of dietary inclusion of starch, sucrose and xylose on the utilization of dietary energy in sheep 21(1988)11

Y

Yam
Raw and steam-pelleted cassava, sweet potato and yam *cayenensis* as starch sources for ruminant and chicken diets 3(1978)73

Yeast
Utilization of yeast grown on $_n$-paraffins in the feeding of heavy pigs, from weaning to slaughter 1(1976)25

Nutritional evaluation, using rats, of diets based on cereals and ethanol-grown yeast 6(1981)309

The utilization of torula yeast in diets for White Leghorn birds during growth and laying periods 7(1982)185

The effect of pelleting on the utilisation of yeast grown on n-alkanes by broiler chicks 12(1984/1985)285

Yellow turnip
Studies on the nutritive value of roots, cabbage and grass silage for growing commercial rabbits 13(1985)299

Z

Zinc
The effect of age of regrowth on content and release of manganese, iron, zinc and copper from four tropical forages incubated in sacco in rumen of sheep 20(1988)171